

Regular Article -Theoretical Physics

t-channel dark matter models – a whitepaper

Chiara Arina , Benjamin Fuks^{2,a}, Luca Panizzi^{3,4}, Michael J. Baker⁵, Alan S. Cornell⁶, Jan Heisig⁷, Benedikt Maier⁸, Rute Pedro⁹, Dominique Trischuk¹⁰, Diyar Agin², Alexandre Arbey¹¹, Giorgio Arcadi^{12,13}, Emanuele Bagnaschi¹⁴, Kehang Bai¹⁵, Disha Bhatia¹⁶, Mathias Becker^{17,18,19}, Alexander Belyaev^{20,21}, Ferdinand Benoit², Monika Blanke^{22,23}, Jackson Burzynski²⁴, Jonathan M. Butterworth²⁵, Antimo Cagnotta²⁶, Lorenzo Calibbi²⁷, Linda M. Carpenter²⁸, Xabier Cid Vidal²⁹, Emanuele Copello¹⁷, Louie Corpe³⁰, Francesco D'Eramo^{18,19}, Aldo Deandrea^{6,11}, Alaman Desai³¹, Caterina Doglioni³², Sunil M. Dogra³³, Mathias Garny³⁴, Mark D. Goodsell², Sohaib Hassan³⁵, Philip Coleman Harris³⁶, Julia Harz¹⁷, Alejandro Ibarra³⁴, Alberto Orso Maria Iorio^{37,38}, Felix Kahlhoefer²², Deepak Kar^{39,40}, Shaaban Khalil⁴¹, Valery Khoze⁴², Pyungwon Ko⁴³, Sabine Kraml⁴⁴, Greg Landsberg⁴⁵, Andre Lessa⁴⁶, Laura Lopez-Honorez^{47,48}, Alberto Mariotti^{48,49}, Vasiliki A. Mitsou⁵⁰, Kirtimaan Mohan⁵¹, Chang-Seong Moon³³, Alexander Moreno Briceno⁵², Maria Moreno Llacer⁵⁰, Leandre Munoz-Aillaud², Taylor Murphy^{2,53}, Anele M. Ncube⁶, Wandile Nzuza³⁹, Clarisse Prat³⁹, Lena Rathmann⁷, Tait⁵⁷, Andrea Thamm⁵, Michel H. G. Tytgat⁴⁷, Zirui Wang⁵⁸, David Yu⁵⁹, Shin-Shan Yu⁶⁰

- ⁴ INFN-Cosenza, 87036 Arcavacata di Rende, Cosenza, Italy
- ⁵ Department of Physics, University of Massachusetts, Amherst, MA 01003, USA
- ⁶ Department of Physics, University of Johannesburg, PO Box 524, Auckland Park 2006, South Africa
- ⁷ Institute for Theoretical Particle Physics and Cosmology, RWTH Aachen University, Sommerfeldstr. 16, 52074 Aachen, Germany
- ⁸ Imperial College, London SW7 2AZ, UK
- ⁹ Laboratório de Instrumentação e Física Experimental de Partículas, Lisbon, Portugal
- ¹⁰ Department of Physics, Brandeis University, Waltham, MA, USA
- ¹¹ Université Claude Bernard Lyon 1, CNRS/IN2P3, IP2I UMR 5822, 4 rue Enrico Fermi, 69100 Villeurbanne, France
- ¹² Dipartimento di Scienze Matematiche e Informatiche, Scienze Fisiche e Scienze della Terra, Universita degli Studi di Messina, Via Ferdinando Stagno d'Alcontres 31, 98166 Messina, Italy
- ¹³ INFN Sezione di Catania, Via Santa Sofia 64, 95123 Catania, Italy
- ¹⁴ INFN, Laboratori Nazionali di Frascati, Via E. Fermi 40, 00044 Frascati, RM, Italy
- ¹⁵ University of Oregon, 1585 E 13th Ave, Eugene, OR 97403, USA
- ¹⁶ Instituto de Física, Universidade de São Paulo, São Paulo, Brazil
- ¹⁷ PRISMA+ Cluster of Excellence & Mainz Institute for Theoretical Physics, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany
- ¹⁸ Dipartimento di Fisica e Astronomia, Università degli Studi di Padova, Via Marzolo 8, 35131 Padua, Italy
- ¹⁹ INFN, Sezione di Padova, Via Marzolo 8, 35131 Padua, Italy
- ²⁰ School of Physics and Astronomy, University of Southampton, Southampton SO17 1BJ, UK
- ²¹ Particle Physics Department, Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, UK
- ²² Institut für Astroteilchenphysik, Karlsruhe Institute of Technology, Hermann-von-Helmholtz-Platz 1, 76344 Eggenstein-Leopoldshafen, Germany
- ²³ Institut für Theoretische Teilchenphysik, Karlsruhe Institute of Technology, Engesserstraße 7, 76128 Karlsruhe, Germany
- ²⁴ Simon Fraser University, 8888 University Dr W, Burnaby V5A 1S6, Canada
- ²⁵ Department of Physics and Astronomy, University College London, London, UK
- ²⁶ Texas A&M University, College Station, USA

Published online: 12 September 2025

- ²⁷ School of Physics, Nankai University, Tianjin 300071, China
- ²⁸ Department of Physics, The Ohio State University, Columbus, OH 43210, USA
- ²⁹ Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain
- ³⁰ Université Clermont Auvergne, CNRS/IN2P3, LPCA, 63000 Clermont-Ferrand, France
- ³¹ Department of Physics, The University of Adelaide, Adelaide, SA 5005, Australia



¹ Centre for Cosmology, Particle Physics and Phenomenology (CP3), Université catholique de Louvain, 1348 Louvain-la-Neuve, Belgium

² Laboratoire de Physique Théorique et Hautes Énergies (LPTHE), UMR 7589, Sorbonne Université & CNRS, 4 place Jussieu, 75252 Paris Cedex 05, France

³ Dipartimento di Fisica, Università della Calabria, 87036 Arcavacata di Rende, Cosenza, Italy

975 Page 2 of 96 Eur. Phys. J. C (2025) 85:975

- ³² School of Physics and Astronomy, University of Manchester, Manchester, UK
- ³³ Kyungpook National University, Daegu, South Korea
- ³⁴ Department of Physics, TUM School of Natural Sciences, Technical University of Munich, James-Franck-Str. 1, 85748 Garching, Germany
- ³⁵ University of Bergen, Bergen, Norway
- ³⁶ Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139-4307, USA
- ³⁷ Dipartimento di Fisica E. Pancini, Universitá di Napoli Federico II, Complesso Universitario di Monte Sant'Angelo, Via Cintia, Naples, Italy
- ³⁸ INFN, Sezione di Napoli, Naples, Italy
- ³⁹ School of Physics, University of Witwatersrand, Johannesburg, South Africa
- ⁴⁰ Royal Society Wolfson Visiting Fellow at the University of Glasgow, Glasgow, UK
- ⁴¹ Center for Fundamental Physics, Zewail City of Science and Technology, 6th of October City, Giza 12578, Egypt
- ⁴² IPPP, Durham University, Durham, UK
- ⁴³ School of Physics, Korea Institute for Advanced Study (KIAS), 85 Hoegi-ro, Dongdaemun-gu, Seoul 02455, Republic of Korea
- ⁴⁴ Laboratoire de Physique Subatomique et de Cosmologie (LPSC), Université Grenoble-Alpes, CNRS/IN2P3, 53 Avenue des Martyrs, 38026 Grenoble, France
- ⁴⁵ Department of Physics, Brown University, 182 Hope St., Providence, RI 02912, USA
- ⁴⁶ Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, Santo André, SP 09210-580, Brazil
- ⁴⁷ Service de Physique Théorique, C.P. 225, Université Libre de Bruxelles, Boulevard du Triomphe, 1050 Brussels, Belgium
- ⁴⁸ Theoretische Natuurkunde & The International Solvay Institutes, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium
- ⁴⁹ IIHE/ELEM, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium
- ⁵⁰ Instituto de Física Corpuscular (IFIC), CSIC-Universitat de València, C/ Catedrático José Beltrán 2, 46980 Paterna, Spain
- ⁵¹ Department of Physics and Astronomy, Michigan State University, 567 Wilson Road, East Lansing, MI 48824, USA
- ⁵² Universidad Antonio Nariño, Bogota, Colombia
- ⁵³ Department of Physics, Miami University, 500 E. Spring St., Oxford, OH 45056, USA
- ⁵⁴ Department of Physics, University of Cape Town, Cape Town, South Africa
- 55 Sydney Consortium for Particle Physics and Cosmology, School of Physics, The University of New South Wales, Sydney, NSW 2052, Australia
- ⁵⁶ Department of Physics and Astronomy, Sam Houston State University, Huntsville, TX, USA
- ⁵⁷ University of California, Irvine, USA
- ⁵⁸ Fudan University, Shanghai 200443, People's Republic of China
- ⁵⁹ Department of Physics, State University of New York at Buffalo, Buffalo, NY 14260, USA
- ⁶⁰ Department of Physics, The Catholic University of America, Washington, DC 20064, USA

Received: 23 April 2025 / Accepted: 9 August 2025 © The Author(s) 2025, corrected publication 2025

Abstract This report, summarising work achieved in the context of the LHC Dark Matter Working Group, investigates the phenomenology of *t*-channel dark matter models, spanning minimal setups with a single dark matter candidate and mediator to more complex constructions closer to UV-complete models. For each considered class of models, we examine collider, cosmological and astrophysical implications. In addition, we explore scenarios with either promptly decaying or long-lived particles, as well as featuring diverse dark matter production mechanisms in the early universe. By providing a unified analysis framework, numerical tools and guidelines, this work aims to support future experimental and theoretical efforts in exploring *t*-channel dark matter models at colliders and in cosmology.

Contents

- 2 Benchmark models for LHC phenomenology

	2.3	Dark matter simplified models inspired by com-						
		positeness						
	2.4	Towards UV completions – frustrated dark matter. New gauge interactions to connect the dark sec-						
	2.5							
		tor to the Standard Model						
3	annel dark matter at the LHC							
	3.1	Minimal simplified models – prompt decays						
		3.1.1 Generalities about the signal of quark-						
		philic dark matter						
		3.1.2 First generation simplified models						
		3.1.3 Second generation simplified models						
		3.1.4 Third generation simplified models						
		3.1.5 Simplified models: considerations on sig-						
		nal modelling						
		3.1.6 Leptophilic models						
	3.2	Non-minimal models – prompt decays						
		3.2.1 Flavoured dark matter at the LHC						
		3.2.2 Composite dark matter at the LHC						
		3.2.3 Frustrated dark matter at the LHC						
		3.2.4 A fermionic portal to a non-Abelian dark						
		sector						
	3.3	Long-lived t-channel mediators						

2.2 Flavoured dark matter: dark minimal flavour



^a e-mail: fuks@lpthe.jussieu.fr (corresponding author)

Eur. Phys. J. C (2025) 85:975 Page 3 of 96 975

3.3.1 Generalities						
3.3.2 Current LLP searches						
3.3.3 Coverage of current searches						
3.3.4 Gaps in coverage						
3.3.5 Advancing reinterpretation methods						
4 Cosmology						
4.1 Generalities						
4.2 Dark matter relic density						
4.2.1 Canonical freeze-out						
4.2.2 Conversion-driven freeze-out						
4.2.3 Freeze-in						
4.2.4 SuperWIMP						
4.2.5 Non-perturbative effects						
4.2.6 Altered cosmological histories						
4.3 Dark matter constraints						
4.3.1 Indirect detection						
4.3.2 Direct detection						
4.3.3 Early universe constraints						
4.4 Cosmological constraints for benchmark models						
4.4.1 Minimal simplified models						
4.4.2 Non-minimal models						
5 Conclusion						
Appendix A: Implementation of the ATLAS-EXOT-						
2018-06 search in MadAnalysis5						
1. Description of the analysis						
2. Validation of the implementation						
References						
	•					

1 Introduction

Despite compelling evidence for its existence [1–4], dark matter (DM) remains elusive in direct searches using nuclear and electronic recoil experiments, indirect probes via cosmicray and gamma-ray spectra analyses, and collider searches for missing transverse energy signatures. Understanding the nature of dark matter is therefore among the top priorities in particle physics, astrophysics and cosmology. Currently, observational evidence confirms the existence of a gravitationally-interacting form of matter that constitutes approximately 25% of the Universe's energy budget. However, the fundamental properties of DM such as its particle nature, its interactions with Standard Model (SM) particles if any, its mass and its spin, are still unknown despite decades of intense efforts. From a theoretical perspective, a vast array of models predicting the existence of DM has been developed over the years. The null results from direct, indirect, and collider DM search experiments have imposed increasingly stringent constraints on these models, reducing the viable regions in the associated parameter spaces.

These bounds are usually derived through two complementary approaches. First, a model-specific approach in which dark matter is embedded within a comprehensive theo-

retical framework built from first principles could be considered. Here, despite current data and related constraints, viable scenarios span orders of magnitude in terms of dark matter masses and couplings. The goal of this report is not to provide an exhaustive overview of all theoretical aspects of these DM models, for which excellent reviews are already available (see for instance the recent work [4] and references therein). Alternatively, experimental results could be interpreted following a phenomenological approach based on simplified models which capture essential features of diverse classes of SM extensions while remaining agnostic to the underlying high-energy theory. Simplified models are particularly useful as they allow for a systematic and model-independent exploration of viable DM scenarios, facilitating the comparison of theoretical predictions with experimental data from varied experiments and guiding the design of future experiments. In this report, we thus adopt the simplified model paradigm [5,6], generically assuming the existence of a DM particle that interacts minimally with the SM through a limited number of well-defined couplings and additional new physics states.

In the most simplified models for dark matter, the dark sector is assumed to consist of a single massive DM particle that interacts with the Standard Model via a single new 'mediator' particle. In addition, the stability of the DM candidate is ensured by imposing a discrete or continuous symmetry under which the DM state transforms differently from the SM states. The interactions of the new physics sector are then determined by the quantum numbers of the mediator and its representations under both the new symmetry and the SM gauge group. In simplified s-channel DM model configurations (with thus a single DM state and a single mediator state), introduced about a decade ago [7–13], the mediator transforms under the new symmetry like the states from the SM sector. In contrast, in the *t*-channel models central to this report, the mediator's properties relative to the new symmetry are similar to those of the DM particle. For instance, under a discrete \mathbb{Z}_2 symmetry, an s-channel setup might involve a \mathbb{Z}_2 -odd DM particle, with the SM and mediator states being \mathbb{Z}_2 -even. In this case, the mediator couples both to a pair of SM particles and to a pair of DM particles.

In a t-channel scenario, however, the mediator would be \mathbb{Z}_2 -odd and thus couple to one DM particle and one SM particle. Furthermore, in the simplest cases, t-channel mediators must decay into DM and SM particles on timescales shorter than cosmological ones, as otherwise, they would appear stable and subsequently violate observational constraints as carrying colour or electric charge. While specific incarnations of minimal t-channel models have been largely explored in the past, the corresponding studies often focused on particular representations and spin configurations for the new states. For instance, this class of model was initially introduced in the context of Majorana or Dirac dark matter



975 Page 4 of 96 Eur. Phys. J. C (2025) 85:975

coupling to quarks [14–19]. Complementarily, some studies also embedded the minimal *t*-channel construction sketched above within a UV-complete framework like supersymmetry.

One of the primary goals of this report is to provide guidance for collider searches targeting final states with missing transverse energy originating from t-channel DM scenarios. This includes not only the minimal framework described above but also non-minimal setups that bring the models closer to UV-complete theories, and that incorporate additional features not captured by the minimal models that could be highly relevant for collider searches. We classify the considered t-channel models according to their degree of minimality. We begin with the most simplified frameworks, extending the SM by a single \mathbb{Z}_2 -odd DM particle and a single \mathbb{Z}_2 -odd mediator. Within this setup, we allow for different spin configurations and interactions with the SM, considering all possible mediator-SM couplings involving quarks and leptons (but one at a time). This approach builds on previous studies [20–25], where efforts were made to simultaneously simplify the new physics parameter space and systematically explore variations in the quantum numbers of the new fields. Next, we extend our analysis to more complex models that involve multiple mediators and DM candidates, while still relying on t-channel exchanges for DM annihilation. These models include scenarios with dark sector flavour symmetries [26], constructions inspired by compositeness [27–29] and frustrated DM frameworks [30]. Another approach to non-minimality is to retain a minimal field content while allowing the fields to transform under non-trivial representations of either the SM gauge group, or an extended gauge group. Examples include the so-called Minimal DM model of [31] or the Minimal Consistent DM scenarios of [32]. In this report, we focus on the latter and explore a setup featuring a non-minimal gauged dark sector [33]. Within all these models, a wide range of DM coupling values leads to cosmologically viable scenarios, which correspond to varied possibilities for the mediator decay width. We analyse cases where mediators decay promptly into DM and SM particles, as well as scenarios with small mediator decay widths, resulting in long-lived particle (LLP) signatures at the LHC. The latter case is particularly motivated by DM production mechanisms in the early universe beyond the standard freezeout paradigm, such as conversion-driven freeze-out [34,35], freeze-in [36,37], or superWIMP production [38,39].

In all these scenarios, we discuss the implications for collider searches, cosmology and astrophysics. By leveraging the complementarity of these observations, we aim to provide a comprehensive framework for probing the parameter spaces of a wide range of t-channel models in the near future. To this end, we additionally provide numerical tools, including those necessary for high-precision Monte Carlo simulations for LHC studies, along with detailed instructions on how to use them. Furthermore, simulated Monte Carlo sam-

ples for simplified models can also be obtained upon request, in the form of grids of mediator and dark matter masses, in order to facilitate reinterpretation and signal modelling. We hope that these efforts will subsequently enable a systematic reinterpretation of both existing and future experimental results in terms of more complex DM models.

This report, that summarizes work achieved in the framework of the LHC Dark Matter working Group, is structured as follows. In the first part (Sect. 2), we provide a comprehensive overview of the t-channel dark matter models under consideration. This includes simplified scenarios discussed in Sect. 2.1, as well as more complex and non-minimal theories explored in Sect. 2.2 (flavoured dark matter), 2.3 (composite dark matter), 2.4 (frustrated dark matter) and 2.5 (non-Abelian gauged dark sectors). Subsequently, we address the phenomenology of these models, focusing on collider studies in Sect. 3 and cosmological implications in Sect. 4. The collider phenomenology is discussed separately for scenarios involving promptly decaying mediators and those with long-lived mediators, while similarly, the cosmological phenomenology distinguishes between canonical dark matter freeze-out and alternative DM production scenarios, such as conversion-driven freeze-out, the superWIMP mechanism, and freeze-in production. We conclude with a summary of our findings and perspectives for future studies in Sect. 5.

2 Benchmark models for LHC phenomenology

In this section, we introduce various classes of models to be explored in this whitepaper. We begin with minimal setups featuring *t*-channel dark matter (DM), that we discuss in Sect. 2.1. Additionally, we delve into non-minimal models incorporating diverse phenomenological features not addressed in the minimal case. These encompass flavoured dark matter models, composite realisations, frustrated dark matter scenarios and non-Abelian gauged dark sectors, which are detailed in Sects. 2.2, 2.3, 2.4 and 2.5 respectively.

2.1 Minimal options and their implementation in high-energy physics software

In a minimal realisation of a t-channel simplified model for dark matter, the field content of the Standard Model (SM) is extended to include one DM candidate X, assumed to be a colourless electroweak singlet, and one mediator state Y. Additionally, to ensure DM stability, an $ad\ hoc\ \mathbb{Z}_2$ symmetry is imposed such that all SM fields are even while the dark matter and the mediator are odd. Following the DMSimpt framework introduced in [40], no assumptions are made regarding the spin of the DM and the mediator, the flavour structure of their interactions, and the quantum numbers of the mediator that depend on how it couples dark matter to the SM sector.



Eur. Phys. J. C (2025) 85:975 Page 5 of 96 975

Under these conditions, the model presents three possible spin configurations for the X particle: it could be a scalar (either the complex state S or the real state \tilde{S}), a fermion (either the Dirac fermion χ or the Majorana fermion $\tilde{\chi}$), or a vector (either the complex state V or the real state \tilde{V}). For bosonic dark matter, the mediator is then a fermionic object, that we denote as ψ , whereas for fermionic dark matter, the mediator is a scalar field φ . We leave the case of vector mediators to future work, as achieving next-to-leading order QCD simulations in models with coloured vector states is far from trivial, even with existing automated tools such as those discussed in this report.

Expanding upon the setup introduced in [40], the full Lagrangian incorporating the interactions of these fields with the SM can be expressed as follows:

$$\mathcal{L} = \mathcal{L}_{SM} + \mathcal{L}_{kin} + \mathcal{L}_{XY}. \tag{1}$$

Here, \mathcal{L}_{SM} represents the SM Lagrangian and \mathcal{L}_{kin} encompasses gauge-invariant kinetic and mass terms for all new fields. The final term \mathcal{L}_{XY} incorporates the interactions between the mediator and the DM with the SM. This term involves a significant number of free coupling parameters, particularly in the absence of assumptions regarding the flavour structure of the couplings. Separating the six possibilities for the properties of the dark matter state, the Lagrangian \mathcal{L}_{XY} takes the following form:

$$\begin{split} \mathcal{L}_{XY}^{\text{F3S}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f f_R \, \tilde{S} + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} Q_L \, \tilde{S} + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F1S}} &= \lambda_{\ell} \, \bar{\psi}_{\ell} \ell_R \, \tilde{S} + \lambda_L \, \bar{\psi}_L L_L \, \tilde{S} + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3C}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f f_R \, S + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} Q_L \, S + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F1C}} &= \lambda_{\ell} \, \bar{\psi}_{\ell} \ell_R \, S + \lambda_L \, \bar{\psi}_L L_L \, S + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{S3M}} &= \sum_{f=u,d} \lambda_f \, \overline{\tilde{\chi}} f_R \, \varphi_f^\dagger + \lambda_{\mathcal{Q}} \, \overline{\tilde{\chi}} Q_L \, \varphi_{\mathcal{Q}}^\dagger + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{S1M}} &= \lambda_{\ell} \, \overline{\tilde{\chi}} \ell_R \, \varphi_\ell^\dagger + \lambda_L \, \overline{\tilde{\chi}} L_L \, \varphi_L^\dagger + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{S3D}} &= \sum_{f=u,d} \lambda_f \, \overline{\chi} f_R \, \varphi_f^\dagger + \lambda_{\mathcal{Q}} \, \overline{\chi} Q_L \, \varphi_{\mathcal{Q}}^\dagger + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{S3D}} &= \sum_{f=u,d} \lambda_f \, \overline{\chi} \ell_R \, \varphi_\ell^\dagger + \lambda_L \, \overline{\chi} L_L \, \varphi_L^\dagger + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3V}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R \, \tilde{V}_\mu + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} \gamma^\mu Q_L \, \tilde{V}_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \lambda_{\ell} \, \bar{\psi}_{\ell} \gamma^\mu \ell_R \, \tilde{V}_\mu + \lambda_L \, \bar{\psi}_L \gamma^\mu L_L \, \tilde{V}_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R \, V_\mu + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} \gamma^\mu Q_L \, V_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R \, V_\mu + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} \gamma^\mu Q_L \, V_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R \, V_\mu + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} \gamma^\mu Q_L \, V_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R \, V_\mu + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} \gamma^\mu Q_L \, V_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R \, V_\mu + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} \gamma^\mu Q_L \, V_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R \, V_\mu + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} \gamma^\mu Q_L \, V_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R \, V_\mu + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} \gamma^\mu Q_L \, V_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R \, V_\mu + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} \gamma^\mu Q_L \, V_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R \, V_\mu + \lambda_{\mathcal{Q}} \, \bar{\psi}_{\mathcal{Q}} \gamma^\mu Q_L \, V_\mu + \text{H.c.}, \\ \mathcal{L}_{XY}^{\text{F3W}} &= \sum_{f=u,d} \lambda_f \, \bar{\psi}_f \gamma^\mu f_R$$

In these Lagrangians, the symbols F3S, S3M, F3V (F1S, S1M, F1V) refer to models with real scalar, fermionic and vector dark matter through the symbol endings S, M and

 $\mathcal{L}_{YY}^{\text{F1W}} = \lambda_{\ell} \, \bar{\psi}_{\ell} \gamma^{\mu} \ell_{R} \, V_{\mu} + \lambda_{L} \, \bar{\psi}_{L} \gamma^{\mu} L_{L} \, V_{\mu} + \text{H.c.}$

v, while the beginning of the symbol names F3 and S3 (F1 and S1) refer to colour-triplet (colour-singlet) fermionic and scalar mediators. Similarly, the designations F3C, S3D and F3W (F1C, S1D and F1W) are utilised for the complex dark matter cases through the ending C, D and W. In our notation, Q_L and L_L represent the weak doublets of left-handed SM quarks and leptons, respectively, while u_R , d_R and ℓ_R denote the corresponding weak singlets. The fermionic mediators ψ_Q , ψ_L , ψ_u , ψ_d and ψ_ℓ interact exclusively with the Q_L , L_L , u_R , d_R and ℓ_R fields, respectively. Furthermore, we introduce one fermionic mediator for each generation of SM fermions. Consequently, they lie in the same representation as their SM fermionic partners, and they are vector-like. Similarly, the scalar mediators φ_Q , φ_L , φ_u , φ_d and φ_ℓ only interact with Q_L , L_L , u_R , d_R and ℓ_R respectively.

The scalar and fermionic mediators all possess a flavour index. Therefore, while in the Lagrangians of Eq. (2) all flavour indices are omitted for clarity, it is important to note that the coupling matrices λ_Q , λ_L , λ_u , λ_d and λ_ℓ are 3×3 matrices in the flavour space. Unless stated otherwise, these matrices are considered real and flavour-diagonal to prevent loop-induced mixing between the SM fermions. Each mediator is hence associated with a given generation of SM partners.

Most of the results presented in this whitepaper are obtained through the joint usage of various standard highenergy physics packages, following the tool chain outlined in [40]. This process involved implementing a *single* model file that incorporates simultaneously all Lagrangians described above in FeynRules [41,42]¹, which we utilised alongside Mogre [43], NLOCT [44] and FeynArts [45] to generate a next-to-leading order (NLO) UFO [46,47] model with five flavours of massless quarks. The resulting UFO model, termed the DMSimpt 2.0 model, is an extension of the implementation designed in [40], now incorporating leptonic fields. This implementation is available from the FeynRules model database (see https://feynrules.irmp.ucl. ac.be/wiki/DMsimpt or the GitHub repository https://github. com/BFuks/DMSimpt). Like the previous implementation, the available NLO UFO model includes λ couplings that are flavour-diagonal and real. However, the FeynRules model allows for a more general structure, which we relied on to generate a leading-order (LO) UFO model which is used for some searches currently conducted by the ATLAS and CMS collaborations. While an NLO extension of this more general model is feasible, it necessitates a more intricate renormalisation procedure to be implemented within Mogre. We leave this task to future work.

¹ Specifically, we use the feynrules-dev-bsm branch of FeynRules available from the GitHub repository (https://github.com/FeynRules/FeynRules).



975 Page 6 of 96 Eur. Phys. J. C (2025) 85:975

Table 1 Model information for the F3S and F3C models. We provide the name of the DM particle and the mediators used in the FeynRules implementation, together with the associated PDG identifiers, spin quantum number and representation under $SU(3)_c \times SU(2)_L \times U(1)_Y$. We recall that three generations of mediators are included, which

requires sets of three PDG identifiers. We additionally provide the new physics couplings linking the DM, the mediators and the SM sector. Each coupling is given together with the associated FeynRules symbol and the reference Les Houches block to be used in the parameter card

DM	Field Š S	FeynRules Xs Xc	PDG 51 56	Spin 0 0	Repr. (1, 1, 0) (1, 1, 0)
Mediators	Field	FeynRules	PDG	Spin	Repr.
	$\psi_{\scriptscriptstyle \mathcal{Q}} = egin{pmatrix} \psi_{\scriptscriptstyle \mathcal{Q}}^{(u)} \ \psi_{\scriptscriptstyle \mathcal{Q}}^{(d)} \end{pmatrix}$	$YF3Q = \begin{pmatrix} YF3Qu \\ YF3Qd \end{pmatrix}$	5910002 5910004 5910006 5910001 5910003 5910005	1/2	$(3, 2, \frac{1}{6})$
	ψ_u	YF3u	5920002 5920004 5920006	1/2	$(3, 1, \frac{2}{3})$
	ψ_d	YF3d	5920001 5920003 5920005	1/2	$(3, 1, -\frac{1}{3})$
	$\psi_{\scriptscriptstyle L} = egin{pmatrix} \psi_{\scriptscriptstyle L}^{(u)} \ \psi_{\scriptscriptstyle L}^{(\ell)} \end{pmatrix}$	$\mathtt{YF1L} = \begin{pmatrix} \mathtt{YF1Lv} \\ \mathtt{YF1Le} \end{pmatrix}$	5910012 5910014 5910016 5910011 5910013 5910015	1/2	$(1, 2, -\frac{1}{2})$
	ψ_ℓ	YF1e	5920011 5920013 5920015	1/2	(1, 1, -1)
Couplings	Parameter	FeynRules	Les Houches Block		
	$\lambda_{\mathcal{Q}}$	lamF3Q	DMF3Q		
	λ_u	lamF3u	DMF3U		
	λ_d	lamF3d	DMF3D		
	λ_L	lamF1L	DMF1L		
	λ_ℓ	lamF1e	DMF1E		

We present information on the six simplified models in Tables 1, 2 and 3 for models with scalar, fermionic and vector DM respectively. For each model, we list the new included fields, their representation under the SM gauge group, their spins, the particle names used in the FeynRules implementation, and the Particle Data Group (PDG) identifiers [48]. Additionally, we detail the conventions for the coupling parameters, including their implementation names and the corresponding Les Houches blocks [49] where numerical values are stored.

The generated UFO model can be used within the MG5aMC platform [50] for achieving computations relevant for to collider phenomenology. Both LO and NLO simulations are feasible for models with a flavour-diagonal structure. However, for models with off-diagonal couplings in the flavour space, only LO simulations are achievable. Additionally, we have produced LO model files in which all flavours of quarks are massive. They are available both in UFO and Calchep [51] formats, enabling their use with micromegas [52,53] and MadDM [54–56] for evaluating cosmological observables. The implemented FeynRules model and the corresponding generated UFO libraries encompass all possibilities for the DM candidate, thus incorporating all Lagrangians of Eq. (2). In order to reduce the number of free parameters compared to the general case, we include specific restrictions tailored for minimal t-channel DM simplified models. One particular restriction involves the activation of either a single mediator or a set of mass-degenerate mediators. Consequently, all new physics states are decoupled and non-interacting, except for a single DM candidate and specific mediators.

We first consider 'universal' possibilities (referred to as XYZ_uni with XYZ being one of the six symbols introduced in Eq. (2) to represent the various DM scenarios), where all mediators of the model are active but assumed to be degenerate, with interaction strengths that are both flavour-conserving and universal. The coupling parameters of such a scenario therefore satisfy the condition:

$$(\lambda_Q)_{ij} = (\lambda_L)_{ij} = (\lambda_u)_{ij} = (\lambda_d)_{ij} = (\lambda_\ell)_{ij} = \lambda \, \delta_{ij}, \tag{3}$$

where λ represents the sole, universal, free coupling parameters and i, j = 1, 2, 3 are flavour indices. Alongside the universal mediator mass (M_Y) and the DM mass (M_X) , the model is defined by three new physics parameters. Furthermore, we can construct a second class of universal scenarios in the context of leptophilic dark matter. These scenarios, termed XYZ_1R models, exclusively involve DM couplings to all three SM right-handed leptons, generalising the framework studied in [57] to accommodate various spin configurations. In this case, the coupling parameters are defined by

$$(\lambda_{Q})_{ij} = (\lambda_{L})_{ij} = (\lambda_{u})_{ij} = (\lambda_{d})_{ij} = 0$$

and
$$(\lambda_{\ell})_{ij} = \lambda \, \delta_{ij},$$
 (4)



Eur. Phys. J. C (2025) 85:975 Page 7 of 96 975

Table 2 Same as in Table 1 but for the S3M and S3D models

DM	Field $ ilde{\chi}$ χ	FeynRules Xm Xd	PDG 52 57	Spin 1/2 1/2	Repr. (1, 1, 0) (1, 1, 0)
Mediators	Field	FeynRules	PDG	Spin	Repr.
	$\varphi_{\mathcal{Q}} = \begin{pmatrix} \varphi_{\mathcal{Q}}^{(u)} \\ \varphi_{\mathcal{Q}}^{(d)} \end{pmatrix}$	$YS3Q = \begin{pmatrix} YS3Qu \\ YS3Qd \end{pmatrix}$	1000002 1000004 1000006 1000001 1000003 1000005	0	$(3, 2, \frac{1}{6})$
	$arphi_u$	YS3u	2000002 2000004 2000006	0	$(3, 1, \frac{2}{3})$
	$arphi_d$	YS3d	2000001 2000003 2000005	0	$(3, 1, -\frac{1}{3})$
	$arphi_L = egin{pmatrix} arphi_L^{(u)} \ arphi_L^{(\ell)} \end{pmatrix}$	$YS1L = \begin{pmatrix} YS1Lv \\ YS1Le \end{pmatrix}$	1000012 1000014 1000016 1000011 1000013 1000015	0	$(1, 2, -\frac{1}{2})$
	$arphi_\ell$	YS1e	2000011 2000013 2000015	0	(1, 1, -1)
Couplings	Parameter	FeynRules	Les Houches Block		
	$\lambda_{\mathcal{Q}}$	lamS3Q	DMS3Q		
	λ_u	lamS3u	DMS3U		
	λ_d	lamS3d	DMS3D		
	λ_L	lamS1L	DMS1L		
	λ_ℓ	lamS1e	DMS1E		

Table 3 Same as in Table 1 but for the F3V and F3W models

DM	Field <i>V</i> <i>V</i>	FeynRules Xv Xw	PDG 53 58	Spin 1 1	Repr. (1, 1, 0) (1, 1, 0)
Mediators	Field	FeynRules	PDG	Spin	Repr.
	$\psi_{\mathcal{Q}} = egin{pmatrix} \psi_{\mathcal{Q}}^{(u)} \ \psi_{\mathcal{Q}}^{(d)} \end{pmatrix}$	$YF3Q = \begin{pmatrix} YF3Qu \\ YF3Qd \end{pmatrix}$	5910002 5910004 5910006 5910001 5910003 5910005	1/2	$(3, 2, \frac{1}{6})$
	ψ_u	YF3u	5920002 5920004 5920006	1/2	$(3, 1, \frac{2}{3})$
	ψ_d	YF3d	5920001 5920003 5920005	1/2	$(3, 1, -\frac{1}{3})$
	$\psi_{\scriptscriptstyle L} = egin{pmatrix} \psi_{\scriptscriptstyle L}^{(u)} \ \psi_{\scriptscriptstyle L}^{(\ell)} \end{pmatrix}$	$\mathtt{YF1L} = \begin{pmatrix} \mathtt{YF1Lv} \\ \mathtt{YF1Le} \end{pmatrix}$	5910012 5910014 5910016 5910011 5910013 5910015	1/2	$(1, 2, -\frac{1}{2})$
	ψ_ℓ	YF1e	5920011 5920013 5920015	1/2	(1, 1, -1)
Couplings	Parameter	FeynRules	Les Houches Block		
	$\lambda_{\mathcal{Q}}$	lamF3Q	DMF3Q		
	λ_u	lamF3u	DMF3U		
	λ_d	lamF3d	DMF3D		
	λ_L	lamF1L	DMF1L		
	λ_ℓ	lamF1e	DMF1E		

and the model is again defined by three free parameters, the DM and (universal) mediator masses M_X and M_Y , as well as the coupling parameter λ . Such a class of models are however not examined in this report. In all these universal setups, the presence of several mediators associated with a specific generation of SM fermion prevents strong constraints from flavour-changing-neutral-current processes, both in the quark and lepton sectors. This contrasts to models featuring mediators coupling to several generations of fermions.

Next, we focus on models in which a single class of mediator is considered. One such scenario is exemplified by the XYZ_uR possibilities investigated in [24,25], wherein the mediator couples exclusively to the right-handed up quark. In this setup, the coupling parameters are constrained to

$$(\lambda_u)_{11} = \lambda$$
 and $(\lambda_f)_{ij} = 0$ for all other couplings
(with $f = Q, L, u, d, \ell$). (5)

In contrast to the universal case, λ represents the coupling strength between the mediator and the right-handed up quark,



975 Page 8 of 96 Eur. Phys. J. C (2025) 85:975

and this restricted scenario is characterised by three free parameters: the masses of the dark matter and the mediator M_X and M_Y , and the coupling λ . In the present work, we will examine scenarios in which the new particles couple to any flavour of right-handed quark (referred to as XYZ_dR, XYZ_cR, XYZ_sR, XYZ_bR and XYZ_tR models) and the righthanded muon (referred to as XYZ_muR models), motivated by the specificities inherent to each generation. In particular, we keep in mind the special role that the second generation could play with respect to new physics (to solve, for instance, the anomalous magnetic moment of the muon puzzle or some of the remaining flavour anomalies), as well as that of the third generation due the heavier corresponding fermion masses and their possible connection to the electroweak symmetry breaking mechanism. In contrast, we only outline in our cosmological analysis the complementarity of astroparticle searches to LHC-based ones, relying on previous studies of models related to the first and third generations of quarks. The reasons is that in this case, the flavour of the quark is not relevant for the constraints that can be imposed on the model so that a smooth extrapolation of the results for models relevant to first-generation quarks can be applied to models relevant to second-generation and bottom quarks.

2.2 Flavoured dark matter: dark minimal flavour violation

While minimal models serve as a valuable parametrisation for a broad range of t-channel DM scenarios, they often fail to provide a comprehensive description of non-minimal features. Therefore, it is crucial to explore models beyond the minimal setups, to ensure thorough theoretical and experimental investigations without overlooking potential loopholes.

One way to expand upon the minimal framework outlined in Sect. 2.1 involves introducing multiple flavours of DM (i.e. a field X_i carrying a flavour index i) and assuming that they transform under a certain flavour symmetry. Early studies of such flavoured DM models [58–65] have applied the concept of minimal flavour violation (MFV) [66–71] to the dark sector. In MFV models, an approximate flavour symmetry $U(3)^5 \equiv U(3)_O \times U(3)_u \times U(3)_d \times U(3)_L \times U(3)_e$ is imposed, treating the five SM fundamental fermion fields Q_L , u_R , d_R , L_L , and ℓ_R as flavour triplets whose components rotate into each other via five separate U(3) groups. This flavour symmetry is then broken by Yukawa interactions. In minimally flavour-violating dark matter models, the DM field $X \equiv (X_1, X_2, X_3)$ is assumed to transform as a triplet under one of these U(3) symmetries. The coupling matrix λ is then determined following the MFV assumption, and its structure follows from an expansion in terms of the SM Yukawa couplings. Additionally, X is required to be a complex field. Unlike the simplified model discussed in Sect. 2.1, the mediator field in flavoured DM models does not carry a flavour index. However, in less minimal 'skew-flavoured' constructions [72], both the DM and the mediator carry flavour indices.

Models featuring dark minimal flavour violation (DMFV) [26] extend this concept further. The flavour symmetry group is promoted to $U(3)^6$ (or $U(3)^5 \times O(3)$ for real DM fields), with additional symmetry transformations acting on the DM field $X(\chi)$ in the notation of [26]), rendering it a flavour triplet. For models with a single mediator, we introduce a t-channel mediator $Y(\phi)$ in the notation of [26]) that couples the dark flavour triplet X to one of the representations of SM fermions f. The Lagrangian governing the interaction between the dark matter and the mediator is generically expressed as

$$\mathcal{L}_{XY} = \lambda_{ij} \,\bar{f}_i X_j Y,\tag{6}$$

where i and j represent flavour indices. Adhering to the DMFV principle, the complex 3×3 coupling matrix λ serves as the sole source of flavour symmetry breaking apart from the Standard Model Yukawa couplings. In contrast to the Lagrangians introduced in Eq. (2), the second flavour index of the coupling λ_{ij} is here associated with the dark matter, rather than the mediator for which only a single state is considered. In accordance with the DMFV principle, the mass matrix M_X of the DM states is not an arbitrary parameter in the theory, but is instead determined by a spurion expansion based on the flavour-violating coupling λ . For example, for X being a Dirac fermion it reads

$$M_X = m_X \Big[\mathbb{1} + \eta \lambda^{\dagger} \lambda + O(\lambda^4) \Big], \tag{7}$$

where η is a real expansion parameter. This relationship ties the dark mass spectrum directly to the flavour structure of the coupling λ that connects the visible and dark sectors. Moreover, the approximate U(3) flavour symmetry in the dark sector reduces the complexity of the coupling matrix λ that can, for complex DM, be characterised by three diagonal coupling strengths, three mixing angles, and three complex phases. It is worth noting that in quark-flavoured DMFV models (i.e. models in which X and Y couple to quarks) featuring Dirac DM, the stability of the lightest DM flavour is ensured by a residual \mathbb{Z}_3 symmetry, which persists after the flavour symmetry is broken.

This has motivated previous studies that have investigated the phenomenology of DMFV models with Dirac dark matter coupling to a specific quark flavour [26,73–75]. Scenarios with right-handed up, down, charm and bottom quarks have hence been explored, as well as setups in which the DM state predominantly couples to the doublet of left-handed top and bottom quarks. This corresponds to $f_i = u_R$, d_R , c_R , b_R and $(t_L, b_L)^T$ in Eq. (6). Furthermore, the possibility of Majorana DM has been examined for the case of $f = u_R$ [76,77]. Here, the presence of a smaller dark flavour symmetry group O(3) results in a coupling matrix λ containing more free



Eur. Phys. J. C (2025) 85:975 Page 9 of 96 975

parameters. Additionally, the absence of the DM-stabilising \mathbb{Z}_3 symmetry necessitates the introduction of an *ad hoc* \mathbb{Z}_2 symmetry to ensure DM stability.

The correspondence between the general DMFV parametrisation discussed above and the simplified model parametrisation outline in Sect. 2.1 can be established in two cases: either when all physical dark matter states are nearly mass-degenerate, or when one DM state is much lighter than the others. In such scenarios, the DMFV model effectively comprises one dark matter state X and one mediator state Y featuring couplings to all generation of SM fermions. Consequently, the coupling matrices λ appearing in Eq. (2) reduce to vectors in the flavour space and only one mediator state is active. Various DMFV-inspired simplified models can then be constructed, depending on the SM fermion representation f to which the DM state couples, and the particle nature of the new physics fields X and Y. While the DMSimpt 2.0 UFO model incorporates such a flexibility, we employ dedicated LO UFO libraries available from the DMSimpt GitHub repository https://github.com/lena-ra/ Flavored-Dark-Matter.

Non-minimally flavoured dark matter can also be introduced to couple to the lepton sector [78–80], as studied in the context of Dirac and complex scalar DM coupling to right-handed charged leptons. In these constructions, a discrete \mathbb{Z}_2 symmetry is always necessary to stabilise dark matter. Additionally, less minimal models [81] featuring two mediators have been studied. Such a setup can provide an explanation to the longstanding deviations between theoretical predictions and experimental measurements relevant to the anomalous magnetic moment of the muon. While the matrices of couplings λ_1 and λ_2 associated with the two mediators are related by DMFV, the DM spectrum becomes instead free. Finally, flavoured DM models beyond DMFV have also been found to offer a promising avenue for achieving successful baryogenesis [82].

2.3 Dark matter simplified models inspired by compositeness

Another compelling possibility for exploring t-channel dark matter models beyond minimal frameworks emerges from composite scenarios of new physics [83–85]. These models typically postulate the existence of additional coloured and non-coloured resonances, arising as bound states from an underlying fermionic construction beyond the SM featuring a new strong dynamics. While some of these resonances are closely tied to the top quark, elucidating the question of its large mass via mechanisms like partial compositeness [86], others could be stable, electrically neutral, and colourless, thereby potentially serving as candidates for dark matter [27–29]. In such frameworks, DM stability is ensured by a residual discrete \mathbb{Z}_2 symmetry stemming from the breaking of the

new strong dynamics. The DM state is assigned an odd parity under this new \mathbb{Z}_2 symmetry, while all SM fields are assigned an even parity. Meanwhile, the particle spectrum comprises both even and odd new states.

An example of next-to-minimal setup, just slightly more complex than the simplified models introduced in Sect. 2.1, involves the inclusion of one real scalar DM state together with two fermionic top partners, or mediators as per the terminology used in this manuscript. Specifically, this setup encompasses one \mathbb{Z}_2 -even state denoted as Y_t' (or T' in the notation of [28]), and one \mathbb{Z}_2 -odd state labelled as Y_t (or T' in the notation of [28]). Both these states share the same quantum numbers as the SM right-handed top quark field t_R , and their interactions with the real scalar DM state X (or S in the notation of [28]) can be captured within an effective Lagrangian resembling those introduced in Eq. (2). This Lagrangian can be expressed as

$$\mathcal{L}_{XY} = \lambda_t \, \bar{Y}_t t_R \, X + \lambda_t' \, \bar{Y}_t Y_t' \, X + \text{H.c.}$$
 (8)

Here, the parameters λ_t and λ_t' represent two new physics couplings, that were noted \tilde{y}_t and $\tilde{y}_{T'}$ in the notation of [28]. The list of free parameters additionally includes the dark matter mass M_X and the two mediator masses M_Y and $M_{Y'}$. The FeynRules model and associated UFO libraries are available from a dedicated GitHub repository, located at https://github.com/BFuks/CompositeDM.

A less minimal effective DM model, still inspired by composite constructions but closer to UV completions compared the simplified setup of Sect. 2.1, could entail a sufficient number of mediator fields to generate the top quark mass via partial compositeness. In such a scenario, both weak singlet and doublet mediators are necessary [87]. Assuming all these mediators to be \mathbb{Z}_2 -odd for simplicity, an effective interaction Lagrangian incorporating their coupling to dark matter would be given by

$$\mathcal{L}_{XY} = \lambda_t \, \bar{Y}_t t_R \, X + \lambda_Q \, \bar{Y}_Q Q_L \, X + \text{H.c.} \quad \text{with}$$

$$Y_Q = \begin{pmatrix} Y_{Q,t} \\ Y_{Q,b} \end{pmatrix} \quad \text{and} \quad Q_L = \begin{pmatrix} t_L \\ b_L \end{pmatrix}. \tag{9}$$

This model introduces a certain number of free parameters, including two new physics couplings λ_Q and λ_t , one dark matter mass M_X , and three mediator masses M_{Y_t} , $M_{Y_{Q,t}}$ and $M_{Y_{Q,b}}$. Although these masses are predictable in effective setups incorporating partial compositeness, for simplicity we treat them as free parameters. Consequently, such a configuration can be mapped to the DMSimpt 2.0 simplified model, which is provided with an associated restriction termed F3S -VLQ, recalling that the choice of scalar DM and fermionic mediators is the relevant one for composite constructions.



975 Page 10 of 96 Eur. Phys. J. C (2025) 85:975

2.4 Towards UV completions – frustrated dark matter

Simple UV-complete dark matter models, which avoid the theoretical shortcomings or arbitrary motivations of certain simplified models, can often be integrated into larger and well-motivated theoretical frameworks [32]. Consequently, they could serve as non-minimal simplified models for dark matter that offer the advantage of a more complex phenomenology including a variety of astrophysical and collider signatures. Frustrated dark matter (fDM) models [30] exemplify these simple UV-complete setups. They consist of models where interactions between dark matter and the visible sector arise at one-loop order, rather than at tree level. Such loop-level couplings can emerge in many UV-complete theories, and they yield signatures that are highly sensitive to the details of the mediator-SM interactions. The fDM framework encompasses a broad class of models where the mediators carry SM charges, but the interactions of dark matter are 'frustrated' in the sense that the specific mediator assignments preclude tree-level interactions with the SM.

In this framework, as is customary, the dark matter field X is assumed to transform as a singlet under the SM gauge symmetry. However, the SM gauge charges of all mediator fields coupling both to X and to the Standard Model are chosen to forbid renormalisable gauge-invariant contact interactions between X and any SM fermion. For fermionic dark matter, these models necessitate a pair of mediators to couple the dark state X to the SM. A schematic representation of this family of models, featuring SM-singlet Dirac dark matter coupled to a pair of mediators, takes the form:

$$SM \longleftrightarrow mediators [\varphi, \psi] \longleftrightarrow DM [X],$$
 (10)

where, as in Sect. 2.1, φ denotes a scalar mediator and ψ a fermionic one. Since the mediators carry SM charges, they can generally decay into SM particles which has the extra benefit to circumvent cosmological and phenomenological issues associated with new stable particles. Therefore, one or both mediators should have renormalisable interactions with the SM, hence allowing for mediator decays.

Within the scope of this whitepaper, we focus on fDM models where mediators carry $SU(3)_C$ quantum numbers, as particles with non-trivial colour charges are anticipated to exhibit the highest production cross sections at the LHC. Depending on the details of the mediator sector, these mediators may also carry non-trivial $SU(2)_L$ charges and $U(1)_Y$ hypercharges. Among the numerous possible $SU(3)_C$ charge assignments for the mediating sector, only certain choices permit direct renormalisable interaction between the mediators and the SM. For instance, only the colour triplet, sextet, and octet options allow for renormalisable interactions between mediators and pairs of quarks. Notably, colour-octet

mediators can enjoy such couplings without the necessity of non-trivial electroweak charges.

We consider a particular fDM realisation where the mediators are $SU(3)_C$ sextets and weak singlets, this last condition being imposed for simplicity, and where the DM is a Dirac fermion. In this scenario, only the scalar mediator directly couples to the SM at tree level, specifically to a pair of quarks of possibly the same electric charge. The hypercharge Y of the mediators depends on the structure of these quark-messenger couplings. While a comprehensive examination of all possible low-dimensional couplings is provided in [88], only one of them is renormalisable. This interaction couples the scalar sextet to a uu, ud, or dd quark pair pursuant to its hypercharge, and its phenomenological consequences have been explored in [89–91]. For simplicity, we only consider a sextet scalar with Y = 4/3 so that the corresponding Lagrangian \mathcal{L}_{XY}^2 is given by

$$\mathcal{L}_{XY} = \lambda_X \varphi^{\dagger} \bar{X} \psi + \lambda_{qq} \bar{K}_6 \varphi^{\dagger} \bar{u}_R^c u_R + \text{H.c.}, \tag{11}$$

where all indices have been omitted for clarity. Here, λ_X represents the mediator-DM coupling, and λ_{qq} is the mediator-SM coupling matrix in the flavour space. Additionally, \bar{K}_6 denotes the sextet Clebsch–Gordan coefficient tensor, i.e. an elementary colour tensor with one antisextet colour index and two fundamental colour indices. Furthermore, we impose a \mathbb{Z}_2 symmetry on the DM and messenger sector to ensure DM stability. The dark matter state X and one mediator (in this case, ψ) are odd, while all other fields, including the SM ones and the other mediator (φ), are even. Consequently, only one of the mediators needs be heavier than the DM state as its decay involves the DM particle. In contrast, the other mediator decays directly into SM fields so that its mass is unconstrained by DM stability requirements.

The free parameters of this model include the three particle masses M_X , M_ψ , and M_φ , the DM-mediator Yukawa coupling λ_X , and the mediator-quark coupling matrix λ_{qq} . In our analysis, we restrict the entries of λ_{qq} to be real, while allowing flexibility in choosing the flavour structure. This flexibility is constrained by stringent limits on flavour-changing neutral currents (FCNCs), which can be enhanced by the presence of a colour-sextet scalar field coupling to uptype quarks [92]. Additionally, certain couplings to charm quarks are tightly constrained [93], leading to the following bounds:

$$(\lambda_{qq})_{11}(\lambda_{qq})_{22} \le 9.3 \left(\frac{M_{\varphi}}{1 \text{ TeV}}\right)^2 \times 10^{-7},$$

$$\sum_{I=1}^{3} (\lambda_{qq})_{I2}(\lambda_{qq})_{I1} \le 2.5 \left(\frac{M_{\varphi}}{1 \text{ TeV}}\right) \times 10^{-3}.$$
(12)



² We use the notation \mathcal{L}_{XY} for any Lagrangian including DM-mediator interactions, regardless of the model, for the purpose of uniformity.

Eur. Phys. J. C (2025) 85:975 Page 11 of 96 975

A straightforward way to comply with these constraints, as explored in [30], is to assume a flavour-diagonal coupling scheme with $(\lambda_{qq})_{22}=0$, effectively yielding a charmphobic scenario. However, completely excluding charm couplings, while convenient, is neither necessary nor symmetry-driven. On the other hand, as discussed in Sect. 3.2.3, large couplings involving third-generation quarks lead to intriguing heavy-flavour phenomenology. To balance these considerations, we adopt a coupling scheme that reflects the Standard Model quark mass hierarchy while respecting the FCNC constraints in (12).

In the investigations carried out in this white paper, we generate a UFO version of the above model with FeynRules, to be used with MG5aMC and MadDM for collider and cosmology phenomenology, respectively. This model is available upon request.

2.5 New gauge interactions to connect the dark sector to the Standard Model

Models with a vector DM field, particularly those involving a non-Abelian gauge sector, remain among the least explored extensions of the SM, despite being well-motivated. Here, gauge principles provide natural constraints and guidance limiting possible theoretical constructions (see, e.g., [94–119] for discussions of non-Abelian DM in different setups, including scenarios with non-renormalisable kinetic mixing terms or Higgs portal couplings). In this section, we highlight a recently proposed minimal framework that extends the SM gauge sector by introducing a non-Abelian gauge group under which all SM particles are singlets and for which no renormalisable kinetic mixing terms are allowed. For further details, we refer the reader to [33], while here we summarise the construction and key properties of this framework.

We consider the simplest non-Abelian group, denoted as $SU(2)_D$, to connect the SM to the dark sector, and we label the associated gauge bosons as

$$V_{\mu}^{D} \equiv \begin{pmatrix} V_{D+\mu}^{0} \\ V_{D0\mu}^{0} \\ V_{D-\mu}^{0} \end{pmatrix}, \tag{13}$$

where the superscripts refer to the field electric charges and the subscripts denote their isospin under $SU(2)_D$ (D-isospin). The spontaneous breaking of the $SU(2)_L \times U(1)_Y$ and $SU(2)_D$ gauge symmetries is achieved through two scalar doublets,

$$\Phi_{H} = \begin{pmatrix} \phi^{+} \\ \phi^{0} \end{pmatrix} \qquad \Phi_{D} = \begin{pmatrix} \varphi^{0}_{D+\frac{1}{2}} \\ \varphi^{0}_{D-\frac{1}{2}} \end{pmatrix}$$
 (14)

whose lower components get the vacuum expectation values (vevs) v and v_D . The scalar potential, that was introduced in [95], is given by

$$V(\Phi_H, \Phi_D) = -\mu^2 \Phi_H^{\dagger} \Phi_H - \mu_D^2 \Phi_D^{\dagger} \Phi_D + \lambda (\Phi_H^{\dagger} \Phi_H)^2 + \lambda_D (\Phi_D^{\dagger} \Phi_D)^2 + \lambda_{\Phi_H} \Phi_D \Phi_H^{\dagger} \Phi_H \Phi_D^{\dagger} \Phi_D,$$
(15)

and involves various bilinear (μ_i^2) and quartic (λ_i) couplings. This potential ensures the degeneracy and stability of the $SU(2)_D$ gauge bosons due to the custodial symmetry inherent to the scalar Lagrangian. Moreover, the interaction between the scalar fields via the portal term $\lambda_{\Phi_H\Phi_D}$ induces scalar mixing and modifies the couplings of the Higgs boson to the SM states, which therefore provides strong constraints on the model [120].

A new mechanism for communication between the dark and visible sectors is introduced through a vector-like fermion doublet.

$$\Psi = \begin{pmatrix} \psi_D \\ \psi \end{pmatrix},\tag{16}$$

which is charged under $SU(2)_D$ but singlet under $SU(2)_L$, and where both components of Ψ share the same hypercharge quantum numbers as an SM right-handed fermion.⁴ The mass and interaction terms for the fermion Ψ are given by

$$\mathcal{L} = -M_{\Psi}\bar{\Psi}\Psi - \left(y'\bar{\Psi}_{L}\Phi_{D}f_{R}^{SM} + \text{H.c.}\right),\tag{17}$$

where $f_R^{\rm SM}$ denotes a generic SM right-handed fermion, and y' is a new Yukawa coupling connecting the SM fermion to the dark fermion Ψ via the scalar dark doublet Φ_D . The stability of the DM is ensured by the absence of an additional Yukawa term $y''\bar{\Psi}_L\Phi_D^cf_R^{\rm SM}$, and is protected by an unbroken global $U(1)_D\equiv e^{i\Lambda Y_D}$ symmetry. Without this symmetry, the term involving the y'' coupling would be unavoidable since the scalar doublet Φ_D lies in a pseudo-real representation. The symmetry-breaking pattern in this framework is $SU(2)_D\times U(1)_D\to U(1)_D^d$. Assigning $U(1)_D$ charges as $Y_D=1/2$ for the dark scalar and fermion doublets and $Y_D=0$ for the vector triplet, there remains an invariance under the discrete subgroup $\mathbb{Z}_2\equiv (-1)^{Q_D}$, where $Q_D=T_D^3+Y_D$.

Among all new particles, the lightest \mathbb{Z}_2 -odd particle is stable. Our construction features two potential candidates, namely the $V_{D\pm}^0$ or ψ_D states, with different implications for cosmology [33]. In the present work, we focus on the scenario where the lightest \mathbb{Z}_2 -odd particle is the $V_{D\pm}^0$ boson, which we designate as the Fermion Portal Vector Dark Matter (FPVDM) framework. The theory predicts six massive

⁴ Vector-like portals have been studied for scalar DM candidates in [27,121], and for vector DM states in [119,122]. In these contexts, simplifying assumptions include neglecting new Yukawa couplings [122] or introducing an extended particle content [119].



³ Gauge kinetic mixing terms may arise at loop level, depending on the Higgs sector structure, but these correspond to suppressed higherdimensional operator contributions.

975 Page 12 of 96 Eur. Phys. J. C (2025) 85:975

gauge bosons (the $Z,\,W^\pm,\,V^0_{D0}$, and $V^0_{D\pm}$ vector bosons), whose longitudinal components correspond to six Goldstone bosons. The remaining two scalar degrees of freedom include the Standard Model Higgs boson and an additional CP-even scalar. In the unitary gauge, the scalar mass terms in the Lagrangian take the form

$$\mathcal{L}_{m}^{\mathcal{S}} = \left(h_{1} \varphi_{1}\right) \begin{pmatrix} \lambda v^{2} & \frac{\lambda \Phi_{H} \Phi_{D}}{2} v v_{D} \\ \frac{\lambda \Phi_{H} \Phi_{D}}{2} v v_{D} & \lambda_{D} v_{D}^{2} \end{pmatrix} \begin{pmatrix} h_{1} \\ \varphi_{1} \end{pmatrix}, \tag{18}$$

where h_1 and φ_1 are defined from $\phi^0 = (v + h_1)/\sqrt{2}$ and $\varphi^0_{D-1/2} = (v_D + \varphi_1)/\sqrt{2}$. Diagonalising the above mass matrix yields the scalar mass eigenvalues:

$$m_{h,H}^{2} = \lambda v^{2} + \lambda_{D} v_{D}^{2} \mp \sqrt{(\lambda_{D} v_{D}^{2} - \lambda v^{2})^{2} + \lambda_{\Phi_{H} \Phi_{D}}^{2} v^{2} v_{D}^{2}},$$
 (19)

with a mixing angle given by

$$\sin \theta_S = \sqrt{2 \frac{m_H^2 v^2 \lambda - m_h^2 v_D^2 \lambda_D}{m_H^4 - m_h^4}}.$$
 (20)

In the dark sector, the Ψ component with $T_{3D}=+1/2$ ($Q_D=+1$) does not mix with any other fermion, and its mass, $m_{\Psi_D}=M_{\Psi}$, is determined solely by the vector-like mass term in (17). In contrast, the masses of the other fermions depend on the vevs of both scalars, as well as on the SM Yukawa interactions and the one introduced in (17). The corresponding fermionic mass Lagrangian is given by

$$\mathcal{L}_{m}^{f} = \begin{pmatrix} \bar{f}_{L}^{\text{SM}} \ \bar{\psi}_{L} \end{pmatrix} \begin{pmatrix} y \frac{v}{\sqrt{2}} & 0 \\ y' \frac{v_{D}}{\sqrt{2}} \ M_{\Psi} \end{pmatrix} \begin{pmatrix} f_{R}^{\text{SM}} \\ \psi_{R} \end{pmatrix}. \tag{21}$$

The corresponding mass eigenvalues are easily obtained, and read

$$m_{f,F}^2 = \frac{1}{4} \left[\Delta \mp \sqrt{\Delta^2 - 8y^2 v^2 M_{\Psi}^2} \right], \tag{22}$$

with $\Delta = y^2v^2 + y'^2v_D^2 + 2M_\Psi^2$. Here, f represents the SM fermion connected to the dark sector, while F denotes its heavier partner, and they always satisfy the hierarchy $m_f < m_{\psi_D} \le m_F$. Let us note that in principle, a vector-like fermion may interact with one or more SM flavours, and multiple vector-like fermions may be included in the theory. In addition, the $SU(2)_D$ gauge bosons are degenerate in mass at tree level, with $m_{V_D} \equiv m_{V_{D\pm}^0} = m_{V_{D0}^0} = g_D v_D/2$ (and g_D being the $SU(2)_D$ gauge coupling). However, this degeneracy is lifted by fermionic loop corrections, which account for the differing \mathbb{Z}_2 parities of the $SU(2)_D$ gauge bosons. For simplicity, we label from now on the states $V_D \equiv V_{D\pm}^0$ of mass m_{V_D} , and $V' \equiv V_{D0}^0$ of mass $m_{V'}$. The leading contribution to the radiative mass splitting $\Delta m_V = m_{V_D} - m_{V'}$

is driven by F and ψ_D loops,

$$\Delta m_V = \frac{\varepsilon^2 g_D^2 m_F^2}{32\pi^2 m_{V_D}} + O(\varepsilon^2), \quad \text{where } \varepsilon = \frac{m_F^2 - m_{\psi_D}^2}{m_F^2}.$$
(23)

Notably, Δm_V vanishes in the limit $y' \to 0$.

To analyse the simplest realisation of the FPVDM framework, we assume that the new vector-like fermions interact exclusively with one SM flavour, that we take to be the top quark. The new physics sector is thus described by six independent input parameters,

$$g_D$$
, m_{V_D} , m_H , $\sin \theta_S$, $m_T \equiv m_F$, $m_{t_D} \equiv m_{\psi_D}$. (24)

We further simplify the parameter space by enforcing no mixing between the two scalars h and H (i.e. $\theta_S = 0$), so that the SM Higgs sector remains unaffected by new physics at tree level and the dark side of the potential mirrors the structure of the SM potential. It is important to note that for $y' \rightarrow 0$, the quartic coupling $\lambda_{\Phi_H \Phi_D}$ cannot be generated radiatively. This implies that the scalar mixing is induced solely by y' effects through fermionic loops, and can thus be safely neglected. In this configuration, the fermion sector satisfies the mass hierarchy $m_t < m_{t_D} \le m_T$, while the mass of the heavy Higgs boson H can take any value consistent with experimental bounds, including values below that of the SM Higgs boson. In this study we test this realisation of the model against multiple observables from cosmology, DM direct and indirect detection and LHC searches. For this purpose the Lagrangian has been implemented in Lanhep [123] and FeynRules [42] to generate model files for Calcher [51], in the UFO [46,47] format, as well as for FeynArts [45]. They are available from the HEPMDB database [124].

3 t-channel dark matter at the LHC

Contributions from D. Agin, C. Arina, E. Bagnaschi, K. Bai, M.J. Baker, M. Becker, A. Belyaev, F. Benoit, M. Blanke, J. Burzynski, J.M. Butterworth, A. Cagnotta, L. Calibbi, L.M. Carpenter, A.S. Cornell, L. Corpe, F. D'Eramo, A. Deandrea, A. Desai, B. Fuks, M.D. Goodsell, J. Harz, J. Heisig, A.O.M. Iorio, D. Kar, S. Kraml, A. Lessa, L. Lopez-Honorez, A. Mariotti, A. Moreno Briceño, L. Munoz-Aillaud, T. Murphy, A.M. Ncube, W. Nzuza, L. Panizzi, R. Pedro, C. Prat, L. Rathmann, T. Sangweni, D. Sengupta, W. Shepherd, A. Thamm, D. Trischuk

This section explores the phenomenology of a wide class of t-channel DM models at colliders. We begin, in Sect. 3.1, by examining the set of simplified models introduced in Sect. 2.1. In Sect. 3.1.1 we describe the general features of collider signals typical of these t-channel simplified mod-



Eur. Phys. J. C (2025) 85:975 Page 13 of 96 975

els, and explain why a straightforward naive approach is generally insufficient. Additionally, we provide details on leveraging standard Monte Carlo event generators, such as MadGraph5 aMC@NLO, to simulate these signals by matching fixed-order matrix elements at NLO accuracy in QCD with parton showers. We also discuss the use of public tools to extract current experimental constraints on these models. Next, we consider in Sect. 3.1.2 simplified models where the DM candidate couples to right-handed quarks of the first generation (i.e. u_R and d_R) and present updated results for the existing LHC constraints obtained using state-of-the-art simulations. This analysis is extended in Sects. 3.1.3 and 3.1.4, where we investigate scenarios involving couplings to quarks of the second generation (i.e. c_R and s_R) and third generation (i.e. t_R and b_R), respectively. The unique features of the signals are examined in Sect. 3.1.5, where we demonstrate again the importance of considering all signal components across different mass spectra. Finally, we briefly discuss leptophilic models in Sect. 3.1.6.

In Sect. 3.2, we leave the minimal assumption and investigate a few non-minimal setups. In particular, Sects. 3.2.1, 3.2.2, 3.2.3, and 3.2.4 delve into flavoured dark matter constructions, composite *t*-channel DM models, frustrated *t*-channel DM models, and models where the dark sector is linked through a new non-Abelian gauge interaction, respectively.

We close the discussion on *t*-channel collider phenomenology by considering, in Sect. 3.3, scenarios where the mediator is long-lived, highlighting their distinctive signatures.

3.1 Minimal simplified models – prompt decays

3.1.1 Generalities about the signal of quark-philic dark matter

The phenomenology of the simplified t-channel DM models under study must account for a description of the signal kinematics that is as accurate as possible. It should therefore include all contributions. At tree level and for models in which DM couples to quarks, this involves all diagrams leading to the production of any pair of new physics states. Specifically, this includes the production of a pair of dark matter states $(pp \to XX)$, with a squared matrix element proportional to λ^4), the associated production of a DM particle and a mediator $(pp \to XY + X\bar{Y})$, with a squared matrix element proportional to $\lambda^2\alpha_s$), and the production of a pair of mediator particles or antiparticles $(pp \to YY + Y\bar{Y} + \bar{Y}\bar{Y})$.

In the following, we label the contributions from the first two processes by XX and XY respectively, while for the last process we must distinguish the different contributing components. The pair production of a mediator and an antimediator can originate from QCD diagrams (labelled by

 $Y\bar{Y}_{QCD}$, with a squared matrix element proportional to α_s^2), t-channel DM exchange diagrams (labelled by $Y\bar{Y}_t$ with a squared matrix element proportional to λ^4), and the corresponding interference (labelled by $Y\bar{Y}_t$, with a matrix element proportional to $\alpha_s\lambda^2$). Conversely, the production of two mediators or two anti-mediators can only be induced by t-channel dark matter exchanges in models where the DM state is real, with the corresponding matrix elements being proportional to λ^4 . The total new physics cross section $\sigma_{\rm BSM}$ is thus given, with the dependence on the new physics coupling λ factorised and as a function of the new physics masses M_X and M_Y , by:

$$\sigma_{\text{BSM}} = \lambda^2 \, \sigma_{XY}(M_X, M_Y) + \lambda^4 \, \sigma_{XX}(M_X, M_Y)$$

$$+ \, \sigma_{Y\bar{Y}_{\text{QCD}}}(M_Y) + \lambda^4 \, \sigma_{Y\bar{Y}_t}(M_X, M_Y)$$

$$+ \lambda^2 \, \sigma_{Y\bar{Y}_t}(M_X, M_Y)$$

$$+ \lambda^4 \, \sigma_{YY_t}(M_X, M_Y) + \lambda^4 \, \sigma_{\bar{Y}\bar{Y}_t}(M_X, M_Y).$$

$$(25)$$

Considering a signal region of an analysis sensitive to the signal, the total efficiency ε depends not only on the mass spectrum but also on the coupling, as the latter can alter the relative contributions of the various signal components. The fiducial new physics cross section $\hat{\sigma}_{\rm BSM}$ corresponding to that region is thus given, once again with the dependence on the new physics masses M_X and M_Y introduced explicitly, by:

$$\hat{\sigma}_{\text{BSM}} = \lambda^2 \, \sigma_{XY}(M_X, M_Y) \, \varepsilon_{XY}(M_X, M_Y) + \lambda^4 \, \sigma_{XX}(M_X, M_Y) \, \varepsilon_{XX}(M_X, M_Y) + \sigma_{Y\bar{Y}_{\text{QCD}}}(M_Y) \, \varepsilon_{Y\bar{Y}_{\text{QCD}}}(M_X, M_Y) + \lambda^4 \, \sigma_{Y\bar{Y}_t}(M_X, M_Y) \, \varepsilon_{Y\bar{Y}_t}(M_X, M_Y) + \lambda^2 \, \sigma_{Y\bar{Y}_t}(M_X, M_Y) \, \varepsilon_{Y\bar{Y}_t}(M_X, M_Y) + \lambda^4 \, \sigma_{YY_t}(M_X, M_Y) \, \varepsilon_{YY_t}(M_X, M_Y) + \lambda^4 \, \sigma_{\bar{Y}\bar{Y}_t}(M_X, M_Y) \, \varepsilon_{\bar{Y}\bar{Y}_t}(M_X, M_Y).$$
(26)

In this expression, fiducial cross sections include mediator decays, $Y \to Xq$, where q represents the relevant SM quark species for the model considered. In the simulation framework used throughout this work, we always assume a small mediator width, ensuring that the narrow-width approximation (NWA) is valid, allowing mediator production and decay to be considered in a factorised way [125]. Consequently, as shown by the above equation, the kinematics originating from each component to the signal depend solely on the masses M_X and M_Y , with the coupling λ serving only to globally rescale each contribution.

Specifically, the XX component has a strong dependence on the new physics coupling λ , to the fourth power, but its modelling requires considering an additional hard visible object in the final state to be detectable. On the other hand, the various contributions involving two mediators or anti-



975 Page 14 of 96 Eur. Phys. J. C (2025) 85:975

mediators in the final state $(YY, Y\bar{Y})$ and $\bar{Y}\bar{Y}$) exhibit different dependencies on λ . Moreover, these are generally phasespace disfavoured due to the large mediator mass required to satisfy LHC constraints on coloured particles and the fact that we deal with pair production. Equation (26) also shows that, even in feebly-coupled scenarios, the QCD contribution serves as a baseline component independent of the DM mass. Conversely, other (anti-)mediator pair channels become increasingly relevant with higher λ values, and some of them can benefit from parton density enhancements. This is specially relevant for scenarios where DM couples to up or down quarks. In such cases, processes like $uu \rightarrow YY$ and $dd \rightarrow YY$ can involve a pair of initial valence quarks. Finally, XY production has a weaker λ dependence, only to the second power, but is phase-space favoured owing to the production of a lighter DM state. Consequently, there is no clear model-independent hierarchy between the different contributions, necessitating their inclusion in signal simula-

In this work, hard-scattering simulations are performed with MadGraph5_aMC@NLO [50] (version 2.9.18) using the MadSTR plugin [43]⁵ to handle resonant contributions appearing at NLO, where an intermediate mediator is produced on-shell and subsequently decays into a DM state and a SM quark. This approach ensures that such contributions are not double-counted across the three new physics processes XX, XY, and YY. For instance, at NLO, the process $pp \rightarrow XY \rightarrow X(Xi)$ includes configurations where the intermediate mediator is on-shell (to be included in XY simulations) as well as off-shell contributions (to be included in the real-emission corrections to XX production). Furthermore, LO and NLO matrix elements are determined (with MadGraph5_aMC@NLO) using the tree-level Feynman rules, R_2 contributions, and counterterms relevant to the models introduced in Sect. 2 and encoded in the UFO format [46,47] as detailed above. These matrix elements are convolved with the LO and NLO sets of NNPDF4.0 parton densities [126,127], respectively. Additionally, decays of heavy unstable particles are handled with MadSpin [128] and MadWidth [129] to retain off-shell propagation and spin correlation effects. Finally, parton showering (PS) and hadronisation effects are simulated with Pythia (version 8.306) [130] and matched with fixed-order calculations according to the MC@NLO procedure [131].

In practice, the event generation procedure is divided into dedicated simulation runs, each associated with a specific component of the new physics signal. Except for the QCD production of a pair of mediator particles and antiparticles $(Y\bar{Y}_{QCD})$ and its interference with the *t*-channel diagrams $(Y\bar{Y}_i)$, all simulations rely on the MadSTR plugin. To start the

⁵ The MadSTR plugin can be downloaded from https://code.launchpad.net/~maddevelopers/mg5amcnlo/MadSTR Plugin.



MadGraph5_aMC@NLO package with MadSTR, we need to type the following command in a shell:

```
<MG folder > / bin / mg5_aMC -- mode = MadSTR
```

Here, <MG folder> represents the directory where MadGraph5_aMC@NLO is installed. We then define the new physics process in the command line interface of MadGraph5_aMC@NLO by typing:

```
import model DMSimpt_v2_0 -<restriction>--modelname
define xx = <X state>
define yy = <Y state>
define yy~ = <Ybar state>
define yyy = yy yy~
<generate command>
```

In this snippet of script, <restriction> specifies the model restriction considered (see Sect. 2), and generic labels are introduced for the dark matter (xx) and mediator states (yy, yy~, and yyy). The placeholders <X state>, <Y state>, and <Ybar state> refer to the FeynRules names of the relevant dark matter state and mediator particles and antiparticles, as listed in Tables 1, 2, and 3.

The event generation commands for the XX, XY, $Y\bar{Y}_t$, YY_t , and $\bar{Y}\bar{Y}_t$ processes are as follows:

```
generate p p > xx xx / <excluded states> [QCD]
generate p p > xx yyy / <excluded states> [QCD]
generate p p > yy yy~ DMT=2 QCD=0 QED=0 / <excluded
    states> [QCD]
generate p p > yy yy DMT=2 QCD=0 QED=0 / <excluded
    states> [QCD]
generate p p > yy yy DMT=2 QCD=0 QED=0 / <excluded
    states> [QCD]
generate p p > yy~ yy~ DMT=2 QCD=0 QED=0 / <excluded
    states> [QCD]
```

Here, <excluded states> is a sequence listing all particles irrelevant to the chosen model restriction. We emphasise once again that it is important to address these production modes individually, as their relative importance depends on the model (particularly, the YY and $\bar{Y}\bar{Y}$ modes are relevant only for setups with self-conjugate dark matter) and the benchmark scenario chosen (i.e. the values of the mass and coupling parameters). Event generation is further performed normally, as stated in the MadSTR documentation [43]. Technically, we remove all resonant diagram contributions squared potentially arising at NLO, which corresponds to the istr=2 option of the MadSTR configuration that we set in the run_card.dat configuration file of MadGraph5_aMC@NLO.

Additionally, event generation for QCD-induced mediator pair production $(Y\bar{Y}_{QCD})$ is performed by starting MadGraph5_aMC@NLO normally, without using the MadSTR plugin as there is no resonant diagrams appearing at NLO. This is achieved by typing the following command:

```
generate p p > yy yy~ / <excluded states> [QCD]
```

Furthermore, the interference between the QCD diagrams and the *t*-channel ones for $pp \to Y\bar{Y}$ is performed at LO with the command:

```
generate p p > yy yy\sim DMT^2==2 / <excluded states>
```



Eur. Phys. J. C (2025) 85:975 Page 15 of 96 975

As it is technically not possible to handle event generation for this contribution at NLO, the corresponding predictions are rescaled with a constant K-factor $K_{Y\bar{Y}_i}$. This K-factor is defined as the geometric mean of the $Y\bar{Y}_{QCD}$ and $Y\bar{Y}_t$ K-factors,

$$K_{Y\bar{Y}_{i}} \equiv \sqrt{K_{Y\bar{Y}_{t}}} K_{Y\bar{Y}_{\text{QCD}}} = \sqrt{\frac{\hat{\sigma}_{Y\bar{Y}_{t}}^{\text{NLO}}}{\hat{\sigma}_{Y\bar{Y}_{t}}^{\text{LO}}}} \frac{\hat{\sigma}_{Y\bar{Y}_{\text{QCD}}}^{\text{NLO}}}{\hat{\sigma}_{Y\bar{Y}_{t}}^{\text{LO}}} \hat{\sigma}_{Y\bar{Y}_{\text{QCD}}}^{\text{NLO}}}.$$
 (27)

where $\hat{\sigma}^{LO}$ and $\hat{\sigma}^{NLO}$ respectively refer to cross sections evaluated at the LO and NLO accuracy in QCD.

For correct pole cancellation between the virtual and real emission contributions appearing at NLO in the above processes, it is crucial to allow non-coloured DM states to run in virtual diagrams. By default, this is forbidden in the default settings and internal mechanisms of MadGraph5_aMC@NLO. To fix this, a few core files of the code need to be modified, as detailed in appendix A of [132]. First, we add the following lines in the function is_perturbating implemented in the file base_objects.py⁶

```
is_dm = abs(self.get('pdg_code')) in
      [51,52,53,56,57,58]
if order in int.get('orders').keys() and is_dm:
return True
```

Next, we modify the function user_filter in the file loop_diagram_generation.py as follows,

As mentioned above, mediator decays into a dark matter and a quark are handled using MadSpin and MadWidth. This is achieved by updating the MadSpin card at runtime, specifying explicitly that the mediator must decay through the only open channel,

```
set max_weight_ps_point 400
decay <Y state> <X state> <q state>
decay <Ybar state> > <X state> <qbar state>
launch
```

As before, the placeholders <Y state>, <Ybar state>, <X state> and <q state> refer to the model's labels for the mediator, anti-mediator, DM particle and the relevant SM quark for the considered model restriction (see Tables 1, 2 and 3). In the case of the interference contribution between the *t*-channel and the QCD diagrams, the command set spinmode none must be additionally included. Parton showering and hadronisation are then performed using the default Pythia cards generated by MadGraph5_aMC@NLO.

Constraints on the new physics signal emerging from the considered models can be obtained by reinterpreting the results of various experimental searches for new physics signatures comprising missing transverse energy (E_T^{miss}) and jets. This procedure, in which constraints from existing LHC analyses are derived for new models not originally considered in experimental publications by implementing the analysis logic in some ad-hoc code, is commonly referred to as *recasting*. Specifically, we focus on recent exclusive LHC Run 2 searches that impose stringent requirements on a small number of jets and more inclusive searches with looser requirements enforcing the presence of a larger number of jets: ATLAS-EXOT-2018-06 [133], ATLAS-SUSY-2018-17 [134], ATLAS-CONF-2019-040 [135], CMS-SUS-19-006 [136] and CMS-EXO-20-004 [137]. The selection criteria defining these analyses have been designed to observe a signal of new physics characterised by a substantial amount of missing transverse energy, energetic jets, and no leptons. While the cuts across the different analyses are largely similar, their differences define signal regions yielding varying sensitivities depending on the signal details. In other words, for a given new physics theoretical framework, the various searches are expected to be sensitive in different parts of the model's parameter space. For instance, the requirements on the number of jet candidates in the final state (N_i) along with the definition of such jet candidates in terms of pseudorapidity $(\eta(j))$ and transverse momentum $(p_T(j))$ slightly differ, as illustrated by Table 4 where we summarise the main requirements included in the different analyses.⁸

The CMS-EXO-20-004 analysis is hence sensitive to a softer monojet-like signature compared to the ATLAS-EXOT-2018-06 or ATLAS-CONF-2019-040 analyses by virtue of a milder cut on the p_T of the leading jet, thus offering complementary sensitivity to signals with less hadronic activity. Additionally, all analyses involve different cuts on the missing transverse momentum (or the missing hadronic activity $H_T^{\rm miss}$, which is the norm of the vector sum of the transverse momenta \mathbf{p}_T of all jets in the event, in the case

⁸ We refer to the relevant publications for the exact definition of all signal regions of the considered analyses. In particular, different regions of a given analysis may involve different cut thresholds on a given observable, Table 4 indicating in this case the typical softest threshold.



⁶ The exact location of this function in the files generated by MadGraph5_aMC@NLO: depends on the version of the program.

⁷ This term will be used throughout this report.

975 Page 16 of 96 Eur. Phys. J. C (2025) 85:975

Table 4 Summary of the typical event selection cuts included in the five jets+ E_T^{miss} analyses considered. Detailed information can be found in the experimental publications [133–137]

Cuts	ATLAS EXOT-2018-06	ATLAS SUSY-2018-17	ATLAS CONF-2019-040	CMS SUS-19-006	CMS EXO-20-004
$\overline{N_j}$	∈ [1, 4]	∈ [8, 12]	≥ 2	≥ 2	<u>≥</u> 1
$ \eta(j) $	< 2.8	< 2.0	< 2.8	< 2.4	< 2.4
$p_T(j_1)$	> 150 GeV	> 50 GeV	> 200 GeV	> 30 GeV	> 100 GeV
$p_T(j_2,, j_{N_i})$	> 30 GeV		> 50 GeV		> 20 GeV
$E_T^{ m miss}$	> 200 GeV	_	> 300 GeV	_	> 250 GeV
$H_T^{ m miss}$	_	_	_	> 300 GeV	_
$E_T^{\rm miss}/\sqrt{H_T}$	_	$> 5 \sqrt{\text{GeV}}$	$> 10 \sqrt{\text{GeV}}$	_	_
$\Delta\Phi(j_i, \mathbf{p}_T^{\mathrm{miss}})$	> 0.4	_	> 0.2	> 0.3	> 0.5
$m_{ m eff}$	_	_	> 800 GeV	_	_
H_T	_	_	_	> 300 GeV	_

of CMS-SUS-19-006) and its properties, such as its separation from the jets or its significance $E_T^{\rm miss}/\sqrt{H_T}$ (with H_T being defined below). For instance, the ATLAS-EXOT-2018-06 analysis features the smallest threshold on the $E_T^{\rm miss}$ requirement, albeit with a rather hard selection on the leading jet transverse momentum. In addition, the more inclusive searches include cuts on global observables like the effective mass $m_{\rm eff}$ or the hadronic activity H_T defined by

$$m_{\text{eff}} = E_T^{\text{miss}} + \sum_j p_T(j) \text{ and } H_T = \sum_j p_T(j).$$
 (28)

We derive constraints on the quark-philic t-channel models introduced in Sect. 2 by relying on the implementation of the above analyses in the MadAnalysis 5 framework [138–140] (version 1.10.12), which uses FastJet [141] (version 3.3.4) and its implementation of the anti- k_T algorithm [142], as well as Delphes 3 [143] (version 3.5.0) and the SFS framework [144] for the simulation of the LHC detectors. Details regarding the integration of these implementation into MadAnalysis 5, along with corresponding validation notes⁹, can be found on the public analysis database of MadAnalysis 5 [145], as well as in [137,146–150]. 10

Bounds on new physics can also be obtained by means of detector-corrected particle-level measurements at the LHC instead of detector-level searches, which thus does not require an approximation of the detector response. Contur [156,157] is currently the only tool which uses this, exploiting that measurement publications from the collaborations systematically provide associated Rivet [158] routines. The

¹⁰ The different codes can be obtained from the MadAnalysis 5 dataverse [151–155].



library of reusable measurements is thus always growing organically, without the need for further work from the phenomenological community to exploit the results. The workflow for Contur starts with a UFO file [46,47] which encodes a chosen new physics model, that is then passed to an event generator to produce HEPMC [159] files. These events are next analysed with Rivet, which outputs a set of histograms displaying where the signal events would have shown up in the bank of LHC measurements having Rivet routines available. Finally, one can stack the predicted signal (properly scaled relative to cross sections) on top of the predictions from the SM, and compare to the observed data in each measurement to derive exclusions, without the need for smearing since everything is done at particle-level. In practice, measurements are grouped into orthogonal pools (defined by final state, experiment and centre-of-mass energy) to avoid double counting: only the best exclusion from a given pool is conserved. In the context of the t-channel DM models discussed in this work, the relevant pools of measurements focus on the $\ell^+\ell^-\gamma$ final state [160], the $E_T^{\rm miss}$ + jet final state [161,162], the hadronic $t\bar{t}$ final state [163] and the $\ell + E_T^{\text{miss}} + \text{jet}$ final state [164, 165].

3.1.2 First generation simplified models

We begin our phenomenological analysis of DM *t*-channel models at colliders by exploring the LHC constraints that can be imposed on models belonging to the XYZ_UR and XYZ_dR classes. We hence extend the studies of [24,25] to scenarios where DM couples to any first-generation right-handed quark, and we additionally incorporate reinterpretations of a broader set of LHC analyses. Given that complex dark matter models are disfavoured by cosmological observations, at least when DM couplings to light generations of SM fermions are considered (see [25] and Sect. 4.4.1.1), we focus exclu-

⁹ The implementation of the ATLAS-EXOT-2018-06 search in MadAnalysis 5 has been carried out in the context of the work done for the present whitepaper, and has therefore not been documented in any peer-reviewed publication. Validation details are consequently provided in appendix A.

Eur. Phys. J. C (2025) 85:975 Page 17 of 96 975

sively on scenarios involving a real, self-conjugate DM particle (i.e. models of type F3S, F3V, and S3M). The collider analyses that we recast are discussed in Sect. 3.1.1, and include ATLAS-EXOT-2018-06, ATLAS-SUSY-2018-17, ATLAS-CONF-2019-040, CMS-SUS-19-006, and CMS-EXO-20-004.

We first present the bounds obtained for various choices of the S3M_uR and S3M_dR model's free parameters in Fig. 1. These scenarios are supersymmetry-inspired, featuring a scalar mediator in the fundamental representation of $SU(3)_C$ (analogous to a right-handed up or down squark) and a fermionic Majorana DM particle (resembling a neutralino). The results are shown in the mediator and DM mass plane, (M_Y, M_X) , for several different setups. The top row of the figure corresponds to a fixed coupling value of $\lambda = 3.5$, such a large value being motivated by cosmological considerations (see Sect. 4), while the bottom row assumes a mediator width-to-mass ratio of $\Gamma_Y/M_Y = 0.05$, which guarantees that the narrow-width-approximation is valid (a necessary condition for using MadSpin for mediator decays). Exclusion limits at the 95% confidence level are displayed as solid and dashed orange lines for NLO and LO simulations respectively. The left panel of the figure shows results for scenarios where DM couples to the first-generation right-handed up quark u_R , while its right panel displays exclusions for DM couplings to the first-generation right-handed down quark d_R . Additionally, grey dotted isolines are included to represent constant λ values (in the case where $\Gamma_Y/M_Y = 0.05$) or constant Γ_Y/M_Y values (for fixed $\lambda = 3.5$). Regions of the parameter space where λ becomes too large for a perturbative treatment (although our calculations are only at first-order in λ) or where the size of Γ_Y/M_Y challenges the validity of the narrow-width approximation are shaded to indicate theoretical and/or technical limitations.

In all the figures, mass configurations to the left of the solid (dashed) orange exclusion line are ruled out at the 95% confidence level, based on state-of-the-art simulations at NLO+PS (LO+PS). For scenarios with $\lambda = 3.5$ (top row), the mediator mass is constrained to be larger than approximately 3.2 - 3.5 TeV for up-like setups and 2.5 - 2.7 TeV for down-like setups, provided that the spectrum compression, defined as $r = M_Y/M_X - 1$, is greater than 0.6 - 0.7. This corresponds to an LHC sensitivity to DM masses M_X smaller than approximately 2 TeV and 1.5 TeV for the uplike and down-like cases, respectively. The loss in sensitivity for decreasing r values (i.e. scenarios closer to the diagonal in the figures) has a twofold origin. First, the XX contribution (red contours), despite being enhanced by the large λ^4 factor as shown in (26), is only significant for very light new physics spectra. Consequently, for heavier mass spectra, such as those close to the current bounds, mediator production (both pair and associated) and subsequent decay drives the constraints. The final-state jets originating from mediator decays thus become softer for increasingly compressed spectra, leading to a reduction in signal selection efficiencies and consequently weaker bounds. Specifically, our predictions show that the LHC currently has no sensitivity to models where the DM mass $M_X \gtrsim 2.2$ TeV for up-like setups, and $M_X \gtrsim 1.5$ TeV for down-like setups.

A key feature emerging from our results is that, in general, the LHC sensitivity is primarily driven by mediator pair production (YY), as indicated by the near-overlap of the blue and orange exclusion lines. Only in scenarios with lighter DM masses (and heavy mediator mass typical in the vicinity of the current bounds) does the associated production channel (XY) begin to contribute significantly, as shown by the green contours. This contribution arises from phase-space enhancements relevant for light dark matter, combined with the large imposed value of the new physics coupling. This large coupling value $\lambda = 3.5$ also induces a hierarchy among the different components of the YY channel. Contributions from OCD diagrams are negligible, as evidenced by the teal contours, which exclude only a small portion of the parameter space. In contrast, t-channel diagrams, with their λ^4 dependence and the existence of a $q_v q_v \rightarrow YY$ sub-process driven by two valence quarks, dominate and yield exclusions equivalent to those of the full YY channel. However, caution is required for scenarios with a split spectrum, or equivalently featuring large r values such as r > 2, where the mediator width-to-mass ratio Γ_Y/M_Y generally exceeds 20%. Such a high ratio signals an ill-defined new physics setup, and challenges the validity of calculations within the narrow-width approximation.

In the bottom row of the figure, we address this last issue by considering a second class of scenarios. Here, the two new physics masses M_X and M_Y remain free parameters, but the coupling λ is dynamically computed to enforce a fixed mediator width-to-mass ratio of $\Gamma_Y/M_Y = 0.05$. Consequently, this approach eliminates any issues related to the use of MadSpin for modelling mediator decays. The resulting exclusion contours exhibit a distinct shape compared to the case where λ was fixed to 3.5. For scenarios where the mediator width-to-mass ratio is fixed, the exclusion bounds (orange line) are nearly independent of the dark matter mass, reaching 2.5-3 TeV and 1.8-2.1 TeV in the S3M_uR and S3M_dR cases, respectively. As before, the exclusion is primarily driven by the YY channel, except for scenarios involving lighter dark matter masses and a heavy mediator, where associated XY production begins to contribute. This behaviour is consistent with the dependence of the mediator pair production rates on the DM mass, as detailed in the analysis of related matrix elements and cross sections in [25]. Interestingly, the bounds become stronger for increasingly compressed spectra, but this trend arises from the progressively larger values of λ required to maintain $\Gamma_Y/M_Y = 0.05$. In such regions of the parameter space, the perturbativity



975 Page 18 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 1 Exclusion limits at 95% confidence level from the reinterpretation of several Run 2 ATLAS and CMS searches [133–137]. The results are shown for the S3M_uR (left) and S3M_dR (right) real dark matter scenarios described in Sect. 2.1, considering two configurations: $\lambda=3.5$ (top row) and $\Gamma_Y/M_Y=0.05$ (bottom row). For scenarios with $\lambda=3.5$, dotted grey lines represent isolines of constant Γ_Y/M_Y value. Conversely, for scenarios with $\Gamma_Y/M_Y=0.05$, these lines correspond to isolines of fixed λ value. Individual contributions to the

bounds are displayed for processes XX (red), XY (green), and YY (dark blue), with the YY process further decomposed into its purely QCD part (YY_{QCD} , teal) and its t-channel part (YY_t , turquoise). The yellow gradient highlights regimes where either the perturbative approach becomes increasingly invalid due to large coupling values or the narrow-width approximation loses validity due to a large mediator width-to-mass ratio

assumption underlying the entire calculation no longer holds so that the exclusion bounds should be interpreted with caution.

Another notable feature of our results is the impact of higher-order QCD corrections on the exclusion bounds. As outlined in [40], NLO QCD corrections influence not only the overall rate of the new physics signal (i.e. the com-

bined contributions from the XX, XY, and YY channels) but also significantly alter the shapes of key observables, such as the missing transverse momentum $\mathbf{p}_T^{\text{miss}}$ and the missing energy E_T^{miss} used in all LHC analyses considered (see also Sect. 3.1.5). As a result, the exclusion contours at LO and NLO are not merely related by a simple translation of each other, but exhibit more complex differences. This is



Eur. Phys. J. C (2025) 85:975 Page 19 of 96 975

evident in Fig. 1 for all scenarios considered, where the LO bounds (dashed lines) and NLO bounds (solid lines) can be compared. For S3M_uR models, global NLO effects are generally mild, except for scenarios involving a heavy mediator $(M_Y \in [2.5, 3.5] \text{ TeV})$ and a light dark matter particle ($M_X \lesssim 1 \text{ TeV}$). In this region of parameter space and for decreasing M_X values, the XY channel begins to contribute significantly, and it turns out that it exhibits a strong sensitivity to NLO corrections due to the strong coupling already entering in the LO matrix elements. Moreover, while NLO contributions are highly relevant for the YY_{OCD} and XXchannels, these channels contribute negligibly given the mass ranges probed by current LHC analyses. Conversely, QCD corrections to the dominant YY_t channel are mild, except for setups with light DM masses, where the dependence of the matrix elements on the DM mass becomes non-trivial [25]. Consequently, NLO contributions influence only a specific portion of the parameter space for S3M_uR models. The situation is markedly different for S3M dR models. Here, the distinct parton distribution functions (PDFs) involved in the YY_t and XY matrix elements lead to much stronger NLO effects for these channels, resulting in exclusion bounds that are significantly more sensitive to higher-order corrections than in the S3M_uR case. This is clearly illustrated in the figures, where shifts in the mediator mass limits approach 500 GeV for a fixed DM mass below approximately 1 TeV.

As previously outlined, significant differences are observed between the bounds applicable on S3M_uR and S3M_dR models, despite the overall similarity in the shapes of their exclusion contours. The primary distinction lies in the mass range covered. In the region of parameter space where the current bounds reside, the signal is in both cases predominantly driven by contributions from the YY_t and XY channels. These two channels are directly influenced by the flavour of the initial-state quark, with the subprocesses $uu \rightarrow YY$ and $ug \rightarrow XY$ dominating in the S3M_uR case, and $dd \rightarrow YY$ and $dg \rightarrow XY$ dominating in the S3M_dR case. These subprocesses are strongly affected by partonic luminosities and potential enhancements from valence quarks. Since the upquark content in the proton is substantially higher than the down-quark content, the bounds on S3M_dR models are consequently weaker by approximately 500-700 GeV for a given dark matter mass.

In Fig. 2, we explore the dependence of the total exclusion for each mass point, accounting for all signal contributions evaluated at NLO, on the analysis driving it. Two illustrative setups are considered: S3M_uR benchmark scenarios featuring a fixed mediator width-to-mass ratio $\Gamma_Y/M_Y=0.05$ (left panel), and S3M_dR scenarios characterised by a fixed new physics coupling value $\lambda=3.5$ (right panel). Scenarios excluded by signal regions from more than one analysis, with exclusion levels equivalent to 4.5 standard deviations or greater, are displayed in grey. These scenarios typically

exhibit a much lighter new particle spectrum compared to the current exclusion limits, represented by the solid red line. For scenarios near or beyond these limits, we use a colour code to indicate the most sensitive analysis among those considered. The figure highlights the strengths of leveraging multiple LHC analyses, each providing an optimal sensitivity to different mass configurations.

The CMS monojet analysis (CMS-EXO-20-004) is particularly sensitive to split spectra, where the dark matter state is at least a few hundred GeV lighter than the mediator. These configurations are shown in yellow in the figure. Additionally, the ATLAS-EXOT-2018-06 analysis, which has a similar selection (as depicted in Table 4), is equally sensitive to such configurations and is, in some cases, even more constraining (the relevant spectrum configuration being shown in green). Notably, these two analyses together provide the best sensitivity to heavier first-generation S3M scenarios compare to existing bounds, making them strong candidates for follow-up investigations during the LHC Run 3 and its highluminosity phase. Moreover, the more inclusive ATLAS-CONF-2019-040 analysis proves to be the most sensitive for scenarios where both the mediator and the dark matter are relatively heavy, but with a significant mass gap, as indicated by the blue regions in the figure. Conversely, the highly inclusive CMS-SUS-19-006 analysis demonstrates strong sensitivity to compressed spectra, as highlighted by the orange regions displayed in the figure. The combined sensitivity of these four analyses helps to clarify the shape and sharp features of the exclusion contour that had been found in Fig. 1. Importantly, it is known that some signal regions across these analyses are uncorrelated in light of the targeted signal, despite all these analyses targeting a final state comprising jets and missing transverse momentum [166, 167]. There thus exists some potential for their combination, as demonstrated by the TACO approach introduced in [166] and applied to a supersymmetric scenario resembling the S3M model in [167], or the approach of [168, 169]. A comprehensive reassessment of the current exclusion via analysis combination is, however, left for future work.

In Fig. 3, we turn our attention to the F3s class of models, in which the dark matter particle is a real scalar state and the mediator is a vector-like fermion lying in the fundamental representation of $SU(3)_C$. We remind that such a setup is typical of composite constructions in theories beyond the Standard Model. Similar to our study of the S3M models above, the dark matter and mediator masses are treated as free parameters, while two complementary approaches are used to define the coupling λ of dark matter to quarks. In the first approach, λ is fixed to a value of 4.8, motivated by cosmological considerations (see again Sect. 4). In the second approach, the mediator width-to-mass ratio is set to $\Gamma_Y/M_Y=0.05$, and the corresponding λ value is then dynamically determined. The obtained exclusion contours exhibit a similar shape to



975 Page 20 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 2 Search including the most sensitive signal region for the new physics signal originating from the combination of all processes at NLO for S3M_uR models with a fixed width/mass ratio (left), and S3M_dR models with a fixed coupling (right)

those of the S3M models, although they are shifted toward significantly higher mediator mass values. This shift originates from the larger cross sections associated with vectorlike quark pair production compared to coloured scalar pair production for a given mediator mass and coupling value [43, 170]. Additionally, scenarios with a fixed coupling value naturally result in higher exclusion limits than in the S3M case due to the increased chosen value for λ . Consequently, mediator masses as high as 3.5-4 TeV are excluded for F3S_uR models, and 2.7-3.5 TeV for F3S dR models. These bounds are almost insensitive to the dark matter mass, as the relevant matrix elements only depend on M_X for much larger dark matter masses. As a result, the $qq \rightarrow YY$ subprocess (with q representing either an up- or down-type quark) plays an even more dominant role compared to the S3M models when consider the parameter space as a whole. Enhanced by the high partonic luminosity associated with valence quarks, this subprocess alone is indeed sufficient to drive the exclusion bounds across the entire parameter space, with the only mild exception occurs for dark matter masses $M_X \lesssim 500$ GeV.

Finally, in Fig. 4, we examine the F3V class of models. The mediator is, as in the F3S models, a vector-like quark, but the dark matter particle is this time a vector resonance. Such a setup is again characteristic of composite constructions beyond the Standard Model. Unlike the other exclusion bounds computed in this subsection, the results are based solely on simulations at LO+PS, this limitation arising from practical constraints in treating massive vector states at NLO in our computational toolchain. As in previous analyses, we consider two options for the choice of the new physics coupling λ . In the first, λ is fixed to 1, consistent with cosmolog-

ical constraints, while in the second, λ is dynamically determined by requiring $\Gamma_V/M_V = 0.05$. Overall, the findings align with those obtained for the other two model classes. In particular, the exclusion is dominated by the YY_t channel across the entire relevant parameter space. Consequently, bounds are stronger for up-type scenarios compared to downtype scenarios due to the higher partonic luminosity associated with up quarks. In addition, the XY channel contributes marginally in scenarios with a heavy mediator and light dark matter. This is evident in the results shown for F3V_dR scenarios with $\Gamma_Y/M_Y = 0.05$ in the parameter space region defined by $M_X \lesssim 500$ GeV. However, for fixed λ scenarios, only the YY_t channel contributes for all represented contours. This is due to an artifact of the plots, where exclusion limits have been removed from regions where the particle width becomes extremely large, and thus unphysical (that we have arbitrarily chosen to correspond to $\Gamma_Y/M_Y \gtrsim 0.30$). Such large widths challenge not only the validity of the narrowwidth approximation, but also the very definition of a particle. Consequently to all these considerations, for scenarios with $\lambda = 1$ we obtain exclusion bounds for mediator masses up to 4 TeV and 3 TeV in the F3V_uR and F3V_dR cases, respectively, and for dark matter masses below 2 TeV and 1 TeV. Conversely, in scenarios enforcing a narrow mediator resonance (with $\Gamma_Y/M_Y = 0.05$), the bounds are nearly independent of M_X , this independence being related to the functional dependence of the associated matrix elements on the new physics masses [25].



Eur. Phys. J. C (2025) 85:975 Page 21 of 96 975




Fig. 3 Same as in Fig. 1, but for the F3S_uR (left) and F3S_dR (right) real dark matter scenarios. For scenarios with fixed λ values, we adopt $\lambda = 4.8$

3.1.3 Second generation simplified models

In this section, we present the results obtained when the DM and mediator particles interact with SM quarks of the second generation, namely charm and strange quarks. Following the approach used for the first-generation case, Fig. 5 shows the bounds that are applicable to the S3M_cR and S3M_sR models where the dark matter is a Majorana fermion and the mediator a coloured scalar state. In fixed-coupling scenarios, the interplay between astrophysical and cosmological constraints favours a different range of λ values compared to the first-generation case, by virtue of the existence of a

large valence up and down quark content in nuclei compared to other quarks: we consequently here focus on benchmarks with $\lambda=2.2$, for which the mediator remains narrow across the entire allowed parameter space.

For both the S3M_cR and S3M_sR classes of models, the bounds are weaker than those obtained for the corresponding first-generation models. For light DM, the mediator mass must exceed approximately 1 TeV for S3M_cR scenarios and 1.2 TeV for S3M_sR scenarios. On the other hand, in the small mass-splitting region, the bounds on the mediator mass reduce to around 500 GeV in both scenarios, while the DM mass is constrained to be larger than 300–400 GeV for lower



975 Page 22 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 4 Same as in Fig. 1, but for the F3V_uR (left) and F3V_dR (right) real dark matter scenarios. For scenarios with fixed λ values, we adopt $\lambda = 1$. Moreover, results are given at LO only

values of M_Y , and $M_Y/M_X - 1 \gtrsim 0.2$. Unlike in the first-generation case, the bounds result from a more intricate interplay between the different production channels due to the absence of a dominant same-charge YY contribution driven by valence-quark PDFs. In second-generation scenarios, the total YY channel receives comparable contributions from the t-channel diagrams (YY_t) and from the QCD ones (YY_{QCD}) , while the XY channel is additionally sensitive to the signal in a similar way as the YY contributions. Consequently, no single contribution entirely dominates, and the location of the overall bounds in the parameter space originates truly from

the combination of the various subprocesses contributing to the signal.

A key distinction between scenarios where the DM couples to charm and strange quarks is the strength of the bounds. Charm scenarios are found to exhibit bounds that are typically weaker by 100–200 GeV, particularly in the light DM parameter space region although this holds for most of the parameter space. Moreover, this feature applies both to results obtained on the basis of LO and NLO simulations. In order to understand the origin of this difference, we examine as a representative example the $YY_{\rm QCD}$ channel, which is independent of λ and dominated by gluon-initiated topolo-



Eur. Phys. J. C (2025) 85:975 Page 23 of 96 975




Fig. 5 Same as in Fig. 1, but for the S3M_cR (left) and S3M_sR (right) classes of models. For scenarios with a fixed coupling value, we adopt $\lambda = 2.2$

gies. Associated cross sections are therefore approximately independent of the nature of the quark to which the DM couples (which also holds for scenarios with couplings to up and down quarks). For benchmark scenarios relevant with the position of the bounds in the framework of light DM, the most sensitive signal region (to a new physics signal only comprising the *YY*_{QCD} contribution) is the SR2j_1600 region of the ATLAS-CONF-2019-040 analysis [135], as shown in Fig. 6 where we display information on the most sensitive analysis for all considered mass configurations. A deep investigation of the associated cut-flow reveals that the preselection criteria disproportionately impact charm scenarios. The corresponding requirements include zero leptons, a leading and

sub-leading jet with a transverse momentum imposed to be above 200 GeV and 50 GeV respectively, missing transverse energy larger than 300 GeV, and an effective mass (defined as the scalar sum of the transverse momenta of all visible objects and the missing transverse momentum) greater than 800 GeV. However, charm quarks tend to produce more leptons and softer jets during hadronisation than strange quarks, subsequently leading to a stronger signal event rejection by the preselection. This is illustrated in Fig. 7 where we show the p_T spectrum of the leading jet (left) and the distribution in the number of leptons (right), both at LO and NLO. These predictions are obtained with the standard tool chain described in Sect. 3.1.1, with a detector parametrisation set to the default



975 Page 24 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 6 Search including the most sensitive signal region for the new physics signal emerging from the YY_{QCD} processes at NLO, for the S3M_cR (left) and S3M_sR (right) models with a fixed width-to-mass ratio. The 95% CL exclusion limit is also reported




Fig. 7 LO and NLO distributions of the transverse momentum of the leading jet (left) and of the number of leptons (right) for the $YY_{\rm QCD}$ channel. We show the number of events obtained at reconstruction level, after hadronisation, detector simulation and before any selection, and for an integrated luminosity of 300 fb⁻¹. The distributions correspond to the

S3M_cR and S3M_sR scenarios with $M_Y=800$ GeV, $M_X=10$ GeV and λ fixed to obtain $\Gamma_Y/M_Y=0.05$. The cross sections reported in the legend are effective to reflect the reduction due to selecting specific final state particles

ATLAS-SFS configuration included in MadAnalysis 5. The charm signal is therefore correspondingly associated with a smaller selection efficiency then the strange signal, thus explaining the weaker exclusion bounds. Additionally, we emphasise that the contribution related to the quark-initiated processes leads to stronger bounds for S3M_SR scenarios than for S3M_CR scenarios across the entire parameter space, primarily due to the larger PDF contributions for strange quarks relative to charm quarks.

A second key difference between charm and strange scenarios lies in the relationship between NLO and LO bounds for the XY and YY_t contributions. For scenarios where the DM couples to charm quarks, NLO bounds are stronger than LO bounds, while the opposite holds for strange quarks. This behaviour is again attributed to PDF effects: as shown in Fig. 8, strange-initiated partonic luminosities are higher at LO than NLO across the entire range of partonic centre-ofmass energies $\sqrt{\hat{s}}$, whereas for charm-initiated processes,



Eur. Phys. J. C (2025) 85:975 Page 25 of 96 975




Fig. 8 NLO/LO ratios of parton luminosities for the NNPDF40 set of parton densities. We consider first the charm-gluon (top left) and strange-gluon (top right) initial states relevant for the *XY* process, and

next the charm-anticharm (bottom left) and strange-antistrange (bottom right) initial states relevant for the $Y\bar{Y}_t$ process. Band thickness refers to PDF uncertainties

the NLO/LO ratio exceeds 1 only for $\sqrt{\hat{s}} \lesssim 1$ TeV. Despite larger PDF uncertainties for charm quarks, this effect significantly impacts the interpretation of results, particularly when combined with the different distributions in the final-state jet multiplicity at NLO. Consequently, it is clear that understanding the impact of the PDFs on the bounds is of utmost importance for the interpretation of the results.

We now move on with the F3S_cR and F3S_sR classes of scenarios in which the mediator is a coloured fermion

and the dark matter a scalar state. The bounds on the simplified models are shown in Fig. 9. As expected, the overall bounds are stronger compared to S3M setups due to the higher number of spin degrees of freedom of the mediator. The latter impacts the rates of the YY_t and XY processes, that get larger than for the production of a scalar mediator, and then contribute dominantly to the combined signal. For light DM, the bounds on the mediator reach 2–2.5 TeV, depending on the scenario. For benchmarks with a fixed coupling value,



975 Page 26 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 9 Same as Fig. 5, but for the F3S_cR (left) and F3S_sR (right) classes of scenarios

caution is nevertheless necessary because of the total mediator width, which easily exceeds 10% of the mediator mass and thus makes the narrow-width approximation less reliable. Moreover, in both the charm and strange cases, the DM mass is constrained to be above approximately 800 GeV as long as the mediator is light enough to get significant bounds from its production at the LHC. In scenarios featuring a fixed width-over-mass ratio (and for DM couplings to either the strange or the charm quark), the bounds are almost entirely driven by the $YY_{\rm QCD}$ channel, at least until the λ -dependent $YY_{\rm t}$ contributions become dominant for setups where the spectrum approaches the kinematic limit where the DM and mediator masses are equal. Finally, we also note that the LO bounds

are stronger than the NLO ones for all considered scenarios, which is compatible with the PDF behaviour shown in Fig. 8.

Finally, we examine in Fig. 10 the LHC bounds imposed on the two F3V classes of models where the dark matter is a vector state, and the mediator a coloured fermion. Here, the YY_t channel dominates almost everywhere in the parameter space, except in regions where its width-to-mass ratios is around or below 1%, for scenarios with a fixed coupling value. There, the $YY_{\rm QCD}$ contribution takes over. For the two classes of models featuring a coupling to charm and strange quarks, the combined bounds exclude DM masses below 500–800 GeV, depending on the mediator mass. Moreover, in scenarios with a fixed width-over-mass ratio, a visible posi-



Eur. Phys. J. C (2025) 85:975 Page 27 of 96 975




Fig. 10 Same as Fig. 5, but for the F3V_CR (left) and F3V_SR (right) scenarios

tive interference effect between the YY_t and YY_{QCD} channels emerges when the DM is light. Numerically, mediator masses are then constrained to exceed 2 TeV almost independently of DM mass until $M_Y/M_X-1\gtrsim 0.3$, where the bounds asymptotically approach the kinematic limit for increasing mediator and DM masses.

3.1.4 Third generation simplified models

Third-generation scenarios differ significantly from those in which the mediator couples to first- or second-generation quarks. The relevant quark parton densities are either entirely absent (for top quarks) or highly suppressed (for bottom quarks). As a result, mediator pair production through QCD

interactions (YY_{QCD}) becomes the dominant contribution to the entire new physics signal. At leading order and in scenarios where dark matter couples to top quarks, YY_{QCD} hence constitutes the sole contribution to the signal, and dark matter pair production (XX) becomes accessible at NLO. For setups where the dark matter couples to bottom quarks, the situation is largely similar, with two notable exceptions. First, associated XY production plays a sub-leading role, particularly for light dark matter masses. Second, mediator pair production via t-channel exchange of a dark matter particle (YY_t) can contribute in regions with a very compressed mass spectrum.

These features are illustrated in Fig. 11, which presents exclusion contours for three classes of third-generation simplified models with fermionic dark matter in the (M_Y, M_X)



975 Page 28 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 11 Same as in Fig. 1, but for the S3M_tR real dark matter scenarios with $\Gamma_Y/M_Y=0.05$ (left), and for the S3M_bR scenarios with either $\lambda=2.2$ (central) or $\Gamma_Y/M_Y=0.05$ (right)

plane. First, the left panel of the figure focuses on supersymmetry-like top-philic S3M_tR scenarios, where the mediator is a coloured scalar similar to a top squark, the dark matter is a Majorana fermion, and the new physics coupling is dynamically set to ensure $\Gamma_Y/M_Y = 0.05$. Since the dependence on the coupling is mild due to the fact that the signal is dominated by its λ -independent QCD component, using a fixed coupling value of $\lambda = 2.2$ as motivated by cosmology would not significantly alter the results. The corresponding figure is therefore omitted from this report. The constraints obtained are relatively weak. The LHC Run 2 is indeed sensitive only to scenarios with mediator masses below 800 GeV and dark matter masses lighter than about 200 GeV. These weaker bounds, compared to those for other scenarios with Majorana DM, stem from the distinct decay patterns of the top quarks produced in mediator decays. The latter hence lead to a variety of final states differing from the simpler (hard-scattering-level) one-jet $+E_T^{\text{miss}}$ or twojet $+E_T^{\text{miss}}$ signatures. In principle, the limits could be significantly improved by incorporating dedicated $t\bar{t}+E_T^{\mathrm{miss}}$ searches in the analysis, as suggested in past studies of composite constructions [27,28]. However, the absence of validated implementations of such full Run 2 searches in public recasting tools limits this possibility. While we could extrapolate predictions from partial Run 2 results available for some time [171], it is unclear whether such a naive extrapolation would provide meaningful new insights beyond the jet+ E_T^{miss} searches already considered, as pointed out in [28]. We therefore refrain from doing so in this report. Finally, as anticipated, the exclusion limits are entirely driven by the YY_{OCD} channel. While XX contributions could in principle play a role, they are found being negligible and only relevant for scenarios with a mass only slightly larger than that of the top quark.

The central and right panels of Fig. 11 explore bottomphilic S3M_bR scenarios. In the central panel, the new physics coupling is fixed to $\lambda = 2.2$, while in the right panel, λ is dynamically determined to satisfy $\Gamma_Y/M_Y = 0.05$. The $Y \rightarrow Xb$ decay, which produces this time always one jet and missing energy (unlike for top-philic DM), impacts the sensitivity of the recast analyses compared to the S3M_tR case and strengthen the bounds, owing to a larger mediator branching ratio into the relevant final states. Additionally, the non-zero bottom-quark parton PDF introduces nonnegligible (albeit suppressed) contributions from processes beyond YY_{OCD} . Specifically, in scenarios with a fixed coupling and light dark matter, XY production slightly tightens the bounds, while in cases with a fixed mediator width-tomass ratio and a compressed spectrum, the larger new physics coupling leads to an increasing contribution from the YY_t channel. However, caution is in order when interpreting predictions in this regime, as it approaches the limits of validity for perturbative treatments. For both S3M_bR scenarios, mediator masses up to 1 TeV are found excluded for dark matter masses up to approximately 400 GeV. Similar to the S3M_tR case, these bounds are conservative, as dedicated $b\bar{b}+E_T^{\rm miss}$ searches have not been included in the analysis due to the lack of validated implementations in public tools.

In Fig. 12, we present exclusion contours for scenarios with a fermionic mediator, specifically for the F3s (top row) and F3V (bottom row) classes of models. For the top-philic cases (leftmost figures), we focus on scenarios where the coupling is dynamically set to satisfy $\Gamma_Y/M_Y=0.05$. As in the S3M_tR case, the exclusion contours in the (M_Y,M_X) plane are nearly identical to those obtained using a fixed coupling value motivated by cosmology, so the latter are omitted for brevity. No new features emerge compared to models with scalar mediators, except for an increase in the exclusion limits driven by the larger mediator pair production cross section, a direct consequence of the fermionic nature of the mediator. As a result, mediator masses up to 1.3 TeV and 1.2 TeV are excluded for the F3S_tR and F3V_tR models, respec-



Eur. Phys. J. C (2025) 85:975 Page 29 of 96 975




Fig. 12 Same as in Fig. 1, but for the F3S_tR (top left), F3V_tR (bottom left) scenarios with a fixed mediator width-to-mass ratio $\Gamma_Y/M_Y = 0.05$, and for the F3S_bR (top row, central and right panels)

and F3V_bR (bottom row, central and right panels) setups. We either fix the new physics coupling to $\lambda = 3.8$ (top central) or 0.9 (bottom central), or derive it from $\Gamma_Y/M_Y = 0.05$

tively, with corresponding dark matter masses constrained to be larger than approximately 800 GeV and 500 GeV. It is worth noting that the weaker exclusions for the F3V_tR model arise from the reliance on LO+PS simulations, lacking thus of important *K*-factor enhancement in the signal rates that cannot be computed due to technical limitations in our toolchain.

The remaining panels in Fig. 12 explore bottom-philic setups. As with the S3M_bR scenarios, we distinguish between cases where the coupling λ is fixed based on cosmological considerations (central figures) and those where $\Gamma_Y/M_Y=0.05$ (rightmost figures). For the fixed coupling case, we use $\lambda=3.8$ for F3S_bR scenarios and $\lambda=0.9$ for F3V_bR scenarios. Again, no new features emerge compared to scalar mediator cases. The limits remain dominated by the YY_{QCD} channel, with sub-leading contributions from XY production for spectrum featuring light dark matter, and from YY_t production for more compressed spectra. Subsequently, mediator masses are excluded up to 1.5-1.6 TeV in F3S_bR scenarios with dark matter masses below approximately 600-700 GeV. Strong constraints are also obtained for compressed spectra in cases where λ is derived from $\Gamma_Y/M_Y=0.05$, but

these correspond to baroque setups where perturbative methods are unreliable, so that predictions could not be trusted. For the F3V_bR model, the bounds are similar but slightly weaker, reflecting the use of LO predictions without (differential) K-factors.

3.1.5 Simplified models: considerations on signal modelling

Designing searches optimised to probe *t*-channel DM scenarios requires a detailed understanding of the distinct kinematic features of the signal final state. This involves weighing the relative importance of individual contributions and examining the effects of NLO corrections at the differential level. Such an approach enables the targeting of dominant contributions in specific signal regions, and determines whether NLO corrections are impactful enough to be experimentally observable. In the following, we present representative kinematic distributions for the different scenarios discussed in the previous sections. The aim is to highlight differences and similarities that can aid in designing new searches. All distributions are computed using the simplified fast detector



simulation (SFS) built within MadAnalysis 5 [144,172]. For these illustrative case studies, we utilised the ATLAS default settings shipped with the code.

In Fig. 13, we consider the S3M uR and F3S uR scenarios, and we show the relative contributions of the various production channels to the $E_T^{\rm miss}$ distribution for two different mass configurations close to the recast bounds at NLO: $M_Y = 3$ TeV and $M_X = 10$ or 1500 GeV. In both cases, the total width of the mediator is set to 5% of its mass (see Fig. 1), with the λ coupling determined accordingly. The differing mass gaps between the mediator and the DM result in significantly distinct shapes for the combined distributions. As already pointed out in [40], at both LO and NLO the XX contribution dominates in the low E_T^{miss} region, but the size of its contribution decreases rapidly with increasing E_T^{miss} values, making other contributions relatively more relevant. In the S3M_uR setup with a large mass gap, the XY contribution becomes dominant around $E_T^{\rm miss} \sim 1$ TeV, peaks between 1.3–1.4 TeV, and then decreases with less than one event per 25 GeV bin above approximately 1.5 TeV for an integrated luminosity of 300 fb⁻¹. In contrast, for the F3S_uR scenario, the PDF-enhanced YY_t contribution (in which same-charge mediators are produced via t-channel DM exchange) dominates above 600 GeV and strongly contributes to shape the E_T^{miss} distribution in the high E_T^{miss} range, as described in [25] (see also, e.g., [15]). When the DM state is half the mediator mass, the YY_t contribution for S3M_uR again dominates, but this time already for $E_T^{\rm miss} \sim 200$ GeV, with a peak around 1 TeV, and a fall below one event per bin above 1.7 TeV for an integrated luminosity of $300 \, \text{fb}^{-1}$. In contrast, for the F3S_uR setup, YY_t is always, and by far, the dominant contribution.

The total signal cross section in the considered S3M_uR cases varies significantly between the two mass configurations. Including all contributions to the signal, it is approximately 40 (50) times larger at NLO (LO) for $M_X = 10 \text{ GeV}$ compared to $M_X = 1500$ GeV. However, we must keep in mind that experimental selections for DM searches often impose strong cuts on the missing transverse energy. In contrast, for the considered F3S_uR scenarios the YY_t contribution scales with a weak dependence on the DM mass (for a fixed width-to-mass ratio of the mediator), while the XX cross section significantly depends on the benchmark point. The latter however peaks in a kinematic regime featuring low E_T^{miss} and low- p_T jets, and the bulk of the related events are thus usually cut away by experimental searches. The impact of the XX channel on the overall number of selected signal events is therefore mild. The markedly different shapes depicted in Fig. 13 motivate a deeper investigation into whether scenarios can be distinguished with sufficient accuracy once appropriate selection criteria and cuts are imposed, particularly for benchmarks with similar effective cross sections. We explore this aspect in detail in the rest of this section.

Figure 14 presents the missing transverse energy distributions obtained in a scenario where $\{M_Y, M_X\} = \{1600, 800\}$ GeV, but this time for the second-generation models F3S_cR (left panel) and F3S_sR (right panel). As discussed in Sect. 3.1.3, PDFs play a critical role in determining the differential K-factors and the bounds on the mediator and the DM states. At the differential level, the impact arises from the varying weights of the different contributions to the signal. While the total cross sections and the NLO distributions are similar in the two scenarios, the F3S_sR LO cross section is about 1.7 times larger than that obtained in the F3S_cR case, primarily due to the same-charge YY_t contribution. Additionally, we can note that the interference between YY_{OCD} and $Y\bar{Y}_t$ has a minimal effect of the order of a few percent, reducing for example the total event count by a few units per bin at the peak.

Figure 15 explores the impact of a simple kinematic cut of $E_T^{\text{miss}} > 200$ GeV. We consider various t-channel DM scenarios in which the λ coupling value is tuned to yield an effective cross section including all signal contributions of $\sigma^{\rm eff} \equiv \sigma \times \varepsilon = 5$ fb, with ε representing the imposed missing energy cut efficiency. In the left panel, we focus on split spectra in which we fix the mediator mass to $M_Y = 1600 \text{ GeV}$ and the dark matter mass to $M_X = 800$ GeV, scanning across a variety of S3M and F3S models featuring couplings to firstgeneration and second-generation quarks. Other scenarios, for such a choice of masses, would require large couplings leading to mediator widths incompatible with the NWA, and these are thus not represented in the figure. The resulting NLO distributions show identical shapes in the E_T^{miss} region above the cut, within statistical fluctuations, making them indistinguishable. Conversely, the LO distributions differ significantly, underscoring the importance of including NLO modelling for accurately describing final-state kinematics and total cross sections.

The right panel of the figure focuses on interactions involving the up quark, with a scenario in which we have fixed M_Y to 1600 GeV and varied M_X across the light regime (1 GeV), intermediate regime (800 GeV), and compressed regime (1595 GeV). For NLO results (with LO cross sections provided in the legend for information), scenarios yielding an identical total number of signal events exhibit visibly distinct kinematic features, already in regions with substantial event counts. This clearly reflects the different weights of the various signal contributions, which depend on the size of the λ coupling (with the exception of YY_{OCD}) and its impact on the determination of the final-state properties. For instance, in the $M_X = 1595$ GeV case, the XX contribution dominates, irrespective of the mediator and DM spins. Similarly, the $M_X = 800$ GeV distributions show consistent shapes across models. In the light DM case, the scalar mediator



Eur. Phys. J. C (2025) 85:975 Page 31 of 96 975




Fig. 13 Differential distributions for the missing transverse energy E_T^{miss} in the case of the S3M_uR (top row) and F3S_uR (bottom row) scenarios, with mass configurations $\{M_Y, M_X\} = \{3000, 10\}$ GeV (left column) and $\{M_Y, M_X\} = \{3000, 1500\}$ GeV (right column). The distributions represent the number of signal events expected for an integrated luminosity of 300 fb⁻¹ of proton-proton collisions at 13 TeV, and

have been obtained using the SFS detector simulation module included in MadAnalysis 5 with the default ATLAS configuration. The contributions of each channel with a cross section larger than 1 ab at both NLO and LO are shown, together with the resulting combined distribution



975 Page 32 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 14 Same as in Fig. 13 but for the F3S_cR (left panel) and F3S_sR (right panel) scenarios, and the mass spectrum $\{M_Y, M_X\} = \{1600, 800\} \text{ GeV}$

(S3M) exhibits a significantly larger YY_t contribution compared to fermion mediators (F3S and F3V), reflecting the different cross section scaling with the DM mass, as also seen in Fig. 13.

3.1.6 Leptophilic models

Before discussing non-minimal models, we briefly consider signals originating from prompt mediator decays in leptophilic scenarios, focusing on a model where Majorana dark matter couples exclusively to right-handed muons (as discussed in Sect. 2.1). In the region of the parameter space where the observed relic abundance is achieved via thermal freeze-out (including potential co-annihilation effects), mediators produced via Drell–Yan processes at the LHC always decay promptly. Consequently, the Yukawa coupling λ does not directly influence the LHC phenomenology.

The main collider signature in this scenario consists of two opposite-sign leptons accompanied by missing transverse energy. If the mass splitting between the X and Y states is small, the resulting leptons tend to be soft, making their reconstruction and identification challenging. This parameter space is best probed using dedicated strategies directly targeting these soft leptons, or through searches exploiting initial-state radiation (ISR) which boosts the system and produces harder leptons. In Fig. 16, we present the most stringent

ATLAS and CMS constraints on promptly decaying mediators within this benchmark scenario. The ATLAS search for slepton pair production in the decoupled region [173] constrains dark matter masses up to approximately 200 GeV (corresponding to mediator masses around 450 GeV) in the regime where the mediator mass is roughly twice that of the dark matter. Additionally, the ATLAS ISR-based search for soft leptons [174] can probe dark matter masses up to about 150 GeV when the mass splitting is around 10%. These constraints significantly improve upon LEP limits; however, viable regions of the parameter space remain, particularly at higher masses and for intermediate mass splittings $(M_Y/M_X-1 \sim 0.1-2)$, which remain challenging to probe.

3.2 Non-minimal models – prompt decays

3.2.1 Flavoured dark matter at the LHC

Compared to the minimal models discussed previously, DM models incorporating DM flavour violation (see Sect. 2.2) feature a significantly richer phenomenology at the LHC due to their more complex flavour structure. Similar to non-flavoured models, mediator pair-production remains the dominant production channel across much of the parameter space. However, both the production cross section and the mediator decay depend on the model's flavour structure.



Eur. Phys. J. C (2025) 85:975 Page 33 of 96 975





Fig. 15 Differential distributions for the missing transverse energy $E_T^{\rm miss}$ for different t-channel DM scenarios in which the mediator width is less than 10% of its mass. The coupling λ is determined to obtain an NLO total signal cross section of 5 fb after the selection cut $E_T^{\rm miss} > 200$ GeV. We either consider NLO and LO predictions for mediator and DM



masses fixed to 1600 GeV and 800 GeV respectively (left panel), or we compute NLO predictions for a scenario in which we assume a mediator mass of 1600 GeV, interactions with the up quark only, and DM masses of 10, 800 and 1595 GeV (right panel). In this case, the LO rates are indicated in the figure's legend





Fig. 16 Current LHC exclusions on the muon-philic benchmark models considered. The exclusion curves correspond to the reinterpretation of the results of the Run 2 ATLAS searches from [173] (blue) and [174]




(orange), as well as from the LHC Run 1 ATLAS results from [175] (dashed red). The dashed green curve represents the limits from the LEP2 SUSY Working Group



In this section, we focus on quark-flavoured DM within the DMFV framework, while for LHC constraints on lepton-flavoured DM models we refer to [78–81]. Moreover, for simplicity, we assume that the DM flavours X_i are approximately mass-degenerate. In contrast, non-degenerate scenarios lead to cascade decays of new particles, which opens the possibility for LLP signatures as those discussed in Sect. 3.3.

Mediator pair-production at the LHC proceeds both via QCD interactions and through t-channel exchange of the DM flavour triplet $X = (X_1, X_2, X_3)$. The latter process becomes particularly relevant when the coupling of X to the first-generation quarks is sizeable. In models with real DM representations, this t-channel exchange also enables samesign mediator pair production [76] that could be enhanced by valence quark-pair contributions as discussed in Sect. 3.1.2. On the other hand, the mediator Y decays into an SM quark and a dark flavour state X_i via the coupling matrix λ , the different branching ratios depending on the relative sizes of the elements of λ . This typically leads to several relevant signatures of mediator pair production. Consequently, LHC constraints on mediator and DM masses are generally weaker than in the single-generation scenarios discussed previously, where only a single decay model prevails.

The case of Dirac flavoured DM coupling to right-handed down-type quarks has been explored in [26]. The strongest LHC constraints arise from searches for supersymmetric squarks in final states with either two light jets or two btagged jets, both accompanied by missing transverse energy. Additionally, the compressed region is constrained by monojet searches. On the other hand, [73,74,176] examined scenarios with couplings to right-handed up-type quarks. Here, the most stringent bounds stem from recasts of supersymmetric squark searches in the $jj+E_T^{\text{miss}}$ and $t\bar{t}+E_T^{\text{miss}}$ final states, with the precise exclusion limits depending on the flavour structure of the matrix λ . The blue shaded regions in Fig. 17 show the resulting constraints, derived from the experimental search [136], on the mediator mass M_Y as a function of the first-generation coupling strength D_1 . The limits are shown for two benchmark DM freeze-out scenarios, a first one featuring quasi-degenerate freeze-out (QDF) and a second one predicting single-flavour freeze-out (SFF), their cosmology being discussed in Sect. 4.4.2.1. QDF setups consist in scenarios in which all three dark flavours are quasi-degenerate, hence participating all together in the thermal freeze-out process through their combined (co-)annihilations. On the other hand, SFF configurations refer to scenarios where only one particle flavour remains thermally active and undergoes freeze-out, while other flavour states are too heavy to contribute. In addition to these flavour-conserving final states, flavoured DM models also give rise to flavour-violating signatures. For instance, [176] proposed search strategies dedicated to single-top final states, which arise in flavoured DM models coupling to up-type quarks and where mediator pair-



Turning to scenarios with Majorana flavoured DM, models with coupling to right-handed up-type quarks have been studied in [76,77]. The key difference between a Majorana scenario and the Dirac case is the possibility of same-sign mediator pair production via t-channel exchange of X states with a mass insertion. If X has significant coupling to firstgeneration quarks, this process is strongly enhanced due to the large up-quark PDF in the proton. As a result, constraints from charge-insensitive searches, such as those exploiting the $jj + E_T^{\text{miss}}$ and $t\bar{t} + E_T^{\text{miss}}$ signatures with zero or one lepton in the final state, are considerably stronger (at least for nonzero DM masses M_X). Additionally, in final states containing top quarks where the charge can be reconstructed, samesign mediator pair production leads to unexplored smokinggun signatures. One such process is the production of two positively charged top quarks in association with missing transverse energy, $tt + E_T^{\text{miss}}$, that could be probed through dileptonic decays of the tt system [76]. However, due to the small leptonic branching fraction of the top quark, a naive estimate suggested that this channel is not competitive with standard searches, necessitating a more refined analysis [77]. A promising alternative could rely on the charge asymmetry a_{ti} of single-top final states [77], defined as

$$a_{tj} = \frac{\sigma(tj + E_T^{\text{miss}}) - \sigma(\bar{t}j + E_T^{\text{miss}})}{\sigma(tj + E_T^{\text{miss}}) + \sigma(\bar{t}j + E_T^{\text{miss}})}.$$
(29)

The single-top search strategy developed in [176] can be directly applied here, as it relies on leptonic top decays. In Dirac models, a_{tj} is then expected to be close to zero, making this asymmetry a powerful discriminator between Majorana and Dirac flavoured DM. As shown in Fig. 19, large positive values, $a_{tj} \sim 1$, are possible in regions of parameter space where the total single-top cross section reaches up to 100 fb. These regions are not excluded by standard LHC searches, as well as by the cosmological probes discussed in Sect. 4.4.2.1.



Eur. Phys. J. C (2025) 85:975 Page 35 of 96 975





Fig. 17 Expected limits from single-top final states on a model with Dirac flavoured DM coupling to right-handed up-type quarks, shown for the QDF (left) and FFS (right) benchmark scenarios of DM freezeout. Solid lines indicate the expected reach in the (M_Y, D_1) plane for $tj + E_T^{\text{miss}}$ and $t + E_T^{\text{miss}}$ analyses, assuming an integrated luminosity of


Fig. 18 Representative Feynman diagrams illustrating the production of two dark matter particles in association with a single top quark. Adapted from [176]




In addition, they can still accommodate the observed DM relic density through canonical freeze-out. Conversely, in conversion-driven freeze-out scenarios, a_{tj} is found to be close to zero due to the specific coupling structure of this mechanism. Such cases should instead be probed through LLP signatures, as discussed in Sect. 3.3.

3.2.2 Composite dark matter at the LHC

The analysis presented so far has focused on the minimal t-channel dark matter scenarios described in Sect. 2.1. In the current section, we explore the impact of non-minimality by considering the composite constructions detailed in Sect. 2.3. We begin this exploration with the class of models described by the Lagrangian (8), which introduces two mediators: a \mathbb{Z}_2 -odd t-channel mediator $Y \equiv Y_t$ and a \mathbb{Z}_2 -even vector-like quark $Y' \equiv Y'_t$. This setup hence corresponds to an F3S_tR scenario extended by the addition of the Y' mediator, whose presence is subsequently expected to modify the bounds derived in Sect. 3.1.4. As previously determined, YY pair production, followed by the decay $Y \to Xt$, dominates the total new physics signal relevant for DM production at the LHC,



137 fb⁻¹ at the 14 TeV LHC. The excluded regions lie to the left of the curves. Dashed lines show the corresponding projections for 300 fb⁻¹, and results for 3000 fb⁻¹ can be found in [176]. The shaded region represents the exclusion derived from a recast of the CMS search [136], and the orange dash-dotted lines indicate parameter values that yield the correct relic abundance. Figure adapted from [176]



with sub-leading contributions from XX dark matter pair production and XY associated production. As a result, scenarios with light dark matter ($M_X \lesssim 400 \text{ GeV}$) are excluded if the mediator mass is smaller than approximately 1.3 TeV. For intermediate dark matter masses ($M_X \in [500, 600] \text{ GeV}$), the constraints that could be imposed on the mediator are significantly weakened. Finally, heavier dark matter scenarios are in principle reachable; for instance, it was shown in [29] that for $M_X \sim 700$ GeV, top partners with masses up to 1 TeV could be excluded, provided that the spectrum is not too compressed (i.e. $M_Y > M_X + M_t$, where M_t is the mass of the top quark). The analysis of the impact of searches for dark matter in the $t\bar{t} + E_T^{\text{miss}}$ channel is however left for future work, due to the absence of validated implementations of corresponding full Run 2 ATLAS and CMS searches in public recasting tools.

The introduction of Y' mediators must comply with ATLAS and CMS searches for vector-like top partners decaying into a SM top quark and an electroweak boson. These searches generally target signatures of the process $pp \rightarrow Y'Y' \rightarrow ttVV$, where V denotes a Z boson, a W boson, or a Higgs boson h, and where the two produced vector-like



975 Page 36 of 96 Eur. Phys. J. C (2025) 85:975





Fig. 19 Predictions for the single-top charge asymmetry a_{tj} in a model of Majorana flavoured DM coupling to the three flavours of right-handed up-type quarks. We present predictions for a_{tj} (colour-coded) at the 14 TeV LHC, compared with current constraints from tops+ $E_T^{\rm miss}$ (light grey) and jets+ $E_T^{\rm miss}$ (dark grey) searches, for $D_2=0$, $M_Y=0$



1200 GeV, and $M_X = 400$ GeV (left); we also show the correlations between predictions for a_{tj} and the total single-top cross section σ_{tot} as defined in the denominator of (29), for viable scenarios featuring canonical DM freeze-out (green) and conversion-driven freeze-out (blue). Here, D_i represent the DM coupling strengths to the i^{th} generation, and the figures have been adapted from [77]

quarks decay similarly. Consequently, we enforce $M_{Y'} \ge$ 1.3 TeV. The presence of the Y' mediator can, in principle, affect dark matter production in association with a jet (i.e. the XX channel) through additional box diagrams involving both Y and Y' mediators in the loops. However, within the parameter space favoured by cosmological considerations, these diagrams contribute negligibly, ensuring that new contributions to monojet or multijet + missing energy signals remain minimal. For scenarios in which $M_{Y'} > M_Y + M_X$, the decay channel $Y' \rightarrow YX$ becomes kinematically open, leading to new contributions to the $t\bar{t}+E_T^{\rm miss}$ signal through the process $pp \to Y'\bar{Y}' \to YX\bar{Y}X \to t\bar{X}X\bar{t}XX$. This can potentially modify the constraints existing on both the Y and Y' states. However, for light dark matter, the mass of the Y mediator is already independently constrained to be at least 1.3 TeV, leaving no room for constraint weakening. For heavier dark matter, the condition $M_Y + M_X < M_{Y'}^{\min} = 1.3 \text{ TeV}$ (the minimum allowed mass for a vector-like quark) is never satisfied for scenarios reachable at the LHC Run 2. As a result, collider bounds on composite dark matter models remain unaffected by this first exploration of non-minimality.

We extend our exploration of non-minimality by considering the Lagrangian (9), which incorporates top partial compositeness into the F3S_tR class of models. This results in a scenario featuring three vector-like quark mediators $(Y_t, Y_{Q,t}, \text{ and } Y_{Q,b})$, one dark matter state (X), and two new couplings $(\lambda_Q \text{ and } \lambda_t)$. To simplify the resulting six-dimensional


parameter space, we assume equal masses for all mediators $(M_Y \equiv M_{Y_t} = M_{Y_{Q,t}} = M_{Y_{Q,b}})$ and equal couplings $(\lambda \equiv \lambda_O = \lambda_t)$.

In the left panel of Fig. 20, we present exclusion bounds derived from the reinterpretation of the full new physics signal in this model, which includes contributions from mediator pair production, mediator-DM associated production, and DM pair production. Each component of the signal implicitly include a sum over all possible mediator combinations, and corresponding simulations are achieved at LO for simplicity. To highlight the regions of parameter space favoured by cosmology, we additionally overlay the region in which the DM relic density predictions agree within 20% with Planck data [178]. The constraints, based on the searches detailed in Sect. 3.1.1 (blue contour), demonstrate that scenarios with light DM (M_X < 500 GeV) and mediators lighter than 1.8 TeV are excluded. These bounds, primarily driven by the ATLAS-CONF-2019-040 search results, are significantly stronger than those reported for the F3S models in Fig. 12. This enhanced sensitivity arises from the combined contributions of the three QCD production channels corresponding to $Y_t, Y_{O,t}$, and $Y_{O,b}$ pair production. Notably, while the results are obtained for a coupling value of 1, the dominance of QCD contributions ensures that this choice has little impact on the exclusion bounds.

For comparison, we also consider constraints derived from recent detector-corrected SM measurements provided by the



Eur. Phys. J. C (2025) 85:975 Page 37 of 96 975






Fig. 20 Exclusion limits at 95% confidence level (CL) for the composite DM model with partial compositeness considered. Results are shown in the (M_Y, M_X) plane, where M_Y is the common mediator mass, M_X the DM mass, and with fixed new physics couplings of 1. The left panel displays the region yielding the correct DM relic abundance (green), and exclusions from LHC new physics searches (blue,

solid) and SM measurements (red, dashed). The right panel shows the most sensitive analysis pool at each grid point, to which we superimpose 95% observed (solid), 68% CL observed (dashed), and 95% CL expected (dotted) exclusions. Analysis pools include $\ell^+\ell^-\gamma$ (light brown), $E_T^{\rm miss}+{\rm jets}$ (green), hadronic $t\bar{t}$ (dark brown), and $\ell+E_T^{\rm miss}+{\rm jet}$ measurements, with the black area denoting the unconstrained region

LHC collaborations (red exclusion). These constraints are somewhat less stringent than those resulting from searches for new physics, as the latter utilise additional kinematic variables to enhance sensitivity. In the right panel of Fig. 20, we identify the most sensitive 'pool' of analyses used by Contur at each point in the parameter space. The majority of the excluded region is dominated by the 'missing energy plus jets' pool, which aggregates several analyses relying on this final-state signature in 13 TeV LHC data. In particular, a 13 TeV ATLAS measurement of the differential cross section for missing energy production [161] is responsible for most of the exclusion. The discrepancy between the expected and observed exclusions in this figure arises from a small excess in the unfolded $E_T^{\rm miss}$ spectrum around 1200 GeV. Consequently, any signal favouring this kinematic regime slightly improves the data/prediction agreement, leading to a marginally weaker exclusion limit.

3.2.3 Frustrated dark matter at the LHC

In this section, we continue with our study of non-minimal models of *t*-channel dark matter. In the fDM model introduced in Sect. 2.4, the only efficient mechanism for DM annihilation involves the *t*-channel exchange of the fermionic mediator ψ , where DM particles annihilate into a pair of scalar mediators, $X\bar{X} \rightarrow \varphi^{\dagger} \varphi \rightarrow qq\bar{q}\bar{q}$. However, this




Fig. 21 Representative Feynman diagram for scalar sextet mediator production in association with a W boson in the fDM model, followed by a leptonic W decay and a φ decay to same-sign top quarks

2 oup 4 process, while relevant for the DM relic abundance, is not directly relevant for the LHC. Instead, the model can be probed at colliders in a number of ways, ranging from jets $+E_T^{\rm miss}$ analyses to direct searches for the colour-charged mediators. The latter category is particularly versatile because colour-charged particles appear in a wide variety of constructions beyond the SM, so many existing searches could be used to constrain this fDM model through reinterpretations. One promising channel is the t-channel single production of the scalar mediator φ in association with a W boson, as illustrated by the representative Feynman diagram of Fig. 21 (that includes specific W and φ decays). Whereas



975 Page 38 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 22 Cross section for scalar mediator production in association with a W boson at the LHC, for the fDM scenario considered. Results are inclusive with respect to the W-boson decay, but the mediator φ is enforced to decay into a same-sign top pair. Calculation applies to the benchmark scenario defined in (31), and for a hadronic centre-of-mass energy of $\sqrt{S} = 14$ TeV. A flat K factor of 1.3 is included to estimate the NLO yields [90]

this t-channel process leads to diverse final states depending on the decays of the φ mediator and the W boson, the figure illustrates a scenario where the W boson decays hadronically, and the scalar sextet mediator decays into like-sign top quarks ($\varphi \to tt$).

The production of same-sign top pairs is quite distinctive compared to conventional SM $t\bar{t}$ production, and it has therefore been previously searched for at the LHC [179]. This channel is particularly relevant to the considered fDM model, since top-philic sextet scalars yield very large associated rates. This enhancement arises from a chirality flip due to the quark exchanged in the t-channel (see the Feynman diagram in Fig. 21): the W boson couples to left-handed quarks, while the scalar φ couples exclusively to right-handed quarks, as shown in the Lagrangian (11). Consequently, the resulting cross section is proportional to the mass of the exchanged up-type quark, which leads to a strong enhancement for the third generation. Specifically, the parton-level cross section $d\hat{\sigma}$ for $W\varphi$ production via ud_f fusion is given, for a specific up-type quark flavour u_f in the t-channel (assuming a diagonal CKM matrix), by

$$\frac{\mathrm{d}\hat{\sigma}}{\mathrm{d}t} \left(ud \to W^{-}\varphi \right) = \frac{\alpha_2}{48} \left(\lambda_{qq} \right)_{1f}^2 \left(\frac{M_{u_f}}{M_W} \right)^2 \\ \left[M_W^2 (M_{\varphi}^2 - 2s - t) - t (M_{\varphi}^2 - s - t) \right], \tag{30}$$

where $\alpha_2 = g_2^2/4\pi$ with g_2 being the $SU(2)_L$ gauge coupling, and where M_W denote the mass of the W boson. The presence of the $M_{u_f}^2$ factor limits the sensitivity of the LHC to this process, unless the exchanged quark is a top quark

 $(f=3, M_{u_f}=M_t)$. Therefore, as alluded in Sect. 2.4, we consider a benchmark sextet-quark coupling matrix λ_{qq} that prioritises sextet couplings to top quarks. Non-flavour-diagonal couplings are however also essential, due to the parton distribution functions involved in the production process. For this purpose, we fix the scalar sextet coupling matrix λ_{qq}

$$\lambda_{qq} \equiv \frac{1}{2} \begin{pmatrix} 2 \times 10^{-5} & 10^{-4} & 0.25 \frac{M_{\varphi}}{1 \text{ TeV}} \\ 10^{-4} & 2 \times 10^{-3} & 10^{-2} \\ 0.25 \frac{M_{\varphi}}{1 \text{ TeV}} & 10^{-2} & 2 \end{pmatrix}, \tag{31}$$

which maximises the production cross section for same-sign top pairs while satisfying the FCNC constraints (12) for any sextet mass. A less aggressive benchmark with a constant value for $(\lambda_{qq})_{13} \equiv \lambda_{13} = 0.325$ is also valid, provided that $M_{\varphi} > 1.3$ TeV. This last bound is anyway almost certainly required in view of the results of LHC searches for the production of four top quarks [30]. The dependence of the hadron-level cross section $\sigma(pp \to W\varphi)$ with $\varphi \to tt$ on the sextet mass M_{φ} , in which the sum with the charge-conjugate process is implicit, is shown for this benchmark in Fig. 22, highlighting a high signal rate that is in principle not impossible to observe at the LHC. Predictions are obtained for a hadronic centre-of-mass energy of 14 TeV, and the LO set of NNPDF2.3 parton densities [180].

The size of the signal cross section, of at most O(1) fb, nevertheless poses a significant challenge for its discovery at the LHC, even with the full high-luminosity dataset of $\mathcal{L} = 3 \text{ ab}^{-1}$. Nonetheless, the distinctive kinematics of the process shown in Fig. 21 could allow the exclusion of a colour-sextet mediator in the mid-TeV mass range, improving upon current limits from searches for pair-produced colour-charged resonances [181]. This t-channel signal produces a pair of boosted top quarks which, if they decay hadronically, can potentially be reconstructed. The invariant mass of the di-top system should then be localised within a narrow window around the mediator mass. Additionally, we enforce the recoiling W boson to decay leptonically. This provides exactly one lepton and missing transverse momentum that could be used to suppress backgrounds such as $t\bar{t}$ events. A $t\bar{t}$ and tt pair cannot indeed be distinguished in the fully hadronic channel, so that other means are in order to control the associated background.

The final state of interest thus consists of multiple jets, including two *b*-tagged jets, a single lepton, and missing transverse momentum. The dominant SM backgrounds for this signature are opposite-sign top-quark pair production in association with a *W* boson, and opposite-sign bottom-quark pair production in association with three electroweak bosons including at least one *W* boson. We produce samples of 10⁵ events for each of the two background processes with MadGraph5_aMC@NLO, that we normalise according to



Eur. Phys. J. C (2025) 85:975 Page 39 of 96 975

cross sections of 769 fb [50,182] and 1211.9 fb [50]. In addition, for the ttW sample, we combine matrix elements featuring up to two additional partons following the MLM procedure [183,184]. Finally, for the signal, we produce samples of 10^5 events for an array of colour-sextet scalar masses $M_{\varphi} \in [1300, 2100]$ GeV.

To motivate the selection criteria for our search, we present two distributions of observables with good discriminating power. These distributions are generated using MadAnalysis 5 [138–140], after performing object reconstruction with its built-in simplified fast detector simulator (SFS) [144]. The latest version of MadAnalysis 5 [185] supports jet reclustering and includes an implementation of HepTopTagger [186,187]. These features enable the creation of two collections, one of narrow jes and one of fat jets. In our analysis, narrow jets are reconstructed using FastJet [141] with the anti- k_t algorithm [142] and a radius parameter of R = 0.4. Fat jets are instead clustered using the Cambridge-Aachen algorithm [188–190] with R = 1.0.

The left panel in Fig. 23 shows the distribution in the transverse momentum $p_T(J_1)$ of the leading fat jet J_1 for the background processes and a representative signal sample with $M_{\varphi} = 1.3$ TeV. In addition, the right panel in this figure presents the invariant mass $m_{J_1J_2}$ of the system made of the two highest-momentum fat jets J_1 and J_2 . These distributions are computed for events passing basic selection criteria, including the presence of at least two fat jets with $p_T > 200$ GeV. As expected (and as a consistent validation of our analysis framework), the $m_{J_1J_2}$ spectrum exhibits a pronounced peak at the mass of the colour-sextet resonance for the $M_{\varphi} = 1.3$ TeV signal, with no analogous structure in the background samples. This finding suggests that $m_{J_1J_2}$ could be a powerful variable not only for suppressing the SM backgrounds, but also for distinguishing between different new-physics scenarios with varying M_{φ} . The complete selection criteria are collected in Table 5.

We now present the results of our mock search. For this analysis, we estimate the expected 95% CL limit under the assumption of no signal observation, using the Asimov approximation for the median signal significance [191],

$$S = \sqrt{2\left[\left(N_{\rm s} + N_{\rm b}\right)\ln\left(1 + \frac{N_{\rm s}}{N_{\rm b}}\right) - N_{\rm s}\right]},\tag{32}$$

where N_s and N_b represent the signal and background yields after selection. In the limit where $N_s \ll N_b$, this expression simplifies to the well-known formula

$$S \to \frac{N_{\rm s}}{\sqrt{N_{\rm b}}} + O(N_{\rm s}^2/N_{\rm b}^2).$$
 (33)

However, this approximation does not hold uniformly across our parameter space following the applied selection criteria. For simplicity, we take S=2 as an approximate threshold

for 95% CL exclusion (the correct value being S=1.64 [191]). It should be noted that our calculation assumes no uncertainties in the signal and background yields, leading to an optimistic estimate of sensitivity. We show in Table 6 the yields and cut-by-cut efficiencies for the background processes and the $M_{\varphi}=1.3$ TeV signal. The most effective selection cuts are those on the number of fat jets (N_J) , the transverse momentum of the leading fat jet, and the final binned cut on the invariant mass of the pair of leading fat jets.


Using analogous results for all signal samples, the left panel of Fig. 24 presents the sensitivity S as a function of the LHC integrated luminosity, $\mathcal{L} \in [139, 3000] \, \mathrm{fb}^{-1}$. Ultimately, our simple cut-and-count strategy can probe coloursextet scalars with masses up to $m_{\varphi} \sim 1.95 \, \text{TeV}$ using the full planned HL-LHC dataset. This represents a best-case scenario for the method that we introduced, as our significance estimate does not account for background uncertainties or the presence of reducible backgrounds. Nevertheless, under these idealised assumptions, as shown in the right panel of Fig. 24, our proposed strategy can exclude mediator masses up to 700 GeV beyond the Run 2 limits imposed on the pair production of colour-sextet scalars in a similar benchmark scenario [30], provided no excess is observed. It should be noted that the cross section in Fig. 24 is not the inclusive cross section shown in Fig. 22, and instead incorporates appropriate branching ratios reflecting the final states targeted by our search. Moreover, the projections exceed the luminosityscaled improvements in new-physics bounds from measurements of the $t\bar{t}t\bar{t}$ production cross section and associated searches for new resonances, which extend to 2 TeV for colour-octet scalars and are expected to be slightly weaker for colour-sextet scalars [192, 193]. Nevertheless, a 5σ discovery in the same-sign top channel at the HL-LHC appears unlikely, although as discussed elsewhere in this report, multiple experimental approaches could be used to further explore frustrated dark matter. While direct LHC searches and astrophysical constraints may offer greater sensitivity, our strategy could serve as a complementary avenue for such studies.

3.2.4 A fermionic portal to a non-Abelian dark sector

Moving further along non-minimality, we discuss now the collider phenomenology arising from the FPVDM construction featuring a non-Abelian dark sector described in Sect. 2.5. Our model implementation in the UFO format has been used in MadGraph5_aMC@NLO [50] for the determination of the LHC constraints, relying on collider simulations at LO using the NNPDF3.0 LO set of parton densities [127,194]. Moreover, a simplified version of the model has been implemented to calculate cross sections at one loop in MadGraph5_aMC@NLO and FormCalc [195].



975 Page 40 of 96 Eur. Phys. J. C (2025) 85:975






Fig. 23 Distribution in the leading fat jet transverse momentum $p_T(J_1)$ (left), and in the invariant mass $m_{J_1J_2}$ of the pair of leading fat jets (right), for a signal scenario with $M_{\varphi} = 1.3$ TeV (i.e. $\varphi_Y \to tt$

production in association with a W^{\pm} boson, followed by a leptonic W-boson decay and hadronic top decays), and the two components of the irreducible SM background

Table 5 Selection criteria in our proposed search for TeV-scale colour-sextet scalar mediator produced in association with a leptonically decaying W boson, and decaying to like-sign top quarks ($\varphi \to tt$) which themselves decay hadronically

Selection criterion	Selection ranges
Narrow jets, anti- $k_{\rm T}$ with $R = 0.4$	$N_j \ge 2$
	$p_{\rm T}(j) > 20 {\rm GeV}, \eta(j) < 2.5$
	Veto any pair with invariant
	$\max m_{jj} \in [81, 101] \text{ GeV}$
b-tagged narrow jets	$N_b \ge 2$
Fat jets, Cambridge-Aachen with $R = 1.0$	$N_J \ge 2$
	$p_{\rm T}(J) > 200 \text{ GeV}, \eta(J) < 2.5$
Leading fat-jet transverse momentum	$p_{\mathrm{T}}(J_1) \ge 475 \mathrm{GeV}$
Lepton from decaying W boson	$N_\ell=1$
	$p_{\rm T}(\ell) > 25 {\rm GeV}, \eta(\ell) < 2.5$
Overlap removal	Veto any narrow jet with $\Delta R(j, \ell) < 0.1$
	Veto any lepton with $\Delta R(\ell, j) < 0.4$
Missing transverse momentum	$E_T^{ m miss} > 15~{ m GeV}$
Binned cut: reconstructed top pair invariant mass	$m_{arphi} - m_{tt}^{ m had} < 250~{ m GeV}$

For the present analysis, we consider a scenario where the new fermions are partners of the top quark. LHC bounds have been derived by confronting the signal originating from t_D pair production, followed by decay into the DM state V_D and top quarks, against the results of CMS searches for top squark pair production decaying into DM and tops with partial Run 2 data [196]. Moreover, we have imposed limits on $T\bar{T}$ production, which are approximately given by $m_T > 1.5$ TeV [197,198], keeping in mind that single T production is less constrained due to rate suppression by the small T-t mixing. Additionally, we have estimated the relevance of V' pair production as well as that of the associated production of a V' boson with the Higgs boson, which occur at LO via fermion

loops. Representative Feynman diagrams for all these processes are displayed in the left panel of Fig. 25.

We present in the right panel of Fig. 25 the regions of the parameter space allowed by LHC constraints. The bounds are projected onto the (m_{t_D}, m_{V_D}) plane for a representative benchmark point with $g_D = 0.3$, $m_T = 1600$ GeV, and $m_H = 1000$ GeV. In the small g_D limit, where the width of t_D is narrow, the t_D pair production cross section depends solely on m_{t_D} , as it is governed by the strong interaction. Consequently, as long as the mass difference between the t_D and V_D states remains well above the top-quark threshold, the exclusion limits are largely independent of m_{V_D} and rule out values of $m_{t_D} \lesssim 850$ GeV. However, as this threshold is approached, the amount of missing transverse energy orig-



Eur. Phys. J. C (2025) 85:975 Page 41 of 96 975

Table 6 Illustrative cut-flow for the selection strategy given in Table 5 and a signal benchmark featuring $M_{\varphi} = 1.3$ TeV, shown together with the cut-flows related to the irreducible backgrounds. We present the

results as yields N_i normalised from an initial number of events set to 10^5 , as well as consecutive selection efficiencies ε_i

Selection	$M_{\varphi} = 1.3 \mathrm{TeV}$	$M_{\varphi} = 1.3 \mathrm{TeV}$		$b\bar{b}$ + triboson		$t\bar{t} + W$	
	$\overline{N_i}$	ε_i	$\overline{N_i}$	ε_i	$\overline{N_i}$	$arepsilon_i$	
Initial	1.00×10^{5}	_	1.00×10^5	_	1.00×10^{5}	_	
$N_j \ge 2$	9.95×10^{4}	0.995	9.74×10^{4}	0.974	9.67×10^{4}	0.967	
No jets in Z window	9.24×10^{4}	0.929	1.63×10^{4}	0.167	8.02×10^{4}	0.830	
$N_b \ge 2$	4.29×10^{4}	0.465	1.11×10^{4}	0.685	2.92×10^{4}	0.363	
$N_J \ge 2$	3.78×10^{4}	0.881	2.51×10^{3}	0.226	6.01×10^{3}	0.206	
$p_{\mathrm{T}}(J_1) \ge 475\mathrm{GeV}$	3.02×10^{4}	0.797	1.98×10^{2}	0.079	1.14×10^{3}	0.190	
$N_{\ell} = 1$	2.41×10^{4}	0.799	3.47×10^{1}	0.175	2.68×10^{2}	0.234	
$p_{\rm T}^{\rm miss} > 15{\rm GeV}$	2.35×10^{4}	0.973	3.26×10^{1}	0.938	2.66×10^{2}	0.993	
$m_{tt}^{\rm had} > 1050 {\rm GeV}$	2.06×10^{4}	0.878	1.55×10^{1}	0.475	1.20×10^2	0.451	




Fig. 24 Projected LHC sensitivity S (left) to the considered fDM colour-sextet scalar signal and associated exclusion (right), following the search strategy proposed. Our sensitivity predictions are given as a

function of the HL-LHC integrated luminosity, the exclusion ones for $\mathcal{L}=3\,\mathrm{ab^{-1}}$, and we take $\mathcal{S}=2$ as an estimate of the 95% CL exclusion threshold

inating from the signal decreases, reducing the sensitivity of the searches considered. This results in scenarios with a small mass gap between t_D and V_D remaining allowed by current data. Subsequently, throughout the entire parameter space, dark matter masses above approximately 500 GeV are hence never excluded by the considered CMS search. In the degenerate region where $m_{t_D} \simeq m_{V_D}$, the t_D state becomes increasingly long-lived, particularly for lower values of g_D , since its width is proportional to g_D . The only available decay mode allowed by the \mathbb{Z}_2 -odd nature of the top partner, $t_D \to V_D t^{(*)}$, implies that the model is testable only through searches dedicated to long-lived particles. Notably, variations in the masses of the T and H states do not significantly affect this qualitative picture.

On the other hand, we find that V' pair production and V'h associated production are only accessible in parameter regions already excluded by cosmological constraints (see Sect. 4.4.2.4), rendering these channels irrelevant for our analysis. For completeness, Fig. 25 also includes cross-

section isolines for hV' and V'V' production, along with the region corresponding to large kinetic mixing, and the non-perturbative region where the corrections to gauge boson masses become larger than 50%.

Finally, an outstanding distinctive feature of the FPVDM scenario, especially when the new fermions are the partners of light quarks or leptons, is the possibility of multiple SM fermion production induced by the pair production of the new fermions, as illustrated in Fig. 26, where they cascade decay into H and V' bosons. This ultimately results in a striking final state featuring ten hard fermions that is worth analysing. This task is however left for future work.

3.3 Long-lived t-channel mediators

3.3.1 Generalities

In theories beyond the Standard Model, particles with macroscopic decay lengths, known as *long-lived particles* (LLPs),



975 Page 42 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 25 Feynman diagram representative of LHC processes relevant for an FPVDM scenario with top-partners (left), and regions of the parameter space excluded by the results of the LHC at the 95% confidence level (right). We consider exclusion limits projected onto the (m_{ID}, m_{VD}) plane for $m_T = 1600$ GeV, $m_H = 1000$ GeV, and

 $g_D=0.3$. Cross section isolines for hV' and V'V' production processes are also displayed, together with the regions related to non-perturbativity and large kinetic mixing (hatched areas). The blue regions correspond to non-physical scenarios where t_D is heavier than T or lighter than the top quark





Fig. 26 Representative Feynman diagram of a multi-fermion final state achievable with the FPVDM scenario

can arise due to phase-space suppression or small (effective) couplings that govern their decay. For a comprehensive



It turns out that the mass splitting $\Delta m \equiv M_Y - M_X$ not only impacts the lifetime of the LLP mediator but also affects the kinematics of the visible objects produced at the LHC from its decay. To address this, we categorise LLP scenarios into two main classes: those with relatively small mass split-



Eur. Phys. J. C (2025) 85:975 Page 43 of 96 975

tings ($\Delta m \ll M_X$, M_Y) and those without particularly small mass splittings. This distinction reflects not only the need for different LLP search strategies, but also corresponds to different DM genesis scenarios. In the CDFO regime, Δm is typically less than 10% of the DM mass, resulting in the emission of soft SM particles during the decay $Y \to X + \text{SM}$. These particles thus have relatively low momentum compared to the centre-of-mass energy of the process or the missing energy carried away by the DM particle. Consequently, such kinematics pose challenges for search strategies, as discussed below. Notably, the cosmologically viable parameter space for CDFO scenarios is constrained to mediator masses below a few TeV, exposing this entire parameter space to collider searches in the near future.

In contrast, the superWIMP and freeze-in production mechanisms, which belong to the class of non-thermalised DM scenarios, do not require small mass splittings. Instead, they often exhibit extremely large mass hierarchies between the Y and X states in the bulk of the cosmologically viable parameter space [202–204]. The DM particle in these scenarios can be as light as O(10) keV [204–206], effectively rendering it effectively massless with respect to a typical collider scale in large regions of the parameter space. On the other hand, testable mediator masses at the LHC are typically around the TeV scale. Nevertheless, the full cosmologically viable range of mediator masses extends far beyond the LHC reach, leaving only a small fraction of the parameter space accessible to collider searches.

In general, both the CDFO and freeze-in/superWIMP scenarios are largely unchallenged by conventional WIMP searches via indirect or direct detection (see Sect. 4) due to the highly suppressed interaction rates resulting from their very weak couplings. Consequently, these parameter space regions often represent the only regions in certain *t*-channel models that have not yet been excluded. This highlights the critical role of collider searches in probing these scenarios and underscores the strong motivation for performing LLP searches.

Experimental signatures featuring displaced objects, often accompanied by significant missing trasnverse energy, are key to detecting LLPs predicted by *t*-channel mediator DM models. Experimental results from collider experiments, such as ATLAS and CMS, impose valuable constraints on the mediator parameter space. Although no dedicated searches targeting *t*-channel mediator DM models have been performed to date, existing results can be recast to establish bounds. The current experimental constraints are detailed in the following sections. First, in Sect. 3.3.2, we review existing search strategies for LLPs. Next, in Sect. 3.3.3, we analyse the constraints originating from these searches within the freeze-in/superWIMP scenarios (Sect. 3.3.3.1) and the CDFO scenario (Sect. 3.3.3.2). In Sect. 3.3.4, we identify gaps in the current coverage of LLP searches, emphasising

the need to explore new signatures. Finally, in Sect. 3.3.5, we provide concluding remarks on the reinterpretability of the searches.

3.3.2 Current LLP searches

The LHC has now established a comprehensive programme dedicated to searches for LLPs. Many of these searches directly target DM production alongside objects originating from positions significantly displaced from the collision point. Moreover, other searches, particularly those selecting events with large missing transverse momentum (E_T^{miss}), may also indirectly probe the DM models considered in this report. In this section, we provide an overview of the signatures examined in these searches. We focus both on searches that explicitly involve missing transverse momentum for triggering and/or event selection, and on searches without explicit E_T^{miss} requirements that are inclusive enough to capture signals from the considered set of models. Table 7 lists a selection of recent LHC Run 2 searches representing these types, and we refer to [199] for a broader overview of LLP searches at the LHC.

Searches for heavy stable charged particles (HSCPs) are sensitive to DM models where the mediator carries electric charge and/or hadronises into charged hadrons, and then decays outside the tracker. These searches [207,208] rely on high ionisation losses (dE/dx) of heavy charged particles, and/or anomalous time-of-flight measurements between their production at the collision point and their arrival in the muon system. Furthermore, triggering often relies on calorimetric missing transverse energy, originating from either new physics particles produced in association with the charged LLP or the LLP itself if it decays outside the calorimeter. The reconstructed HSCP mass is then used to suppress backgrounds. HSCP searches are usually sensitive to models in which the electrically charged and/or coloured mediators have proper decay lengths larger than approximately 0.1 m [207]. Since they typically employ model-independent selection criteria, they are easily recastable using tabulated efficiencies like those provided in [219], and HSCP results have thus been reinterpreted not only within supersymmetric models [220,221], but also in frameworks featuring heavy resonances decaying into doubly charged LLPs [222].

In scenarios where charged LLPs decay within the tracker to mostly soft and/or invisible final states, disappearing track (DT) signatures arise. Disappearing track searches [209,210,223] identify short tracks with a few hits in the innermost tracking layers caused by charged particles that decay invisibly or to low-momentum states not reconstructed as tracks. Events are triggered by the large amount of $E_T^{\rm miss}$ generated by the invisible LLP decay, and the searches are usually sensitive to models with small mass splittings Δm and LLP decay lengths between 10 mm and 100 mm. ATLAS



975 Page 44 of 96 Eur. Phys. J. C (2025) 85:975

Table 7 Summary of recent LLP searches at the LHC Run 2 relevant for the signatures discussed in the text, and involving missing transverse energy together with heavy stable charged particles (HSCP), displaced

tracks (DT), dispalced vertices (DV), displaced leptons (DL) and displaced jets (DJ)

Signature	Analysis and reference	\mathcal{L} [fb $^{-1}$]	Decay length [mm]
HSCP	ATLAS-SUSY-2018-42 [207]	139	≥ 300
	CMS-EXO-18-002 [208]	101	$\gtrsim 1000$
DT	CMS-SUS-21-006 [209]	137	100-1000
	ATLAS-SUSY-2018-19 [210]	136	100-300
DV plus E_T^{miss}	CMS-EXO-22-020 [211]	137	0.1–20
•	ATLAS-SUSY-2016-08 [212]	32.8	4–300
Displaced soft tracks	ATLAS-SUSY-2020-04 [213]	140	0.1-10
DL	ATLAS-SUSY-2018-14 [214]	139	3–300
	CMS-EXO-18-003 [215]	113-118	0.1-100
DJ (calorimeter)	ATLAS-EXOT-2019-23 [216]	139	2000-4000
DJ (muon system)	CMS-EXO-21-008 [217]	138	3000-7000
	ATLAS-EXOT-2019-24 [218]	139	4000-8000

and CMS disappearing track searches therefore commonly target simplified supersymmetric scenarios where a long-lived charged wino or higgsino decays into a neutral particle and a low-momentum pion.

In contrast to HSCP and disappearing track analyses, searches for displaced vertices (DVs) focus on visible LLP decays within the tracker [211,212], and they can be very inclusive as long as the LLP decays to SM particles and invisible states. While displaced vertex searches often include stringent $E_T^{\rm miss}$ requirements for triggering reasons, their sensitivity further diminishes in scenarios with small Δm due to the additional requirement for high-momentum displaced objects. Moreover, some displaced vertex searches specifically target resonant displaced hadronic decays without missing energy, although these are less relevant for the (non-resonant) mediator models considered here. Since it is not always possible to determine the sensitivity of this last set of searches to those non-resonant mediator decays, they will not be considered in this report.

In addition, ATLAS has recently performed a search for mildly displaced soft tracks [213]. It relies on a trigger on the missing transverse energy and focuses on events with soft tracks (2 GeV $< p_T < 5$ GeV) exhibiting small displacements ($|d_0| < 10$ mm). This search specifically targets supersymmetric scenarios with compressed mass spectra, such as setups in which charginos decay to neutralinos and soft pions with small displacements. Despite being a challenging signal due to the presence of the soft objects, the search exploits the relatively small associated SM background rates (i.e. related to the production of soft and displaced pions). It thus allows to look for an LLP signal and hence cover previously existing gaps in typical LLP parameter spaces where $\Delta m < 1$ GeV and $c\tau \sim 0.1$ mm -1 cm.

Beyond $E_T^{\rm miss}$ -triggered searches, displaced leptons (DL) searches without any specific requirement on the missing energy are sufficiently inclusive to apply to long-lived mediators decaying to charged leptons and DM [214,215,224–229]. Such searches are typically triggered by either standard lepton triggers, or more specialised ones such as ATLAS's Run 3 displaced lepton triggers [230] (that focuses on the reconstruction of tracks with a large impact parameter) or CMS's data scouting dimuon stream with low p_T thresholds [231]. They are sensitive to a broad range of LLP decay lengths (0.1 mm to 10 m), and to spectra featuring large mass splittings, so that they can then be used to probe both supersymmetric models and scenarios featuring a more generic dark sector.

Finally, displaced jets (DJ) searches target LLPs decaying hadronically within different detector regions, such as the inner detector, the calorimeter, or the muon spectrometer [216–218,232–239]. These searches often rely on multijet triggers or lepton triggers (when jets and/or leptons are produced in association with the LLP), and in some cases, specialised triggers designed for the unique related calorimeter and muon spectrometer signatures. Displaced jet searches primarily probe dark sector scenarios involving LLPs decaying to SM fermions, although supersymmetric models are also considered, commonly in the *R*-parity-violating case.

3.3.3 Coverage of current searches

Figure 27 schematically illustrates the parameter space coverage of the LLP searches discussed in Sect. 3.3.2, along-side cosmological DM production scenarios. This coverage is presented in the plane defined by the LLP proper lifetime $c\tau$ and the associated mass splitting. The parameter space



Eur. Phys. J. C (2025) 85:975 Page 45 of 96 975




Fig. 27 Schematic representation of the coverage of LLP searches in the parameter space defined by the LLP proper lifetime and the associated spectrum mass splitting. Regions corresponding to various cosmological DM production scenarios are also indicated

of canonical freeze-out, which typically results in short lifetimes ($c\tau \lesssim 1$ mm), is associated with prompt signatures. In contrast, the parameter space relevant for CDFO scenarios, predominantly characterised by 1 mm $\lesssim c\tau \lesssim 1$ m and relatively small mass splittings, remains only partially covered by current searches. However, these regions can still be probed from multiple angles using various LLP and even prompt search strategies. Freeze-in/superWIMP scenarios, on the other hand, allow for larger mass splittings, while the mediator lifetime spans values similar to those in the CDFO scenario and extends well into the detector-stable regime. As a result, their parameter space is also exposed to searches for displaced leptons and jets, as well as for HSCP signatures.

One of the significant challenges from a theoretical perspective is the limited availability of recast analyses. In many instances, the absence of reinterpretation material or publicly available recasting codes makes it difficult to rigorously evaluate the coverage of specific searches for the models discussed in this report. We refer to Sect. 3.3.5 for a more detailed discussion on advancing reinterpretation methods. Consequently, the results presented here do not always reflect searches using the full integrated luminosity available. Currently, publicly available recasting and reinterpretation tools include ten LLP searches available from the 11precasting repository [245], three analyses implemented in the MadAnalysis 5 public analysis database [172], six analyses implemented in CheckMATE [246], and twelve analyses included in SModels [247–250]. It should be noted that the numbers above include some overlap between tools and represent only a subset of the LLP analyses conducted by the LHC collaborations. A summary of all the searches considered in this section is provided in Table 8.

3.3.3.1 Freeze-in/superWIMP regime

If the DM-mediator coupling λ is too small to thermalise dark matter in the early universe, the relic density as

observed in data can still be achieved through freeze-in and superWIMP production. In general, both production mechanisms coexist, with their relative contributions being model-dependent. For this reason, we consider them together. As the related cosmologically viable parameter space extends to very large mediator masses, up to around $10^9\,\text{GeV}$, the LHC can only probe a small fraction of it. Nonetheless, as we discuss below, current LLP search strategies offer good sensitivity.

We begin our study with an exploration of quark-philic models. The left panel of Fig. 28 shows the cosmologically viable parameter space for a class of scenarios where a topphilic scalar mediator couples to a Majorana DM particle (i.e. which corresponds to the S3M_tR class of simplified models discussed in Sect. 2.1). At large mediator and DM masses, superWIMP production dominates, with the solid black line marking where this mode alone saturates the observed relic density, $\Omega h^2 = 0.12$. To the right of this line, it would thus overclose the universe, which subsequently excludes that part of the parameter space. To its left, both freeze-in and super-WIMP production contribute to the relic density, with the black dashed contours indicating the relative superWIMP contribution. In addition, the figure includes two isolines in the mediator proper decay length, shown as grey dotted lines. They respectively span the prompt decay and detector-stable cases, covering decay lengths of 1 mm and 10 m respectively. For mediator masses around the TeV scale (i.e. within the reach of the LHC) and DM masses allowed by structure formation ($M_X \gtrsim 10 \text{ keV}$), mediator decay lengths typically exceed a few centimetres. For DM masses close to the exclusion limit, displaced vertex searches are the most relevant, and they exclude the violet-shaded region of the parameter space. For larger DM masses ($M_X \gtrsim 10 \text{ MeV}$), the mediator decays predominantly outside the tracker, making HSCP searches most effective. The corresponding excluded regions of the parameter space are indicated by the blueshaded excluded region of the parameter space. Both the DV and HSCP search strategies therefore restrict the mediator to be quite heavy.

Similar results are expected for other models of the S3M type where quarks of the first and second generation are involved, provided that $\Delta m \gg M_t$. Differences would only be due to the uniqueness of the decay modes of the displaced top quark. For instance, in top-philic models, an additional displacement arises from B-hadron decays, potentially complicating vertex reconstruction due to the extra associated tracks, and leptonic top decays could provide an alternative search avenue that is not available for models featuring quarks of the two lighter generations.


In the right panel of Fig. 28, we perform a similar analysis, but this time for a scenario with a scalar DM state and a fermionic mediator coupling to first-generation quarks. Here, the entire displayed region of the parameter space is consis-



975 Page 46 of 96 Eur. Phys. J. C (2025) 85:975

Table 8 Summary of the LHC searches reinterpreted to constrain LLP scenarios in this report. Those that are accessible through the publicly available llprecasting repository and the MadAnalysis 5, CheckMATE and SModelS programs are indicated with a check mark

Signature	Analysis & reference	\mathcal{L} [fb ⁻¹]	Reinterpretation code
HSCP	CMS-EXO-13-006 [219]	18.8	✓
	CMS-PAS-EXO-16-036 [240]	12.9	\checkmark
	ATLAS-SUSY-2016-32 [241]	36.1	\checkmark
	ATLAS-SUSY-2018-42 [207]	139	\checkmark
DT	ATLAS-SUSY-2016-06 [242]	36.1	\checkmark
	CMS-EXO-16-044 [243]	38	\checkmark
	CMS-EXO-19-010 [223]	101	\checkmark
DV plus E_T^{miss}	ATLAS-SUSY-2016-08 [212]	32.8	\checkmark
DV plus μ	ATLAS-SUSY-2018-33 [244]	136	✓
DJ plus $E_T^{\rm miss}$	CMS-EXO-19-001 [237]	137	\checkmark
DL	CMS-EXO-18-003 [215]	113–118	
	ATLAS-SUSY-2018-14 [214]	139	\checkmark
Monojet	CMS-EXO-20-004 [137]	137	\checkmark






Fig. 28 Constraints on *t*-channel DM models featuring quark-philic mediators in the freeze-in/superWIMP (non-thermalised) DM production regime, adapted from [204,251]. In the left panel, we display the viable region of the parameter space (i.e. with $\Omega h^2 = 0.12$) of a scenario with a top-philic scalar mediator and a Majorana DM state. Shaded regions indicate exclusions from DV (violet, [212]) and HSCP (blue,

[241]) searches, as well as from structure formation (purple) and BBN (red) (see details in Sect. 4). In the right panel, we consider instead a model where a fermionic mediator couples to first-generation quarks and scalar DM, and display again the constraints originating from DV (blue, [212]) and HSCP (purple, [240]) searches. In addition, the grey region represents setups with a hot DM candidate

tent with $\Omega h^2 = 0.12$, which is achieved through freeze-in production. In this case, as the mediator lifetime decreases, the rate for DM production increases, thus requiring smaller DM masses in order to reproduce the observed relic abundance. For $c\tau \lesssim 0.1$ m, the required DM mass even drops below 10 keV, which is excluded by structure formation constraints. Once again, displaced vertex and HSCP searches remain the most sensitive searches to this setup, constraining scenarios with mediator masses ranging up to $M_Y \sim 1.8$ TeV. Compared with the case of a scalar mediator and a fermionic

DM state, the bounds are slightly weaker because of the different production cross section for coloured fermions and scalars [251,252]. Moreover, as for Majorana DM, we expect our results to be qualitatively similar for the other quarkphilic models of this class.

We now move on with leptophilic scenarios that we illustrate by considering a setup in which a scalar mediator couples to Majorana DM and the right-handed muon, with an interaction strength relevant for freezing-in DM. Figure 29 shows contours of correct relic abundance for dif-



Eur. Phys. J. C (2025) 85:975 Page 47 of 96 975




Fig. 29 Isolines of constant relic density $\Omega h^2 = 0.12$ for different choices of the DM mass in a leptophilic Majorana DM model with a scalar mediator coupling to right-handed muons. We show constraints stemming from HSCP searches (red, [241]), ATLAS and CMS displaced lepton searches (green, [214,215]) and disappearing track searches (yellow, [223]). The plot is adapted from [252,253]

ferent choices of the DM mass, assuming a reheating temperature higher than the freeze-in scale. It turns out that low DM masses are excluded by structure formation constraints from Lyman- α measurements [204–206], and HSCP searches constrain large mediator lifetimes. For intermediate mediator lifetimes, displaced lepton searches can probe mediator masses up to approximately 500 GeV, while scenarios with a mediator decay length of O(1) metre are less constrained. In this case, a signature featuring a kinked charged track (made from the combination of the charged scalar track and the muon one) is the primary handle on the model. Reinterpreted disappearing track searches therefore provide the strongest bounds in this regime [252], due to the fact that there is no LHC analysis dedicated to kinked tracks to reinterpret.

The interpretation of the LLP searches as achieved so far depends on the assumption used for the cosmological history [251,252,254–256] (see also Sect. 4.2.6). Here, we have considered reheating temperatures $T_{\rm rh}$ much larger than the masses of the new physics states, which in particularly implies that $T_{\rm rh} \gg M_Y$. Consequently, the collider phenomenology is insensitive to $T_{\rm rh}$. However, for $T_{\rm rh} \lesssim M_Y$, larger couplings would be required to match the observed relic density, expanding the regions of the parameter space accessible to LLP searches at colliders. In such cases, freezein occurs during reheating, with entropy injection diluting the relic abundance. The dilution depends on the reheating potential $(V(\Phi) \sim \Phi^k)$ and on the fermionic or bosonic nature of

the reheating. These dependencies are illustrated in Fig. 30, in particular for early matter domination scenarios corresponding to k=2. Concretely, for $T_{\rm rh}\gg M_Y$, the region below the red solid line in the left panel of the figure would be excluded by structure formation constraints (i.e. imposing $M_X \gtrsim 10 \text{ keV}$) [204–206], thus limiting the relevance of LLP searches within the model. In case of a lower reheating temperatures $T_{\rm rh} < M_Y$, this bound shifts towards smaller decay lengths, revealing new relevant regions of the parameter space that are testable at colliders with displaced signatures, as visible from the left and right panels of the figure. The observation of an associated LLP signal corresponding to a region that would be excluded in the case of a standard cosmological history could thus provide valuable insights into the dynamics of the (inflationary) reheating phase for a given DM model. Conversely, for a given reheating scenario, the measurement of the mediator mass and lifetime would give an absolute upper limit on the reheating temperature consistent with freezing-in DM. In this context, a coherent analysis of specific inflation models (thus with a particular reheating potential) reveals an interesting interplay with cosmological data from Planck [254].

Alternatively, scenarios with a faster-than-standard expansion of the universe also require larger DM couplings to meet relic density constraints, enhancing again the importance of displaced object searches [257].

3.3.3.2 Conversion-driven freeze-out regime

The cosmologically viable regions of the parameter space for CDFO scenarios is characterised by a relatively small mass splitting Δm between the DM and mediator particles, below a few tens of GeV, and very weak DM couplings λ of the order of 10^{-6} [34]. This parameter space region borders the WIMP regime at larger Δm , where significantly stronger couplings are required to match the measured relic density. A striking feature of the CDFO scenario is its prediction of macroscopic mediator decay lengths, typically $1 \text{ mm} \leq c\tau \leq 1 \text{ m}$ for two-body decays, or even larger for three-body-suppressed or four-body-suppressed decays. As an illustrative example, Fig. 31 shows this behaviour for a scenario with a bottom-philic scalar mediator. The shortest lifetimes occur for large Δm near the boundary with the WIMP region, while lifetimes increase as Δm approaches the kinematic threshold of the two-body decay defined by $\Delta m = M_f$ (with f representing the SM fermion coupled to the mediator). In addition, for $\Delta m < M_f$, the mediator is typically detector-stable. The predicted lifetimes thus align well with the sensitivity range of LLP searches at the LHC, making this scenario particularly relevant for experimental exploration.

Within a quark-philic framework where the DM interacts with right-handed bottom quarks (f = b), different LLP searches are sensitive to specific ranges of media-



975 Page 48 of 96 Eur. Phys. J. C (2025) 85:975





Fig. 30 Isolines of constant reheating temperature $T_{\rm rh}$ expressed in the $(M_Y, c\tau)$ plane, for scenarios accounting for the observed DM relic abundance and a DM mass of 12 keV for a muon-philic Majorana DM model (left), and 10 keV for a top-philic model with scalar DM (right). In the left panel, adapted from [254], we study the impact of the reheating potential $V(\Phi) \sim \Phi^k$, and consider $T_{\rm rh} = 20$ GeV (blue), 100 GeV (black) and 10^4 GeV (red, with details of reheating being no

tor decay lengths. These include searches for heavy stable charged particles, disappearing tracks and displaced vertices, as well as searches for signals with missing transverse energy whose sensitivity could extend to longer lifetimes depending on their inclusiveness. Let us note that scenarios where f = u, d, s, c yield a very similar phenomenology, but with a different location of the two-body threshold $\Delta m = M_f$. In contrast, for f = t, the mediator is effectively detector-stable throughout the entire CDFO parameter space as $\Delta m < M_t$, making HSCP searches alone well-suited to probing this case [201]. Each class of searches covers different regions of the CDFO parameter space, as shown in Fig. 32 which illustrates the constraints that could be imposed on the mediator mass and decay length for scenarios compliant with $\Omega h^2 = 0.12$. Bounds from CMS disappearing track searches [223,243] (green), the ATLAS displaced vertex+ E_T^{miss} search [212] (red), and the CMS multijet+ E_T^{miss} search [137] (blue) are indicated in the figure, while those from HSCP searches are not shown. While these searches dominate at large $c\tau \gtrsim 10 \,\mathrm{m}$ or $\Delta m \lesssim M_b$, the corresponding region in parameter space is indeed omitted from the figure. In the detector-stable limit, scenarios with mediator masses up to approximately 1.3 TeV would however turn out to be constrained [207,208,241].

For smaller decay lengths with $\Delta m > M_b$, disappearing track searches (green areas in Fig. 32) become most sensitive, the mass splitting dependence being highlighted in the figure through grey dotted isolines. However, care must



longer relevant for the considered mediator masses). Solid lines apply to k=2, while dashed and dot-dashed lines illustrate fermionic and bosonic reheating scenarios for k=4 potentials respectively. Various constraints from LLP searches are additionally shown as grey-shaded regions. In the right panel, adapted from [252], we consider k=2, vary $T_{\rm rh}$, and show collider constraints from DJ (blue, [237]), DV (green, [212,244]) and DL (dark green, [214]) searches




Fig. 31 Representation of a typical cosmologically viable parameter space for t-channel DM scenarios featuring a bottom-philic scalar mediator [258], adapted from [259]. The black line separates the WIMP and CDFO regimes, where in the latter the mediator decay length is displayed via the colour code. We recall that in the WIMP regime decays are prompt

be taken with the reinterpretation of the experimental disappearing track search results, focusing on colourless chargino decays to DM and an ultra-soft pion, to the CDFO signal. This requires accounting for differences originating from mediator hadronisation into neutral or charged R-hadrons, and from the specific properties of the b-jet emerging from the mediator decays (that is generally not ultra-soft for $\Delta m \gtrsim 10 \, \text{GeV}$). To solve this issue, we have made use of the implementation



Eur. Phys. J. C (2025) 85:975 Page 49 of 96 975




Fig. 32 LHC constraints on the parameter space of a scenario featuring a bottom-philic mediator and DM production in the CDFO regime [259]. We display constraints from LHC searches for disappearing tracks (green, [223]), displaced vertices (red, [212]) and monojet searches (blue, [137])

available in MadAnalysis 5 [172] and enforced the presence of a charged R-hadron and $\Delta R(Y,b) > 0.2$ in order to avoid decay configurations where the b-jet is aligned with the track candidate. For decay lengths below approximately 1 cm, displaced vertex and monojet searches provide the best sensitivity (red in Fig. 32). However, signal jets are usually soft and with a low invariant mass, yielding small signal efficiencies. Consequently, the alternative signature where an extra hard jet would stem from initial state radiation could be tested by conventional monojet searches (blue in Fig. 32). As it is not clear that events featuring displaced jets would be vetoed, we conservatively rejected, in our analysis, events exhibiting jets with a displacement larger than 2 mm and a transverse momentum $p_T > 20$ GeV.

Despite all these efforts, the large mediator production cross section for $M_Y \lesssim 500\,\text{GeV}$ and the potentially prominent LLP signatures, a substantial gap remains in the coverage of the parameter space for intermediate to small lifetimes (mm $\lesssim c\tau \lesssim$ few cm), revealing an uncharted regime that we discuss further in Sect. 3.3.4.

We continue our discussion with leptophilic scenarios, that we illustrate with a setup involving a Majorana DM candidate coupled to a scalar slepton-like mediator [260]. For an interaction strength $\lambda \lesssim 10^{-6}$, neglecting bound state formation effects [258,261] and ignoring Higgs-mediator couplings, the measured DM relic abundance can be achieved through CDFO production for mediator masses below 200 GeV and mass splittings $\Delta m < 2.6\,\text{GeV}$. While such a setup is already stringently constrained by searches at the LHC, introducing a non-zero quartic coupling between the scalar mediator and the Higgs boson significantly enlarges the viable regions in the model parameter space. In Fig. 33, we consider such a





Fig. 33 LHC constraints on the parameter space of a scenario featuring a muon-philic Majorana DM state and a scalar mediator, where the observed relic density is achieved within the CDFO regime [260]. The blue contours correspond to isolines of correct relic abundance for different values of the mass splitting Δm , and we display constraints from LHC searches for HSCP (red, [219,240]), and disappearing tracks (purple and green, [242,243]). For this illustrative case a quartic coupling of $\lambda_H=0.1$ has been assumed between the scalar mediator and the SM Higgs boson

quartic coupling $\lambda_H=0.1$, and illustrate the constraints on the parameter space of a model including interactions with the right-handed muon. The results are as usual given in terms of the mediator lifetime and mass. The upper region of the plot corresponds to long-lived mediators with decay lengths of a few centimetres or more, or equivalently small values of the DM-mediator coupling together with a very compressed mass spectrum (as indicated by the blue contours). For $\lambda_H=0.1$, the CDFO regime extends to mediator masses up to 300 GeV, and necessitates mass splittings $\Delta m < 3$ GeV. We include in the figure the complementary constraints emerging from early LHC Run 2 searches for HSCP [219,240] (red) and disappearing tracks [242,243] (green and purple), that should get a little bit stronger by relying on more recent analyses like [223,241].

Similar to the quark-philic case, we observe a significant lack of experimental coverage in the region with small lifetimes, where the considered scenario predicts soft displaced leptons. Here, the small mass splitting renders the emitted leptons too soft to be efficiently reconstructed in existing displaced lepton searches. This contrasts sharply with leptophilic scenarios involving freezing-in DM in alternative cosmologies (see Fig. 30), where displaced lepton searches are highly effective. Nevertheless, it would be worthwhile to investigate whether recent DL searches like [262] could probe this particular region of the CDFO parameter space. We further discuss this issue in Sect. 3.3.4.



975 Page 50 of 96 Eur. Phys. J. C (2025) 85:975






Fig. 34 Constraints on CDFO flavoured DM scenarios, shown as a function of the proper lifetime and mass of the mediator (left, adapted from [82]) and the heavier dark states (right, adapted from [77]). In the left panel, the blue and red lines represent slices in the parameter space of a leptophilic two-flavour model that simultaneously account for the observed DM abundance and successful leptogenesis (excluding the regions shown with faint colours). The solid shaded areas indicate

current LHC constraints, while the transparent light blue areas illustrate projected sensitivities from dedicated searches. In the right panel, cosmologically viable points are shown for a three-flavour quark-philic model. The green and blue points represent the decay lengths of the heavier DM multiplet states X_2 and X_1 , respectively, as functions of the mass of the heaviest state X_1

CDFO realisations can also be achieved within nonminimal models [77,82,263]. A particularly intriguing example is flavoured DM scenarios, where the DM state X = (X_1, X_2, X_3) consists in a flavour multiplet whose lightest mass eigenstate X_3 serves as the DM candidate. Within these models, the CDFO mechanism has been explored for both quark-philic DM [77] and leptophilic DM [82]. Notably, these frameworks offer the appealing prospect of addressing DM and baryogenesis within a unified framework, specifically through conversion-driven leptogenesis which provides additional motivation for experimental searches. In practice, CDFO DM production can manifest at colliders in multiple ways, potentially leading to signatures from either pairproduced long-lived mediators or from the production of long-lived heavier dark multiplet states X_1 or X_2 . In the former case, the signatures resemble those of minimal models but with quantitative differences in the preferred lifetimes due to the presence of the additional states. For instance, in conversion-driven leptogenesis scenarios, the preferred decay lengths range from a few centimetres to millimetres, with $\Delta m \lesssim 5$ GeV (see the left panel of Fig. 34). Moreover, these models predict soft, mildly displaced leptons as a challenging but intriguing target for LHC searches [82].

Alternatively, if the coupling matrix λ exhibits significant hierarchies, a heavier multiplet state X_i can interact more

sizeably with the SM and become the LLP produced at the LHC. In such cases, an electrically neutral LLP would not generate detectable tracks but could lead to displaced vertices. Interestingly, the kinematic suppression inherent to the three-body decay via an off-shell mediator can result in large lifetimes for these states. The right panel of Fig. 34 illustrates the dependence of the proper decay lengths of the heavier DM multiplet states X_2 (green) and X_3 (blue) on the mass of the heaviest state in a quark-philic setup [77]. The large lifetimes of these neutral X_i states allow them to escape LHC detectors, making them ideal candidates for detection at dedicated experiments such as MATHUSLA [264].

3.3.4 Gaps in coverage

The scenarios discussed in Sect. 3.3.3 illustrate the importance of LLP searches in testing various DM mechanisms. Despite the broad scope of the LLP search programme at the LHC, specific regions of the various parameter spaces considered remain uncovered by current search strategies. In particular, within the CDFO scenario, configurations with $c\tau \lesssim 1$ cm ($\Delta m \gtrsim 10$ –30 GeV) are not covered because they fall outside the lifetime range addressed by disappearing track searches, and/or the LLP visible decay products are too soft to be detected in conventional displaced vertex



Eur. Phys. J. C (2025) 85:975 Page 51 of 96 975




Fig. 35 In the left panel, we show the two-dimensional distribution in the number of displaced vertices as a function of their invariant mass $m_{\rm DV}$ and their number of tracks $n_{\rm trk}$, for a quark-philic CDFO benchmark scenario with $M_Y = 481$ GeV, $\Delta m = 31$ GeV and $\lambda = 3.9 \times 10^{-7}$

[259]. In the right panel, we examine how the parameter space region excluded by the ATLAS DV search depends on the specific requirements on the displaced vertices, as detailed in the text

searches. This gap in coverage is approximately represented by the white region shown in Fig. 27.

The quark-philic scenario discussed above provides an example of this gap in coverage and how it could be addressed with minor modifications to existing search strategies. For instance, we could consider the ATLAS displaced vertex search [212] (ATLAS-SUSY-2016-08), which relies on a E_T^{miss} trigger and is sensitive to decay lengths between 4 mm and 30 cm. The signal region defined in this search requires at least one displaced vertex with five or more tracks and an invariant mass exceeding 10 GeV. In the left panel of Fig. 35, we present the distribution of the number of displaced vertices as a function of the number of tracks n_{trk} associated with it and of the invariant mass oof the vertex m_{DV} , in the framework of a quark-philic CDFO benchmark model with $M_X = 450$ GeV, $\Delta m = 31$ GeV, and $c\tau = 0.85$ cm. Most vertices have $3 < n_{trk} < 9$ and $m_{DV} < 6$ GeV, making them too soft to satisfy the ATLAS analysis requirements. For this benchmark, the m_{DV} cut reduces the signal efficiency from approximately 6% to 0.1%, therefore significantly diminishing the search sensitivity. Although the high m_{DV} threshold effectively suppresses SM backgrounds, it could likely be lowered without a substantial increase in background contamination. Indeed, the ATLAS search observed zero vertices with $n_{\text{trk}} > 5$ and $m_{\text{DV}} > 4$ GeV so that lowering the $m_{\rm DV}$ requirement, for instance to 5 GeV, could still maintain an effective SM background suppression. Assuming that the SM background associated with the modified selection $m_{\rm DV} > 5~{\rm GeV}$ remains negligible (i.e. fewer than one vertex), we estimate in the right panel of Fig. 35 the sensitivity improvement from reducing the invariant mass threshold from 10 GeV (red exclusion curve) to 5 GeV (orange exclusion curves), as further detailed in [259]. This demonstrates that a significant portion of parameter space becomes accessible when mildly soft displaced vertices are included, this region being in addition not covered by other LLP searches.

Similarly, the strategy employed in the mildly displaced soft track search from [213] could also address a similar gap in the leptophilic CDFO case. Here, soft and mildly displaced leptons arise for moderate Δm values so that, as illustrated in Fig. 33, scenarios with $\Delta m > 3$ GeV are currently unconstrained. Comparable gaps in coverage exist in non-minimal models as well, as seen in the right panel of Fig. 34 and [82] for the leptophilic case, particularly in conversion-driven leptogenesis scenarios, and [77] for the quark-philic case.

Beyond soft displaced objects, additional signatures could be explored to enhance background suppression. These include more sophisticated E_T^{miss} -based discriminators, timing information from the tracker or calorimeters, and the combination of the prompt track of the mediator with the displaced track of its decay products (yielding so-called 'kinked tracks'). In Fig. 36, we present some key distributions relevant for designing such searches, using the quarkphilic model discussed earlier as an example. The upper-left panel shows the track length distributions for two benchmarks with $\Delta m = 31 \text{ GeV}$ ($c\tau \simeq 1 \text{ cm}$) and $\Delta m = 8 \text{ GeV}$ $(c\tau \simeq 30 \text{ cm})$. The colour-charged mediator, which forms charged R-hadrons with SM quarks, produces highly ionising tracks. In addition, those decaying within the tracker but traversing multiple inner layers give rise to disappearing or kinked tracks, depending on the softness of the decay products. Since $\Delta m < 40$ GeV is required by the CDFO mechanism, the displaced jets are in general relatively soft so that hard jets from initial state radiation may be necessary



975 Page 52 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 36 Kinematic properties of an LLP signal relevant to design new LLP searches at the 13.6 TeV LHC, emerging from a typical quark-philic CDFO scenario. We consider the distribution in the charged R-hadron decay length (upper left) for a benchmark with $c\tau \simeq 1$ cm (blue) and 30 cm (orange), the p_T distributions of prompt (blue) and displaced

(orange) jets with $p_T(j) > 25$ GeV and $|\eta(j)| < 5$ (upper right) as well as the associated prompt (blue) and displaced (orange) jet multiplicity spectrum (lower left), and the E_T^{miss} distribution (lower right)

200

 E_T^{miss} (GeV)

200

 $p_T(j_1)$ (GeV)

250

Prompt jets

300

300

350

400

400

Displaced jets

to satisfy trigger requirements. We refer to [259] for further discussion.

3.3.5 Advancing reinterpretation methods

We conclude this chapter with some remarks on the material necessary for the reinterpretation of LLP searches (see also [265] and chapter 6 of [199]).

As emphasised in Sect. 3.3.3, the scarcity of validated recast analyses is a major obstacle to assessing the current coverage of the LLP parameter spaces and to identifying the associated gaps. Therefore, it is pertinent to discuss potential measures to improve the situation, which requires adequate resources and collaboration. Reinterpretation tools are typically developed by small teams, often consisting of only a few researchers. In contrast, each experimental analysis is conducted by a dedicated team of analysts. Consequently, a single 'reinterpreter' must handle the implementation and validation of numerous experimental analyses, frequently resulting in delays between the publication of an analysis and its incorporation into public reinterpretation frameworks. Consequently, by the time at which reinterpretation tools are updated, obtaining further information about the original analysis can become challenging, if not impossible. A more direct interaction between theorists and experimentalists could help alleviate these issues.

Another critical factor is access to data from experimental collaborations, which is essential for proper implementation and validation. For instance, implementing analyses into SModelS requires acceptance times efficiency maps for each signal region for pure simplified models, given as a function of the simplified-model parameters. These can be either provided directly by the experimental collaborations, or derived via simulation-based recasting. Full-fledged recast codes, on the other hand, demand clear and unambiguous object definitions, a step-by-step description of the analysis logic, and truth-level versus reconstruction-level efficiencies. Moreover, recasting LLP searches presents additional challenges due to their reliance on non-standard objects, low-



Eur. Phys. J. C (2025) 85:975 Page 53 of 96 975

level detector inputs not reproducible in fast simulation, and the use of machine learning techniques for optimisation. 11

Significant strides have been made to address these challenges, with several recent examples demonstrating good practices. First, the ATLAS Run 2 search for LLPs decaying into hadronic jets in the calorimeter [216] provided sixdimensional efficiencies for an event to enter Region A (of the ABCD method). These efficiencies parametrise the output of a sequence of boosted decision trees and neural networks as functions of the LLP kinematics, decay type and decay position. An example code illustrating their usage is also included. Second, the CMS Run 3 search for LLPs decaying to a pair of muons [228] published truth-level signal efficiencies for the considered dimuon categories. These efficiencies are functions of the minimum muon transverse momentum p_T and displacement d_0 in three intervals of the generated transverse decay length L_{xy} . The collaboration also provided detailed reinterpretation instructions. As a last example, the updated ATLAS Run 2 search for pairs of neutral LLPs in events with displaced jets and leptons [233] (extending [216]) went a significant step further. It included a BDT trained to compute the overall selection probability in the ABCD plane, using truth-level input of the decay position (L_{xy}, L_z) , kinematics $(p_T, \eta, E_T^{\text{miss}})$, and Child ID. This approach effectively serves as a surrogate model for the complete analysis, incorporating detector effects. The trained BDTs were provided as pickle files alongside example code demonstrating their use. In each of the examples above, the accuracy and range of validity were thoroughly documented, making such material valuable for advancing recasting efforts.

Finally, reliable reinterpretation efforts also require detailed information about the probability models underlying the analyses [267]. This is particularly crucial when combining signal regions in the statistical interpretation or performing combined fits to signal and control regions. Since 2019, ATLAS has been providing comprehensive HistFactory models in JSON format [268], mainly for supersymmetry and top quark analyses. More recently, CMS released its Combine software [269] along with data cards detailing early Higgs boson measurements. These developments represent significant progress, and we hope to see similar initiatives applied to LLP searches.

4 Cosmology

Contributions from C. Arina, M.J. Baker, M. Becker, A. Belyaev, M. Blanke, L.M. Carpenter, E. Copello, B. Fuks, M. Garny, J. Harz, J. Heisig A. Ibarra, S. Khalil, L. Lopez-

Honorez, K. Mohan, A. Moreno Briceño, T. Murphy, L. Panizzi, D. Sengupta, W. Shepherd, A. Thamm

4.1 Generalities

In this section, we explore the phenomenology of dark matter in the *t*-channel models introduced in Sect. 2, which encompass both minimal and non-minimal realisations. We outline the constraints and requirements that a viable dark matter candidate must satisfy from cosmological and astroparticle perspectives. We then assess the primary impact of these constraints on the parameter space of the model, providing explicit examples from the existing literature on *t*-channel models. Among minimal scenarios, a Majorana dark matter candidate with a scalar mediator is the most extensively studied case, particularly when it couples to right-handed third-generation quarks. As a benchmark, we will thus frequently refer to the S3M_tR model.

This section is organised as follows. In Sect. 4.2, we discuss the mechanisms available to achieve the correct relic density in the early universe. Section 4.3 addresses the primary signatures for direct and indirect dark matter searches, as well as the constraints from early-universe physics. Finally, in Sect. 4.4, we examine the associated impact on the parameter space of selected benchmark models.

This section serves as a reference for understanding the complementarity between cosmological and astrophysical searches for dark matter, and collider-based searches. Each subsection is designed to convey the key concepts while providing references for readers seeking a deeper understanding of the topics discussed. For a more general overview of dark matter constraints, models, and features, we direct readers to recent reviews [4,270].

4.2 Dark matter relic density

In *t*-channel mediator models, the DM relic density can be achieved through a variety of production mechanisms, as already sketched in Sect. 3.3.1. These include (by decreasing DM interaction strength) canonical WIMP freeze-out (FO) [271,272], conversion-driven freeze-out (CDFO) [34,35], freeze-in (FI) [36,37,200], and superWIMP (SW) [38,39] production. Describing these mechanisms requires at least two coupled Boltzmann evolution equations for the DM *X* and the mediator *Y*. A priori, these equations should be solved at the level of the distribution functions taking the form:

$$\frac{\mathrm{d}f_i}{\mathrm{d}t} = C(f_i, f_j),\tag{34}$$

where d/dt is the total time derivative and $C(f_i, f_j)$ is the collision operator. The distributions f_i and f_j are the distribution functions for particle species i and j, which, for the



Machine-learning-based analyses have long been considered impossible to recast; see, however, [266] for a discussion of recent progress and guidelines for reusable machine-learning models in LHC analyses.

975 Page 54 of 96 Eur. Phys. J. C (2025) 85:975

purpose of DM relic abundance calculations, can be considered as functions of proper time and of the particle momenta. Non standard cosmology (early matter-dominated era, see e.g. [252,254,255,273–277]) will essentially affect the left-hand side of this equation through a specific Hubble expansion rate, while particle physics processes affect its right-hand side. Unless stated otherwise, a standard cosmological history is assumed, with non-standard cosmology being discussed briefly in Sect. 4.2.6.

In standard cosmology, the universe is radiation-dominated during DM production, with energy density $\rho = g_* \frac{\pi}{30} T^4$ and entropy density $s = g_{*S} \frac{\pi}{45} T^3$ satisfying $sa^3 = \text{constant}$. In these expressions, g_* and g_{*S} are the conventional effective numbers of relativistic degrees of freedom. Assuming a Boltzmann distribution for all particles involved in the DM production process and negligible variation of the number of relativistic degrees of freedom during DM production, the evolution equations can be integrated over the momentum of the particle species i, yielding [278]:

$$\frac{\mathrm{d}Y_{i}}{\mathrm{d}x} = \frac{1}{3Hs^{2}} \frac{\mathrm{d}s}{\mathrm{d}x} \sum_{ij} \times \left[\gamma_{ij} \left(\frac{Y_{i}Y_{j}}{Y_{i}^{\mathrm{eq}}Y_{j}^{\mathrm{eq}}} - 1 \right) - \gamma_{j \to i} \left(\frac{Y_{j}}{Y_{j}^{\mathrm{eq}}} - \frac{Y_{i}}{Y_{i}^{\mathrm{eq}}} \right) \right]. \tag{35}$$

Here the quantities γ_{ij} and $\gamma_{i\rightarrow i}$ represent the reaction densities (described below), $Y_i = n_i/s$ are the comoving number density for the particle species i = X, Y with n_i being the number density, and $Y_i^{\text{eq}} = n_i^{\text{eq}}/s$ are the associated equilibrium comoving density with n_i^{eq} being the i^{th} species number density assuming kinetic equilibrium and zero chemical potential μ_i . If kinetic equilibrium is maintained while the dark matter chemically decouples, as typical of freeze-out scenarios, $n_i/n_i^{\text{eq}} = \exp(-\mu_i/T)$. Moreover, (35) relies on a standard and convenient practice that introduces a dimensionless time variable x inversely proportional to the bath temperature T, thus defined by $x = M_X/T$ (M_Y/T) for freeze-out (FI and SW) scenarios. Finally, we remind that H is the Hubble rate and that the prefactor in (35) simplifies under the assumption of approximately constant g_{*S} , for which $ds/dx \simeq -3 s/x$. This assumption will be implicit in the following (for simplicity), unless specified otherwise.

In order to write (35), we have assumed that all particles involved in (co-)annihilation and conversion processes are in kinetic equilibrium at the time of DM production. For non-thermal DM production discussed in Sects. 4.2.3 and 4.2.4, the relevant relation (42) can be obtained from (35) without making any assumptions on the DM distribution, but after instead enforcing a negligible initial abundance for X and suppressed couplings to Y. Equation (35) is thus also perfectly suitable for evaluating the DM comoving number den-

sity Y_X in FI and SW scenarios for which the loss terms for the dark matter particle X are neglected. ¹² In addition, the semi-classical Boltzmann equation (35) neglects various quantum and thermal effects, such as quantum coherence, quantum statistics, screening, multiple scatterings, and particle interactions with coherent condensates. A systematic framework for deriving quantum kinetic equations that incorporate these effects in relic density computations is provided by the Schwinger-Keldysh formalism (see, e.g., [281–287], as well as [288-291] for related computations in equilibrium and [286] for a lattice-based approach). In particular for freeze-in scenarios quantum and thermal effects can alter the relic density significantly. This was demonstrated in [289] for a Majorana DM model and in [284] for a scalar DM model, respectively. The latter work, which employs nonequilibrium quantum field theory from first principles, quantifies the expected corrections to various approaches commonly used in the literature (freeze-in with only decays, including scattering with and without thermal masses, hard thermal loop approximation, etc.) and provides recommendations for accurate results in phenomenological studies. For freeze-out scenarios and in particular for co-annihilating configurations, significant impact is expected from the Sommerfeld effect and bound state formation, as discussed in Sect. 4.2.5. However, only small corrections are expected due to thermal effects, as detailed in the context of dark matter annihilation in, e.g., [281,292] and Sommerfeld and bound state formation in, e.g., [288,293–298].

The γ_{ij} quantities appearing in (35) represent the reaction densities for (co-)annihilation of the dark species i and j, while $\gamma_{i \to j} = \gamma_{i \to jk}^{\text{dec}} + \gamma_{ik \to jl}^{\text{scat}}$ govern the conversion processes of the dark species i to j through (inverse-)decays or scatterings off SM particles k, l. For l and l and l processes, the reaction densities respectively take the forms:

$$\gamma_{ij\to kl} = \int d\phi_i \, d\phi_j \, f_i^{\text{eq}}(t, p_i) \, f_j^{\text{eq}}(t, p_j)
\int d\phi_k \, d\phi_l \, (2\pi)^4 \delta^4(p_i + p_j - p_k - p_l) \, \left| \mathcal{M}_{ij\to kl} \right|^2,
\gamma_{i\to jk} = \int d\phi_i \, d\phi_j \, d\phi_k \, f_i^{\text{eq}}(t, p_i) \, (2\pi)^4 \delta^4(p_i - p_j - p_k)
\left| \mathcal{M}_{i\to jk} \right|^2,$$
(36)

where f_i and p_i denote the i^{th} species distribution function and momentum, while $|\mathcal{M}|^2$ represents the relevant squared scattering amplitude averaged over the initial-state *and* final-state degrees of freedom, and $\mathrm{d}\phi_i = g_i \mathrm{d}^3 p_i/(2E_i(2\pi)^3)$ stands for a one-body phase-space element. These reaction densities can be expressed in terms of thermally averaged cross sections and decay rates of the particle species i in its



¹² For further discussion on kinetic decoupling, see e.g. [34,279,280].

Eur. Phys. J. C (2025) 85:975 Page 55 of 96 975




Fig. 37 Schematic plot showing the dependence of the relic density, Ωh^2 , on the DM coupling λ for a scenario with a relatively small mass splitting between the DM particle X and the t-channel mediator Y. The blue band indicates the region for which $\Omega h^2 = 0.12$ and depends on the new physics masses M_X and M_Y . The numbers for λ indicate rough orders of magnitude, and are model-dependent. The four characteristic production regimes are further discussed in Sects. 4.2.1, 4.2.2, 4.2.3 and 4.2.4, respectively, in the order of decreasing coupling strength λ . This leads to a distinct phenomenology in each case

rest frame. For instance, for (co-)annihilation processes and for the $i \rightarrow jk$ decay contribution to a conversion processes,

$$\gamma_{ij} = n_i^{\text{eq}} n_j^{\text{eq}} \langle \sigma_{ij} v_{ij} \rangle,
\gamma_{i \to j}^{\text{dec}} = n_i^{\text{eq}} \langle \Gamma_{i \to jk} \rangle = n_i^{\text{eq}} \Gamma_{i \to jk} \frac{K_1(M_i/T)}{K_2(M_i/T)}.$$
(37)

Here $\langle \sigma_{ij} v \rangle$ stands for the thermally averaged (co-) annihilation cross section entering the freeze-out relic abundance calculations, and $\langle \Gamma_{i \to jk} \rangle$ is the thermally averaged decay rate related to the process $i \to jk$ in the rest frame of particle i (with $M_i > M_j + M_k$). Moreover, K_1 and K_2 are the modified Bessel functions of the second kind.

Equation (35) represents general Boltzmann equations that can be simplified according to the dominant processes and the size of the DM coupling to the mediator. Depending on this coupling, certain terms can indeed be neglected or integrated out. Moreover, the size of the coupling strength largely determines the type of DM production mechanism in the early universe, as schematically depicted in Fig. 37 (see [260] for an explicit example). From left to right in the plot, very feebly interacting DM particles achieve their relic density via the SW mechanism. As the coupling λ increases, FI becomes active, followed by conversion-driven freeze-out, and finally standard freeze-out mechanism for couplings of O(1).

Returning to (35), during freeze-out, the DM particle X can be assumed to remain in thermal and chemical equilibrium with the mediator Y, provided that the coupling is sufficiently strong. In this case, the conversion processes occur much faster than the Hubble rate, enforcing $n_X/n_Y = n_X^{eq}/n_Y^{eq}$. Under these conditions, the coupled set of equations (35) reduces to a single equation describing the evolution of the DM abundance, driven by an effective annihilation cross section that accounts for all co-annihilation processes

of *X* and *Y*, weighted by the relevant Boltzmann suppression factors. For more details, see Sect. 4.2.1 and references such as [271,272,278]. Conversely, if the conversion processes between *X* and *Y* are inefficient, the full coupled set of Boltzmann equations must be solved. This situation is discussed in Sect. 4.2.2.

The four primary DM production mechanisms are now described in the following subsections. For further discussion, see also [299,300].

4.2.1 Canonical freeze-out

DM production via standard freeze-out occurs if the interaction between the DM and the SM is strong enough to establish chemical equilibrium between the dark sector and the SM at early times. If chemical equilibrium is also maintained within the dark sector through conversion processes among its particle species, we can approximately write

$$\frac{Y_j}{Y_i^{\text{eq}}} = \frac{Y_i}{Y_i^{\text{eq}}},\tag{38}$$

which allows us to simplify (35). Under this assumption, the evolution of the total comoving number density $\tilde{Y} = \sum_i Y_i$ is governed by

$$\frac{\mathrm{d}\tilde{Y}}{\mathrm{d}x} = \sum_{i} \frac{\mathrm{d}Y_{i}}{\mathrm{d}x} = -\frac{cg_{\star,\mathrm{eff}}^{\frac{1}{2}}}{x^{2}} \left\langle \sigma_{\mathrm{eff}}v \right\rangle \left(\tilde{Y}^{2} - \left(\tilde{Y}^{\mathrm{eq}}\right)^{2}\right), \quad (39)$$

where $c = \sqrt{\pi/45} \, M_{\rm Pl} \, M_X$. In this last expression, the effective annihilation cross section and number of relativistic degrees of freedom are defined by

$$g_{\star,\text{eff}}^{\frac{1}{2}} = \frac{g_{\star S}}{\sqrt{g_{\star}}} \left(1 + \frac{T}{3g_{\star S}} \frac{dg_{\star S}}{dT} \right),$$

$$\langle \sigma_{\text{eff}} v \rangle = \sum_{i,j} \langle \sigma_{ij} v \rangle \frac{Y_i^{\text{eq}} Y_j^{\text{eq}}}{\tilde{Y}^{\text{eq}} \tilde{Y}^{\text{eq}}},$$
(40)

where $g_{\star S}$ is the effective number of relativistic degrees of freedom for entropy density. Here, we assume that the distribution functions of all particles can be expressed as a rescaling of their equilibrium distributions, $f_i = \alpha_i f_i^{\rm eq}$, and that these equilibrium distributions follow a Boltzmann distribution, i.e. $f_i^{\rm eq} = \exp(-E_i/T)$. The annihilation and coannihilation processes inherent to the *t*-channel DM models considered and contributing to $\langle \sigma_{\rm eff} v \rangle$ are depicted in Fig. 38.

Since all dark sector particles heavier than the DM candidate X are assumed to eventually decay into X, the DM relic density is determined by the solution to (39). Inspecting the effective annihilation cross section (40) reveals that annihilations of heavier dark sector particles Y can significantly contribute to the total cross section if $M_Y \lesssim 1.2 \, M_X$, a scenario commonly referred to as a co-annihilation.



975 Page 56 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 38 Representative LO Feynman diagrams contributing to the DM annihilation cross section $\langle \sigma_{\rm eff} v \rangle$. They include t-channel DM annihilations (top left), XY co-annihilations (top central and right), and mediator pair-annihilations (bottom row) with the blob accounting for channels yielding SM electroweak bosons or gluons. The final state q represents any quark flavour, and may include a sum over flavours depending on the

model. For leptophilic models, q can be replaced by a lepton ℓ , and for models with self-conjugate DM, additional YY annihilation diagrams must be added. Moreover, in cases where the S-wave contribution to $\langle \sigma_{\rm ann} v \rangle (X\bar{X} \to q\bar{q})$ is helicity suppressed, NLO corrections become relevant (see the diagrams in Fig. 40)

Within the standard freeze-out mechanism, the final DM relic density Ωh^2 exhibits the characteristic *increasing* trend as the coupling strength λ is *reduced*, as shown on the right-hand side of Fig. 37. Within the co-annihilation regime, this trend levels off when the coupling λ between the DM and the mediator becomes smaller than the gauge couplings relevant for the *t*-channel mediator. In this regime, DM freeze-out is dominated by efficient conversion to the mediator and subsequent freeze-out of mediator pair-annihilations, resulting in the plateau observed in Fig. 37. When λ becomes even smaller, the system transitions into the domain of conversion-driven freeze-out.

The canonical freeze-out picture can be affected by non-perturbative effects, the so-called Sommerfeld effect, as well as the formation and successive decay of bound states, as discussed in detail in Sect. 4.2.5. Specifically, for a mediator that is both coloured and electrically charged, long-range effects via gluon exchange impact the expected viable regions of the parameter space significantly [301,302], as found in the case of exchanges of relatively light scalars [303,304].

4.2.2 Conversion-driven freeze-out

The assumption of chemical equilibrium among the coannihilating partners during freeze-out holds if conversions between X and Y, such as $Y \to AX$ decays or $YB \to XA$ scatterings along with their inverse processes (with A and B denoting SM particles), remain fully efficient, i.e. if $\Gamma_{\text{conv}} \gg H$. This is generally true if the DM coupling λ and the mediator gauge interaction strengths are of comparable magnitude, and if A and B are comparably light SM particles. In such cases, conversion rates are not suppressed, while annihilation processes initiate chemical decoupling due to their Boltzmann suppression from the initial-state heavy particles. As λ decreases, two effects occur. First, direct annihilation processes involving λ become negligible relative to

mediator pair annihilation, while conversions remain efficient. In this regime, the relic density becomes independent of λ , corresponding to the plateau in Fig. 37. However, further reduction of λ eventually renders conversions inefficient, initiating chemical decoupling at $\Gamma_{\rm conv} \sim H$. In this conversion-driven freeze-out scenario [34,35], the relic density rapidly increases with decreasing λ , with typical values of $\lambda \sim 10^{-6}$ for couplings to light quarks, while for topphilic scenarios slightly larger couplings are required due to the involvement of massive top quarks in conversions [201]. Since decays and scatterings contribute to conversions, the condition $\Gamma_{\rm conv} \sim H$ constrains the decay rate $\Gamma_{\rm dec} \lesssim H$ at freeze-out. For DM masses near the weak or TeV scale, this implies $c\tau_Y \gtrsim (10^{-3}-1)$ m, indicating a long-lived mediator with significant collider implications [34] (see Sect. 3.3).

Conversion-driven freeze-out requires solving coupled Boltzmann equations for the DM and mediator abundances. In addition, the weak strength of the DM couplings can induce deviations from kinetic equilibrium, necessitating to solve a momentum-dependent Boltzmann equation [34,35]. However, for the models considered, these deviations have a minor quantitative impact [34], and the system can often be described using the coupled rate equations (35) for i = X, Y, after explicitly accounting for the conversion processes entering the rates $\gamma_{X \to Y}$ and $\gamma_{Y \to X}$. For λ values where $\Gamma_{\text{conv}} \lesssim H$, annihilation processes proportional to λ are typically negligible, leaving mediator pair-annihilation ($Y\bar{Y} \to SM$) as the only relevant channel. In this case, the Boltzmann equations reduce to:

$$\begin{split} \frac{\mathrm{d}Y_X}{\mathrm{d}x} &= \frac{1}{Hxs} \left[\gamma_{Y \to X} \left(\frac{Y_Y}{Y_Y^{\mathrm{eq}}} - \frac{Y_X}{Y_X^{\mathrm{eq}}} \right) + \gamma_{\bar{Y} \to X} \left(\frac{Y_{\bar{Y}}}{Y_{\bar{Y}}^{\mathrm{eq}}} - \frac{Y_X}{Y_X^{\mathrm{eq}}} \right) \right], \\ \frac{\mathrm{d}Y_Y}{\mathrm{d}x} &= \frac{-1}{Hxs} \left[\gamma_{Y\bar{Y}} \left(\frac{Y_YY_{\bar{Y}}}{Y_Y^{\mathrm{eq}}Y_{\bar{Y}}^{\mathrm{eq}}} - 1 \right) + \gamma_{Y \to X} \left(\frac{Y_Y}{Y_Y^{\mathrm{eq}}} - \frac{Y_X}{Y_X^{\mathrm{eq}}} \right) \right], \end{split}$$

$$(41)$$



Eur. Phys. J. C (2025) 85:975 Page 57 of 96 975

where we assume the detailed balance condition $\gamma_{X \to Y} =$ $\gamma_{Y \to X}$. The mediator Y and its antiparticle \bar{Y} are treated separately for clarity, and we have assumed a self-conjugate DM species X. The Boltzmann equation for \bar{Y} is analogous, and under *CP*-conservation $Y_Y = Y_{\bar{Y}}$ and $\gamma_{\bar{Y} \to X} = \gamma_{Y \to X}$. This reduces the system to the two coupled equations given above (see [82] for a CP-violating scenario in the context of barvogenesis). Furthermore, the mediator annihilation rate $\gamma_{V\bar{V}}$ is dominated by gauge interactions, including QCD for coloured t-channel mediators and electroweak interactions for leptophilic mediators. Accordingly, these processes are subject to Sommerfeld enhancement and bound-state formation effects, as discussed in Sect. 4.2.5 [258,261,302]. At very low temperatures, mediator pair-annihilation eventually becomes subdominant due to the double Boltzmann suppression related to the initial-state mediators. Thus, conversiondriven freeze-out arises from the interplay between the freeze-out of the conversion processes $\gamma_{Y \to X}$ due to the small coupling strength λ , and deviations from equilibrium of the mediator itself due to a finite annihilation rate $\gamma_{Y\bar{Y}}$. Both are crucial in determining the DM relic abundance [34].

The cosmologically viable CDFO region in the model parameter space requires the DM to be under-abundant, when assuming, regardless of λ , chemical equilibrium between the X and Y states. This condition implies relatively small mass splitting between the dark matter and the mediator particles. However, it cannot be met above a certain mediator mass for which $\gamma_{Y\bar{Y}}$ is too small to provide sufficient dilution of the dark sector particles, even for mass-degenerate X and Y particles. This limits the allowed mass range for the mediator to a few hundred GeV (a few TeV) for leptophilic [260] (quark-philic [258]) models. CDFO scenarios therefore provide a well-bounded parameter space, that is likely fully testable with collider searches in the foreseeable future. This constitutes a difference from freeze-in or superWIMP scenarios that permit a broader mediator mass range.

Another notable feature of the CDFO mechanism is that, despite weak DM couplings, the DM particle remains thermalised, making the relic abundance independent of the initial conditions, akin to canonical freeze-out. In contrast, freeze-in and superWIMP scenarios require even weaker λ such that DM never reaches thermal equilibrium during the entire evolution of the universe. CDFO has been studied in different minimal *t*-channel mediator models, including quark-philic [34,201,258,259] and leptophilic [260] setups (where the Higgs-portal interactions of the mediator can become important), as well as in non-minimal models like flavoured DM [77,82]. Notably, CDFO may simultaneously achieve baryogenesis [82], motivating further exploration.

4.2.3 Freeze-in

Within the framework of freeze-in production of DM [36,37, 200], we assume that the DM particle X is so feebly coupled ($\lambda \lesssim 10^{-10}$) that it has never reached chemical or kinetic equilibrium (see Fig. 37). Under this assumption and starting with a negligible initial abundance of dark matter, the evolution of the DM population is governed primarily by conversion processes $Y \to X$, which include both decays and scatterings. The Boltzmann equation for the evolution of DM abundance (35) simplifies to

$$\frac{\mathrm{d}Y_X}{\mathrm{d}x} = \frac{1}{Hxs} \gamma_{Y \to X} \frac{Y_Y}{Y_v^{\mathrm{eq}}},\tag{42}$$

where we neglect $X \rightarrow Y$ contributions because the DM abundance is highly suppressed at early times. During freezein production, we assume that the mediator Y remains in chemical and kinetic equilibrium, as most of the DM production process occurs slightly before the freeze-out of the mediator Y ($T \gtrsim M_Y$). Consequently, we can approximate $Y_Y \approx Y_V^{\rm eq}$ all along the production process. This assumption holds for a freeze-in mechanism embedded in a renormalisable theory where production is dominated by infrared (IR) effects, in contrast to models featuring ultraviolet freezein [305]. For the simplified t-channel DM models analysed in this report, freeze-in remains IR-dominated as long as $M_Y < T_{\text{max}}$, where T_{max} is the maximum temperature of the universe. For scenarios with DM production occurring after the chemical decoupling of the mediator, we refer instead to Sect. 4.2.4. However, we must keep in mind that both contributions should generally be taken into account, as discussed, for example, in [202, 204].

Interactions with the thermal plasma can play a crucial role in scenarios of freeze-in dark matter production. This was first demonstrated in [289] for a Majorana DM model, and later extended to scalar DM models in [284,306]. The latter works employed a first-principles non-equilibrium quantum field theory approach, specifically the closed-time path formalism, to quantify the corrections to standard approximations commonly used in the literature for freeze-in dark matter. These approximations include vacuum decays only, decays with thermal masses, decays and scatterings with thermal masses, and the hard thermal loop approximation. It was shown that using decays with vacuum masses alone can underestimate the relic abundance by up to 90% in scenarios with small mass splittings where scattering processes, particularly those involving multiple soft scatterings, dominate. In contrast, for large mass splittings where decay processes dominate, the relic abundance is predicted correctly to within $\pm O(10\%)$, the sign depending on the gauge coupling. For large gauge couplings, the relic abundance is typically underestimated by about 10%, whereas for small gauge couplings it is overestimated by a similar margin. For further details,



975 Page 58 of 96 Eur. Phys. J. C (2025) 85:975

we refer the reader to [284,306]. Scattering processes such as $VY \to Xf$ may contribute slightly to the relic abundance, up to $\sim 25\%$, when V is a gluon and f is a coloured SM fermion. A more precise treatment of such thermal effects, particularly for small mass splittings, may nevertheless be necessary [289]. For simplicity, we focus here on mediator decay contributions during a radiation-dominated era to illustrate the main aspects of freezing-in DM production. In this case, (42) reduces to

$$\frac{\mathrm{d}Y_X}{\mathrm{d}x} = \frac{1}{Hx} \Gamma_{Y \to X} \frac{K_1(x)}{K_2(x)} Y_Y^{\mathrm{eq}},\tag{43}$$

where $\Gamma_{Y \to X}$ is the decay rate of $Y \to Xf$, and $x = M_Y/T$. For DM production via decay, which is also relevant for the superWIMP scenarios discussed in Sect. 4.2.4, it is convenient to define the dimensionless parameter

$$R_{\Gamma}^{\text{prod}} = \frac{M_0(T_{\text{prod}})\Gamma_{Y \to X}}{M_V^2},\tag{44}$$

where $M_0(T)=45M_{\rm pl}/(4\pi^3g_*(T))$ is the rescaled Planck mass entering the Hubble rate during a radiation-dominated era, $H=T^2/M_0(T)$, evaluated at the DM production temperature $T_{\rm prod}$. In addition, $g_*(T)$ denotes the number of relativistic degrees of freedom at a temperature T. In the case of freezing-in DM, $T_{\rm prod} \simeq T_{\rm FI} \simeq M_Y/3$. In this context, $R_{\Gamma}^{\rm prod} \equiv R_{\Gamma}^{\rm FI}$ approximately represents the ratio of the DM production rate to the Hubble rate at $T=M_Y$.

Integrating (43) over time yields

$$\Omega_X h^2|_{\text{FI, dec}} = M_X \times \frac{135}{8\pi^3} \frac{g_Y}{g_*(T_{\text{FI}})} R_{\Gamma}^{\text{FI}} \frac{s_0 h^2}{\rho_{\text{crit}}},$$
 (45)

where $\rho_{\rm crit}=3M_{\rm pl}^2H_0^2/(8\pi)$ is the critical energy density today, s_0 is the entropy density today, h is the rescaled Hubble parameter, $h=H_0/(100\,{\rm km\,s^{-1}Mpc^{-1}})\sim 0.7$, and g_Y is the number of degrees of freedom of the mediator. Increasing the DM coupling λ enhances $\Gamma_{Y\to X}$ (or equivalently $R_\Gamma^{\rm FI}$), increasing the DM production rate and resulting in a larger relic abundance. Finally, non-negligible scattering contributions may also be added to the decay terms in equations (43)–(45).

In summary, the freeze-in production of DM is a cumulative effect of rare decay or scattering processes involving the thermalised mediator particle Y, which produces DM at a very low (sub-Hubble) rate. Consequently, the final DM abundance decreases strongly with a decreasing coupling strength λ , as shown in Fig. 37. However, as λ is reduced further, a plateau in the relic density is reached, as indicated at the leftmost end of Fig. 37. We now turn to this extremely weakly coupled regime.



For extremely small coupling strengths between the mediator and the DM particle, the superWIMP production mechanism [38,39] can account for the observed relic DM density, as shown in Fig. 37. This mechanism typically unfolds in two distinct stages. First, at temperatures above or around the mediator mass, the mediator remains in thermal equilibrium with the SM thermal bath due to its gauge interactions, while the DM particle is decoupled because of its extremely weak interaction strength, assuming a negligible initial abundance. When the temperature drops below the mediator mass, mediator pair annihilation into SM particles becomes Boltzmannsuppressed and the mediator then eventually freezes out, similarly to the usual WIMP mechanism. At this stage, the DM abundance remains negligibly small because of its tiny interaction strength. The evolution of the mediator abundance during this stage is governed by conventional freeze-out via its gauge interactions, while conversion to DM is neglected. The corresponding Boltzmann equation is:

$$\frac{\mathrm{d}Y_Y}{\mathrm{d}x} = \frac{-1}{Hxs} \left[\gamma_{Y\bar{Y}} \left(\frac{Y_Y Y_{\bar{Y}}}{Y_Y^{\mathrm{eq}} Y_{\bar{Y}}^{\mathrm{eq}}} - 1 \right) \right],\tag{46}$$

with a similar equation for the antiparticle \bar{Y} . When the age of the universe becomes comparable to the lifetime of the mediator, the frozen-out distribution of mediator particles decays into DM. For each mediator particle one DM particle is therefore produced. For self-conjugate DM (e.g., a Majorana fermion or a real scalar), this implies

$$Y_X^{\text{SW}} = (Y_Y + Y_{\bar{Y}})|_{\text{FO}}.$$
 (47)

These decays also produce SM particles, and impart a boost to the DM particles whose magnitude depends on the mass ratio between the mediator and the DM particle. This boost can result in a non-negligible DM velocity distribution, potentially leading to constraints that could be derived from structure formation. In particular, Lyman- α forest observations could provide very strong bounds [202,204], as discussed in Sect. 4.3.3, and the long lifetime of the mediator can additionally result in collider signatures of long-lived particle searches and may affect Big Bang Nucleosynthesis (BBN) [202].

The scenario described above can be modified under certain conditions, particularly when non-perturbative effects are taken into account. For instance, the formation of bound states between pairs of mediator particles can significantly enhance their effective annihilation rate, $\gamma_{Y\bar{Y}}$ [258,261,302]. Specifically, for a mediator that is both coloured and electrically charged, bound states formed through gluon exchanges together with electromagnetic transition processes among the different bound state levels can prevent the mediator from freezing out completely. In such cases, the mediator abun-



Eur. Phys. J. C (2025) 85:975 Page 59 of 96 975




Fig. 39 Diagramatic representation of bound state (\mathcal{B}) formation of two coloured particles via the emission of a gluon, and of the Sommerfeld Effect relevant for the initial state. Figure adapted from [302]

dance continues to be depleted even at temperatures much lower than its mass, and this depletion ceases only when the age of the universe becomes comparable to the mediator lifetime. When these effects are significant, the two stages described earlier become interconnected rather than distinct. Further details on this regime can be found in [261]. On the other hand, while bound states are also quantitatively relevant for leptophilic models, they do not qualitatively alter the evolution of the superWIMP mechanism as outlined above.

4.2.5 Non-perturbative effects

The freeze-out of dark sector annihilations typically occurs when the annihilating particles are non-relativistic. This regime allows for significant non-perturbative effects, namely the Sommerfeld Effect (SE) and Bound State Formation (BSF), that affect the annihilation cross section through multiple exchanges of light states between the initial or final-state particles. Non-perturbative corrections in the final state are neglected here, since the annihilation products are much lighter than the dark sector particles so that they are relativistic.

Simplified t-channel DM models always involve a dark sector particle Y carrying SM charges. Consequently, annihilation involving two SM-charged dark sector particles is subject to corrections from multiple exchanges of (massless) gauge bosons. Each exchange of a massless gauge boson contributes parametrically with a factor of α/v where α is the relevant coupling constant and v is the relative velocity between the two initial-state particles, which could be sizeable if $\alpha \sim v$. During the freeze-out of DM annihilations, $v \sim 0.1$. This thus necessitates the resummation of the ladder diagrams shown in Fig. 39 if $\alpha \sim 0.1$, which is particularly relevant for mediator particles Y charged under $SU(3)_C$ (as in the models considered). In the non-relativistic regime, this resummation reduces to solving the Schrödinger equation for the wave functions of the initial-state particles, incorporating a colour potential as discussed in [293,307-310]. Since gluons are massless, the potential at energy scales far above the QCD confinement scale is Coulomb-like, and can be either attractive or repulsive depending on the representation of the initial state particles. In the context of simplified t-channel DM models, the dark sector particles are either singlet or lie in the (anti-)fundamental representation of $SU(3)_C$. Dark sector annihilation therefore leads to the following initial state colour configurations,

$$3 \times \bar{3} = 1 + 8,$$

 $3 \times 3 = \bar{3} + 6.$ (48)

Among these, the 1 and $\bar{3}$ configurations result in an attractive potential, allowing for particle-antiparticle and particle-particle bound states, respectively, while the 6 and 8 configurations are repulsive.

The Sommerfield effect affects both attractive and repulsive initial states by capturing the distortion of the incoming wave functions due to the effective colour potential, which are otherwise modelled as plane waves. For repulsive configurations, the cross section is reduced, while for attractive ones, it is enhanced. This effect can be incorporated by multiplying the perturbatively calculated cross section for each partial wave by the corresponding SE factor S_l [311,312],

$$\sigma_{ij}v_{ij} \to \sum_{l} \left(\sigma_{ij}v_{ij}\right)_{l} S_{l}\left(\frac{\alpha_{\text{eff}}}{v_{ij}}\right),$$
 (49)

where l denotes the lth partial wave.

Bound state formation contributes to the effective annihilation cross section by introducing an additional channel. A bound state forms via the emission of at least one gluon, and it can subsequently either decay into dark sector or SM particles, or be ionised by a gluon from the thermal bath. Only bound states that decay contribute to the effective annihilation cross section by depleting the dark sector. The related contribution in the case of a single bound state is given by [313]

$$\langle \sigma_{\rm eff} v \rangle \to \langle \sigma_{\rm eff} v \rangle + \frac{\Gamma_{\rm dec}}{\Gamma_{\rm dec} + \Gamma_{\rm ion}} \langle \sigma_{\rm BSF} v \rangle,$$
 (50)

where $\langle \sigma_{\rm BSF} v \rangle$ is the thermally averaged cross section for bound state formation under the emission of a gluon [301]. The prefactor depends on the bound state decay and ionisation rates $\Gamma_{\rm dec}$ and $\Gamma_{\rm ion}$, and reflects that only bound state decays into SM particles deplete the dark sector. When excited bound states are included, transitions between different states i and j with rates $\Gamma_{\rm trans}^{i \to j}$ must be considered, which yields [258,261,314]

$$\langle \sigma_{\text{eff}} v \rangle \to \langle \sigma_{\text{eff}} v \rangle + \sum_{i} R_{i} \langle \sigma_{\text{BSF},i} v \rangle,$$
 (51)

with

$$R_{i} = 1 - \sum_{j} \left(\delta_{ij} - \frac{\Gamma_{\text{trans}}^{i \to j}}{\Gamma_{\text{ion}}^{i} + \Gamma_{\text{dec}}^{i} + \sum_{j \neq i} \Gamma_{\text{trans}}^{i \to j}} \right)^{-1}$$



975 Page 60 of 96 Eur. Phys. J. C (2025) 85:975

$$\frac{\Gamma_{\text{ion}}^{j}}{\Gamma_{\text{ion}}^{i} + \Gamma_{\text{dec}}^{i} + \sum_{j \neq i} \Gamma_{\text{trans}}^{i \to j}}.$$
 (52)

When summing the bound-state formation contributions to the cross section over all possible bound states in the final state, unitarity violation however arises [315,316], although a possible solution has been recently proposed in [317].

In summary, the annihilation cross section involving two possibly different coloured particles Y_1 and Y_2 must be modified to include two classes of non-perturbative effects, Sommerfeld effect and bound state formation effects,


$$\langle \sigma_{Y_1 Y_2} v \rangle \rightarrow \sum_{l} \langle S_l \left(\frac{\alpha_{\text{eff}}}{v} \right) \left(\sigma_{Y_1 Y_2} v \right)_{l} \rangle + \sum_{i} R_i \langle \sigma_{\text{BSF}, i} v \rangle.$$
 (53)

The Sommerfeld effect and bound state formation have been extensively studied in *t*-channel scenarios [258,288, 294,295,301–304,318]. Their impact on the interpretation of the constraints existing in the parameter space is in particular explored in [302] for freeze-out scenarios and in [258] for CDFO models.

4.2.6 Altered cosmological histories

The precise measurement of the cosmic abundance of light element relics provides one of the strongest constraints on the properties of the universe at the time of their creation, specifically when BBN occurred at temperatures of approximately $T \sim O(1)$ MeV [319]. Experimental data indicates that the universe must have been in a period of radiation domination by the time of BBN, with the energy density of the relativistic particles in the thermal bath driving the expansion. At earlier times, however, the expansion history of the universe remains uncertain, as no data conclusively establishes the expansion history between inflation and BBN. This period could span up to 20 orders of magnitude in energy (given that the energy scale of inflation must be below 10^{16} GeV), during which various phenomena could have altered the expansion rate, impacting particle interactions and production [273,274]. Among the possibilities are low-reheating temperature scenarios after inflation, early matter domination epochs [320,321], and faster-than-radiation eras such as kination [322, 323]. These phenomena could result from slow decays of the inflaton during reheating [274], decays of heavy particles like moduli fields (scalar fields with generic equations of state), primordial black holes, quintessence models of dark energy, and more. For a comprehensive discussion, we refer the reader to the review [275].

If the production of dark sector particles occurs during such altered cosmological histories, the evolution of their densities can change significantly, leading to predictions for model observables that differ by orders of magnitude [255]. For instance, in scenarios with low-reheating temperatures,



$$T \propto a^{-k},$$
 (54)

where k is a model-dependent real number that under adiabatic conditions would be k=1. If the production of a relic such as dark matter ceases during the reheating era, its abundance undergoes dilution until entropy is conserved again, for instance when the universe transitions back to a radiation-dominated phase. This dilution can span several orders of magnitude, drastically impacting predictions for dark sector particle masses and couplings.

For simplified t-channel DM models, low-reheating temperature scenarios have been studied in [252,254], focusing on freeze-in production for both Majorana and scalar DM models with scalar and vector-like fermion mediators, respectively. Ref. [252] highlights that when the reheating temperature T_{RH} is lower than the mediator mass M_Y , such that DM freezes in during reheating, entropy dilution significantly reduces the comoving number density. Reproducing the observed DM relic density for a fixed mass configuration then requires much stronger Yukawa couplings between DM, the mediator, and the SM fermions. This, in turn, shifts the collider signatures of the mediator from the long-lived particle regime to the prompt regime, making the interpretation of potential signals and constraints from colliders highly dependent on the reheating history. This result was extended in [254] to reheating scenarios with potentials of the form $V(\phi) \propto \phi^k$ ($k \geq 2$) and different inflaton decay channels into bosons and fermions, leading to distinct expansion histories and entropy dilution factors. Such scenarios are motivated by α -attractor models of inflation [324–326], which predict inflationary observables consistent with the constraints from the Planck 2018 analysis [327].

As shown in Sect. 3.3, these findings modify the predictions for LLP signatures, demonstrating the intricate dependence of the results on the reheating potential and the main inflaton decay channel. This arises from the power-law dilution of the relic abundance,

$$Y_X \sim \Gamma_Y \times \begin{cases} \left(\frac{T_{\text{rh}}}{M_Y}\right)^{4k-1} & \text{Bosonic Reheating,} \\ \left(\frac{T_{\text{rh}}}{M_Y}\right)^{\frac{9-k}{k-1}} & \text{Fermionic Reheating,} \end{cases}$$
 (55)

which is true if $M_Y \gtrsim T_{\rm RH}$. Although dilution of the relic abundance necessitates larger mediator decay widths in freeze-in scenarios, it reduces the DM-SM interaction



Eur. Phys. J. C (2025) 85:975 Page 61 of 96 975

strength for thermal freeze-out production. Since freeze-out during reheating has not been explored in the context of simplified t-channel DM models, we refer instead the reader to general discussions in e.g. [277,328,329].

4.3 Dark matter constraints

In this section, we describe the most relevant astrophysical and cosmological constraints that could be applied to the models studied.

In Sects. 4.3.1 and 4.3.2, we focus on astrophysical searches probing DM interactions at the present time, where DM is cold and clustered within halos in galaxies and galaxy clusters. The typical velocities v involved range from approximately $10^{-5}c$ to $10^{-2}c$, with $v \sim 10^{-3}c$ in the Milky Way, and current detection methods primarily include indirect and direct detection approaches. Indirect detection relies on the annihilation of DM into SM particles in regions with high DM density, such as the Galactic Centre and dwarf spheroidal galaxies (dSphs). The SM particles produced through DM annihilations subsequently decay, shower, and hadronise to produce a flux of stable cosmic rays $(\gamma, e^+, \bar{p}, \text{ and } \nu)$, which can be detected by space-based and ground-based telescopes. Direct detection, on the other hand, relies on observing elastic scattering of DM off protons or neutrons within underground detectors, where the scattering produces a measurable recoil of the associated nucleus. However, these detection methods are most effective in probing the canonical freeze-out regime, where the DM annihilation or DM-nucleon scattering rates are sufficiently large. In contrast, scenarios that involve the freeze-in, superWIMP, or conversion-driven freeze-out mechanisms predict extremely small rates that are typically not testable with these methods. Instead, such scenarios often give rise to the interesting long-lived particle signals at colliders that have been discussed in Sect. 3.3.

Cosmological constraints are presented in Sect. 4.3.3. While constraints from the cosmic microwave background (CMB) apply broadly across all DM scenarios, constraints from Big Bang Nucleosynthesis and structure formation are particularly relevant for CDFO and freeze-in and super-WIMP scenarios.

Several computational tools are available to automatically calculate DM observables and constraints such as those originating from the calculation of the relic density, as well as indirect and direct detection predictions. These tools generally perform calculations at LO, with some also offering a handle on NLO or loop-induced processes, and most of them facilitate automated testing of model parameter space against a wide range of experimental observables. The set of most comprehensive and widely used tools includes micromegas [53], MadDM [56], DarkSUSY [330] and GAMBIT [331].

4.3.1 Indirect detection

Predictions for present-day annihilation of DM in dense astrophysical environments, in the context of minimal simplified models coupling DM to fermions (quarks and/or leptons), depend strongly on the spin of the DM particles. The dominant tree-level annihilation process, $XX \rightarrow f\bar{f}$, is illustrated by the Feynman diagram shown in the top left corner of Fig. 38.

For Majorana and real scalar DM, this process is P-wave and D-wave suppressed, respectively, with the S-wave contribution being proportional to the fermion mass. This S-wave contribution is therefore negligible in the chiral limit or for top quarks in scenarios featuring DM and mediator masses above a few hundred GeV. For complex scalar DM, this suppression is still relevant, and it is primarily P-wave. There are however two processes that could enhance the detectability of such DM particles. Virtual internal bremsstrahlung (VIB), where the final-state quark pair or lepton pair is produced alongside a photon emitted by the internal t-channel propagator $(XX \to f \bar{f} \gamma)$, can provide significant corrections to the tree-level annihilation cross section under certain mediator-DM spectrum configurations, overcoming Pwave suppression by orders of magnitude. Moreover, VIB yields a sharp spectral feature at the high-energy end of the gamma-ray spectrum, that could enhance the sensitivity of related experiments [21,332-341]. Additionally, loopinduced annihilation processes, such as $XX \to \gamma \gamma$, γZ , or γH , produce distinctive gamma-ray lines that have the potential to directly reveal the DM mass [342–347]. Representative Feynman diagrams for these two sets of processes are given in Fig. 40, and we emphasise that analytical expressions for the most studied models are available in the literature, such as in [21,201,337,338]. Moreover, automated tools like MadDM could be used to compute predictions for such higher-order processes for any DM model, provided that the Lagrangian is translated in the UFO format at NLO [47].

In freeze-out scenarios where DM self-annihilation dominates and proceeds via S-wave processes (as for Dirac, real vector and most complex DM models except when it is a scalar), indirect detection signals are typically expected in the form of a continuum of photons, positrons, neutrinos and antiprotons. However, in regions where co-annihilation or bound-state formation plays an important role (typically when the mediator mass is less than about 1.2 times the DM mass), the connection between the relic density and indirect detection can weaken or break down entirely. In particular, if the relic density is set by processes involving the pair-annihilation of the co-annihilating partner (such as *YY* annihilations to a pair of SM particles), no indirect detection signal is expected today. Nonetheless, in the mass and coupling ranges considered, annihilation cross sections near



975 Page 62 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 40 VIB and loop-induced processes contributing to DM indirect detection through the production of gamma-ray lines and other features in the gamma-ray spectrum

the canonical value of 10^{-26} cm³/s still arise in considerable parts of the parameter space.


The search for gamma-ray fluxes from DM annihilation in dwarf spheroidal galaxies, as performed by the Fermi-LAT satellite [348,349], is among the most sensitive methods to probe t-channel DM models, and it can exclude a large portion of the parameter space of minimal models [25]. Measurements of cosmic-ray antiproton fluxes by the AMS-02 experiment at the International Space Station [350–352] provide complementary constraints on DM annihilation in our galaxy, particularly for higher DM masses. While gammaray lines and VIB are higher-order contributions, they yield sharp spectral features that can be distinguished from smooth astrophysical foreground emission [333]. These features lead to excellent experimental sensitivity from Fermi-LAT [353] and other gamma-ray observatories, and allow us to probe cross sections well below 10^{-26} cm³/s for a wide range of DM masses. We should however keep in mind that theoretical predictions for these higher-order processes are however also often smaller than this benchmark value.

For DM models involving leptonic interactions, annihilations into neutrinos, either monochromatic or as a continuum, are instead relevant. Although the observation of the associated signal is partially challenged by continuum photon emission [354] and the existing antiproton flux [355], neutrinos produced from DM annihilation can in principle be detected by current and future neutrino telescopes like IceCube [356] and KM3NeT [357]. This has been particularly demonstrated for specific models such as secluded t-channel models and $U(1)_{L_{11}-L_{12}}$ gauge scenarios [358–360].

Finally, gamma-ray line searches by HESS [361,362] and MAGIC [363], along with neutrino searches, provide sensitivity to very heavy DM, offering a complementary probe to collider experiments limited by the current centre-of-mass energies and luminosities.

4.3.2 Direct detection

The standard direct detection signal considered in simplified t-channel models is the scattering of dark matter off nucleons. This signal varies significantly depending on the nature of the DM particle, the charge of the mediator, and its coupling to the SM particles. In this section, we illustrate the



For a Dirac DM particle, interactions with nucleons can be described by the representative Feynman diagrams shown in Fig. 41, which depicts tree-level and one-loop contributions. These include the tree-level scattering process featuring the exchange of the coloured scalar mediator Y (first two diagrams in the bottom row), as well as one-loop penguin diagrams mediated by the photon, the Z boson, or the Higgs boson (two rightmost diagrams in the bottom row). While the tree-level contributions only arise for DM coupling to the up or down quark, the loop-induced subprocesses arise universally, regardless of the choice of SM fermion connecting the dark and SM sectors. Additionally, box and triangle diagrams involving two external gluons appear at one-loop, and are ubiquitous of a model where the DM couples to a coloured mediator. We now discuss below the effective Lagrangian interactions between the DM state and nucleons arising in each case.

If the DM particle is a Dirac fermion and couples at tree level to a first-generation quark, the dominant contribution to the scattering cross section comes from the t-channel exchange of the scalar mediator Y, as depicted in Fig. 41. For the DM antiparticle \bar{X} , scattering off quarks occurs additionally via s-channel exchanges of the mediator. After a Fierz rearrangement of the corresponding matrix elements, these diagrams yield a vector interaction of the X states with quarks, which translates into an effective vector interaction with nucleons N,

$$\mathcal{L}_{\text{eff,tree}} = f_{V,\text{tree}}^{(N)} \bar{X} \gamma^{\mu} X \bar{N} \gamma_{\mu} N. \tag{56}$$

This leads to spin-independent interactions in the non-relativistic limit, with the effective couplings for protons and neutrons appearing in this expression, $f_{V,\text{tree}}^{(p)}$ and $f_{V,\text{tree}}^{(n)}$, being given by

$$f_{V,\text{tree}}^{(p)} = \begin{cases} 2 \, \lambda_V & \text{for coupling to } u_R \\ \lambda_V & \text{for coupling to } d_R \\ 3 \, \lambda_V & \text{for coupling to } (u_L, d_L) \end{cases} \,,$$



Eur. Phys. J. C (2025) 85:975 Page 63 of 96 975




Fig. 41 Illustrative Feynman diagrams contributing to DM-nucleon scattering, where we select as an example the case of fermionic DM. We include all diagrams that account for the scattering off gluons at one-loop and that are always present regardless of the model details

(top row), tree-level scattering (relevant if the DM couples to an up or down quark) and the one-loop exchange of a photon, Z boson or Higgs boson

$$f_{V,\text{tree}}^{(n)} = \begin{cases} \lambda_V & \text{for coupling to } u_R \\ 2\lambda_V & \text{for coupling to } d_R \\ 3\lambda_V & \text{for coupling to } (u_L, d_L) \end{cases}$$
(57)

with

$$\lambda_V = \frac{\lambda^2}{8\left(M_Y^2 - M_X^2\right)}. (58)$$

Due to vector current conservation, similar spin-independent tree-level contributions relevant for DM coupling to second-generation or third-generation quarks vanish, and they are (obviously) irrelevant for DM coupling solely to leptons. These diagrams also generate an axial-vector interaction of the form $f_{A,\mathrm{tree}}^{(N)} \bar{X} \gamma^{\mu} \gamma_5 X \bar{q} \gamma_{\mu} \gamma_5 q$ that leads to spin-dependent interactions. The coupling constants of the axial-vector and the vector interaction terms are generally comparable in size in t-channel models. Therefore, given that the experimental sensitivity to spin-independent interactions is several orders of magnitude better than in the spin-dependent case, this axial-vector contribution can usually be neglected.

For a Majorana DM particle coupled to first-generation or second-generation quarks, the vector current vanishes at tree level. Spin-independent interactions then arise predominantly from higher-order effects, like interactions with gluons via loop diagrams such as those shown in Fig. 41, or higher-order corrections to the conventional mediator exchanges present at tree level. These generate a higher-dimensional effective Lagrangian involving quark and gluon operators,

$$\mathcal{L}_{q}^{\text{eff}} = f_{q} \, \bar{X} X \, O_{q}^{(0)} + \frac{g_{q}^{(1)}}{M_{X}} \, \bar{X} \big(i \partial^{\mu} \gamma^{\nu} + i \partial^{\nu} \gamma^{\mu} \big) X \, O_{q,\mu\nu}^{(2)} \\
+ \frac{g_{q}^{(2)}}{M_{X}^{2}} \, \bar{X} (i \partial^{\mu}) (i \partial^{\nu}) X \, O_{q,\mu\nu}^{(2)}, \\
\mathcal{L}_{g}^{\text{eff}} = f_{G} \, \bar{X} X \, O_{g}^{(0)} + \frac{g_{G}^{(1)}}{M_{X}} \, \bar{X} \big(i \partial^{\mu} \gamma^{\nu} + i \partial^{\nu} \gamma^{\mu} \big) X \, O_{g,\mu\nu}^{(2)} \\
+ \frac{g_{G}^{(2)}}{M_{X}^{2}} \, \bar{X} (i \partial^{\mu}) (i \partial^{\nu}) X \, O_{g,\mu\nu}^{(2)}.$$
(59)

Here, the operators *O* represent the scalar and higher-twist quark and gluon operators defined by

$$\begin{split} O_q^{(0)} &\equiv m_q \bar{q} q, \ O_q^{(2)\mu\nu} \equiv \frac{1}{2} \bar{q} \left(\gamma^{\{\mu} i D_-^{\nu\}} - \frac{1}{4} g^{\mu\nu} i \not D_- \right) q, \\ O_g^{(0)} &\equiv G_{\mu\nu}^A G^{A\mu\nu}, \\ O_g^{(2)\mu\nu} &\equiv -G^{A\mu\lambda} G^{A\nu}_{\quad \lambda} + \frac{1}{4} g^{\mu\nu} (G_{\alpha\beta}^A)^2, \end{split} \tag{60}$$

where we have introduced the standard shorthand notation

$$A^{\{\mu}B^{\nu\}} = \frac{1}{2}(A^{\mu}B^{\nu} + A^{\nu}B^{\mu}) \text{ and } D^{\mu}_{\pm} = D^{\mu} \pm \overleftarrow{D}^{\mu},$$
(61)

with D^{μ} being the usual covariant derivative. These operators yield the nucleon matrix elements detailed in [22,365,366], with the analytical expressions for the corresponding Wilson coefficients f_q , f_G , $g_q^{(1)}$, $g_q^{(2)}$, $g_G^{(1)}$, and $g_G^{(2)}$ given in [22,366,367]. Since these Wilson coefficients are generated from higher-order contributions, the resulting spin-independent and spin-dependent cross sections are comparable. Consequently, constraints from spin-independent cross sections typically dominate when the DM and mediator masses are close, while constraints from spin-dependent rates dominate for DM masses comparably smaller than the mediator mass [366]. Finally, although such contributions also exist for models featuring a Dirac DM fermion, the resulting Wilson coefficients are subdominant and thus often ignored.

If the DM does not couple to first-generation and second-generation quarks, the penguin diagrams in Fig. 41 become more significant. Regardless of the precise SM fermion species involved, the DM interaction with quarks necessarily induces, at the quantum level, DM-nucleon scattering due to penguin diagrams mediated by a photon, a Z boson, or a Higgs boson. The photon-mediated diagram generates electromagnetic moments for the DM particle X, with the most relevant ones in the case of Dirac DM being the magnetic dipole moment μ_X and the charge radius b_X . These are



described by the effective DM-photon Lagrangian¹³

$$\mathcal{L}_{\text{eff},\gamma} = \frac{\mu_X}{2} \bar{X} \sigma^{\mu\nu} X F_{\mu\nu} + b_X \bar{X} \gamma^{\mu} X \partial^{\nu} F_{\mu\nu}. \tag{62}$$

The electromagnetic moments μ_X and b_X are determined by matching the coefficients of this effective Lagrangian to the results of explicit loop diagram calculations. Moreover, in the case of Majorana DM, all vector currents vanish, rendering thus these interactions irrelevant.

Similarly, DM interactions with the Z boson arise at one-loop, and induce effective vector interactions of the form

$$\mathcal{L}_{\text{eff},Z} = f_{V,Z}^{(N)} \bar{X} \gamma^{\mu} X \bar{N} \gamma_{\mu} N, \tag{63}$$

where

$$f_{V,Z}^{(p)} = \left(4s_W^2 - 1\right) \frac{G_F a_Z}{\sqrt{2}}, \quad f_{V,Z}^{(n)} = \frac{G_F a_Z}{\sqrt{2}}.$$
 (64)


Here, G_F represents the Fermi constant, s_W the sine of the electroweak mixing angle, and a_Z is an effective form factor whose analytical expression can be found in [340]. Furthermore, for DM coupling to the doublet of left-handed SM fermions, we have $a_Z^{(u_L,d_L)} = -a_Z^{(u_R)} - a_Z^{(d_R)}$ and $a_Z^{(v_L,e_L)} = -a_Z^{(e_R)}$. The DM effective coupling to the Z boson scales as $(M_f/M_X)^2$, making this contribution subdominant compared to photon exchange, except in scenarios where DM couples to the third-generation fields t_R or (t_L, b_L) .

The last penguin diagram in Fig. 41, i.e. the Higgs-mediated one, induces a coupling of the DM state X to the SM Higgs boson h, which in turn generates the effective DM-nucleon interaction

$$\mathcal{L}_{\text{eff,Higgs}} = f_{\text{S,Higgs}}^{(N)} \bar{X} X \bar{N} N, \tag{65}$$

where expressions for $f_{\rm S, Higgs}^{(N)}$ are provided in [340,369]. Since the coupling of fermions to the Higgs boson is proportional to their mass, this contribution is subdominant for DM coupling to light SM fermions. However, for DM coupling to third-generation quarks, the *Z*-exchange contribution is always dominant, making the Higgs exchange negligible as well. Additional contributions to the DM-Higgs interaction may arise from quartic couplings involving the scalar mediator and the Higgs boson, but even with quartic couplings of O(1), the Higgs-mediated contribution remains subdominant in DM-nucleon scattering.

Finally, in general and for any coupling to a Standard Model fermion, the running of the λ coupling from the electroweak scale down to the GeV scale significantly enhances direct detection cross section predicted values [366], often



In the parameter space of the t-channel mediator models discussed in this work, searches for spin-independent nucleon-DM scattering in data from the experiments XENON1T [371], XENONnT [372], and LZ [373] have achieved unprecedented sensitivity, imposing constraints even on loop-suppressed interactions. For models with couplings to first-generation quarks, next-generation experiments such as DARWIN [374,375] are expected to explore much of the currently viable parameter space, as their reach on the spin-independent scattering cross section exceeds the neutrino background [376]. Regarding spin-dependent proton scattering, the PICO-60 experiment [377] sets strong exclusion limits for DM masses below 500 GeV, even for Majorana DM [368]. In contrast, xenon-based detectors provide the strongest limits for spin-dependent neutron interactions. While this review was being finalised, impressive new results were released by PandaX-4T [378] and LZ2025 [379]. In particular, the LZ2025 exclusion limit for spinindependent DM-nucleus scattering improves upon that of XENONnT by more than an order of magnitude. These updated bounds are not included in the present analysis.

The sensitivity of direct detection experiments to all these operators is constrained by the minimum threshold energy required to produce detectable excitations in the detector material. For nucleon-DM scattering, the recoil energy is well below the detection threshold for sub-GeV DM masses [380,381]. In such cases, electron-DM scattering offers a promising alternative, as the larger available energy allows triggering inelastic atomic processes that produce visible signals [382]. An expanding experimental programme aims to explore this, including Super-CDMS [383], DAMIC [384], SENSEI [385], PandaX-II [386], DarkSide-50 [387], and XENON1T-S2 [388].

4.3.3 Early universe constraints

The early universe provides a wealth of constraints on the DM properties and interactions, through in particular observations of the CMB, BBN and structure formation.

The CMB anisotropies are a powerful tool for probing the history and content of the early universe, and their analysis constrains the DM abundance to $\Omega_X h^2 = 0.120 \pm 0.001$ [178]. Moreover, DM candidates that annihilate or decay into SM particles can alter the reionisation history, thus affecting the CMB anisotropy spectrum. This results in an upper bound on the annihilation efficiency, $p_{\rm ann} \lesssim 3 \times 10^{-28} \, {\rm cm}^3/{\rm s/GeV}$ [178], where $p_{\rm ann} = f_{\rm eff} \langle \sigma v \rangle / M_X$ with $f_{\rm eff}$ representing the fraction of energy from annihilation transferred to ionisation at redshifts relevant to CMB data. This bound, however, may not directly apply to t-channel scenarios where the DM annihilation cross section is P-wave or D-wave sup-



 $^{^{13}}$ The same diagram also induces an anapole moment, corresponding to the effective operator $\bar{X}\gamma^{\mu}\gamma_5X\partial^{\nu}F_{\mu\nu}$. However, this operator is suppressed in the non-relativistic limit by the square of the DM velocity, and is thus always subdominant in the *t*-channel models considered [368].

Eur. Phys. J. C (2025) 85:975 Page 65 of 96 975

pressed (see Sect. 4.3.1). For the scenarios considered here, we should instead rely on the fact that the DM particle itself cannot decay in the early universe, but the mediator can. CMB anisotropies can then constrain the lifetime of the decaying particles, with bounds such as $\tau > 10^{13}$ s [389]. However, stronger additional constraints also stem from Lyman- α forest data (see below). On the other hand, particles featuring shorter lifetimes can be constrained either by CMB spectral distortions or by BBN data (as discussed below too), and the CMB can finally also probe the effective number of relativistic degrees of freedom N_{eff} near the surface of last scattering with similar precision to BBN constraints. Unfortunately, for DM produced with relatively large momenta through the freeze-in or superWIMP mechanisms, the Planck 2018 N_{eff} bounds are always less stringent than those originating from the Lyman- α forest data analysis and applicable to DM freestreaming [204].

CMB spectral distortions also test the nature of the dark sector. These distortions, which arise from non-standard energy injection (e.g. from particle decays or annihilations) that disrupts the thermodynamic equilibrium between photons and free electrons after BBN ($z \lesssim 10^8$), can be probed and thus used as constraints on models [390]. The FIRAS instrument on COBE measured the CMB energy spectrum, and it found it consistent with a perfect black-body spectrum at $T_0 = 2.725 \pm 0.002$ K [391,392]. For mediator decays

Within the context of the t-channel models considered here, feebly interacting DM produced from decays or scatterings off an heavier mediator in the earlier universe can have very large momentum at the time of production compared to their mass, thereby affecting small scale structures and possibly leaving an imprint similar to warm DM (WDM) through free-streaming. The Lyman- α forest data, a typical tracer of small-scale structure clustering, provide stringent upper bound on the mass of thermal WDM, $M_{\rm wdm} \gtrsim 5.3$ keV [396]. Translating these constraints to non-thermally produced DM, via for example freeze-in or the superWIMP mechanism, requires costly hydrodynamical simulations or approximations using velocity dispersion or free-streaming scale comparisons [204–206,397–401]. Another more advanced approach requires to determine the non-cold DM velocity distribution at the time of production, feed it to a Boltzmann solver such as [402], obtain the corresponding linear power spectrum or the associated transfer function, and compare those to the WDM results. Finally, we could also employ the area criterion [204,206,397,398].

Remarkably, for freezing-in and superWIMP DM, the velocity distributions are similar in shape to the thermal WDM case but shifted, allowing a straightforward mapping of WDM bounds to constraints on M_X [204]. The resulting Lyman- α forest constraint can generally be expressed as [204]

$$M_X \gtrsim \begin{cases} 15 \text{ keV} \times \left(\frac{106.75}{g_*(T_{\text{FI}})}\right)^{1/3} & \text{for freeze-in through decays,} \\ 3.8 \text{ GeV} \times \left(R_{\Gamma}^{\text{SW}}/10^{-12}\right)^{-1/2} \times \left(\frac{106.75}{g_*(T_{\text{SW}})}\right)^{1/3} & \text{for superWIMP,} \end{cases}$$

$$(66)$$

with lifetimes $10^4~s < \tau < 10^{13}~s$, CMB spectral distortions provide stronger constraints than CMB anisotropies, though they are still weaker than those originating from BBN. Future experiments such as PIXIE or PRISM have nevertheless the potential to significantly improve the bounds [390].

Big Bang Nucleosynthesis, which occurred at $T_{\rm BBN} \sim$ 0.1 MeV, is another critical probe of dark matter. New particles can affect the primordial abundances of light nuclei by altering the Hubble rate or the entropy density of the universe. This leads to constraints on $N_{\rm eff}$, which are comparable to those originating from the CMB [393], but less stringent than the Lyman- α forest bounds discussed below. In addition, if the mediator has a lifetime longer than approximately 0.1 s, its decay can induce non-thermal nuclear reactions during or after BBN, modifying standard predictions [394,395]. Such constraints are especially relevant for superWIMP production scenarios involving mediators with lifetimes 10⁴ s $<\tau<10^{13}$ s that decay into hadronic final states. For example, in top-philic scenarios, these constraints help close the existing gap between constraints stemming from Lyman- α forest probes and the collider limits [201,204].

where $M_Y\gg M_X$ and $M_Y\gg M_f$, R_Γ is the dimensionless parameter introduced in (44) and proportional to the mediator decay rate and inversely proportional to its mass squared, and g_* denotes the number of relativistic degrees of freedom at production. Notably, Lyman- α forest data can probe DM masses well above the naive keV scale in the case of super-WIMP production, depending on the mediator's lifetime and mass.

4.4 Cosmological constraints for benchmark models

In this section, we summarise the cosmological and astrophysical constraints that could be imposed on several of the benchmark models considered. We start by examining minimal *t*-channel DM models in scenarios where the observed relic density is reproduced through conventional freeze-out, conversion-driven freeze-out and through the freeze-in/superWIMP mechanism in Sects. 4.4.1.1, 4.4.1.2 and 4.4.1.3, respectively. The discussion is thus structured by distinguishing between the different DM production mechanisms responsible for explaining the relic density across



975 Page 66 of 96 Eur. Phys. J. C (2025) 85:975

distinct domains in the model parameter space. Leptophilic models are next investigated in Sect. 4.4.1.5, before we finally turn in Sect. 4.4.2 on non-minimal models. More precisely, Sects. 4.4.2.1, 4.4.2.2, 4.4.2.3 and 4.4.2.4 are dedicated to models featuring multiple DM states X_i (e.g. flavoured DM), multiple mediators (e.g. frustrated DM), and mixed t-channel and s-channel mediator scenarios.

It is important to note that the selection of benchmark models presented here is not exhaustive. The choices are primarily guided by the availability of existing results and do not aim to reflect theoretical preferences for any specific models. For minimal quark-philic models, most studies have focused on couplings to right-handed quarks of the first and third generations. Similarly, investigations of minimal leptophilic models have predominantly concentrated on couplings to muons.

4.4.1 Minimal simplified models

4.4.1.1 Minimal quark-philic models in the canonical freezeout regime

Figure 42 illustrates the cosmologically viable regions of the parameter space for six minimal models described in Sect. 2.1 in which the DM candidate couples to the righthanded up quark u_R , three of them with a real DM candidate and three of them with a complex one. Specifically, these include self-conjugate scalar DM (F3S, top left), Majorana DM (S3M, central left), real vector DM (F3V, bottom left), as well as complex scalar DM (F3C, top right), Dirac DM (S3D, central right), and complex vector DM (F3W, bottom right) candidates. The results are displayed in the plane defined by the DM mass M_X and the relative mass splitting M_Y/M_X-1 . Moreover, the grey-scale colour map indicates the value of the coupling λ required to achieve the observed DM relic density, $\Omega h^2 \simeq 0.12$ [178]. As the DM mass and mass splitting increase, the coupling value necessary to match the relic density also grows. Consequently, the white regions in the upper-right corners of the panels correspond to overabundant DM scenarios, where the annihilation cross section is insufficient within the perturbative regime of the coupling. Conversely, for small mass splittings, co-annihilation effects involving the mediator become increasingly significant. The white regions in the lower-left corners thus represent underabundant scenarios, where mediators remain in chemical equilibrium with the thermal bath and dominate the annihilation process due to the large associated cross section. Although this condition persists for couplings $\lambda \geq 10^{-4}$, cosmologically viable solutions also exist for smaller couplings of the order of 10^{-6} . In such cases, the relic density is determined via conversion-driven freeze-out [34], in which the above chemical equilibrium breaks down due to semiefficient conversion processes between the DM particle and the mediator. This regime, opening a new part of the parameter space, has not been explored in [25], despite that for $\lambda < 10^{-4}$ all astrophysical constraints are naturally evaded.

In the canonical freeze-out regime, direct detection constraints originating from spin-independent DM interactions with nuclei represent the strongest constraints across all six models. They are found to exclude the entire sampled parameter space for all three complex DM classes of scenarios, as shown in the right panels of Fig. 42. This leaves the conversion-driven freeze-out region as the sole viable regime for these models. In contrast, for self-conjugate DM models, portions of the parameter space remain unaffected by direct detection bounds. Nevertheless, combining constraints from direct detection (both spin-independent and spin-dependent bounds), indirect detection via gamma-ray and cosmic-ray antiproton observations, and robust limits from Z-boson visible decay measurements excludes significant portions of the parameter space, as depicted in the left panels of Fig. 42. For real scalar DM, these constraints exclude all scenarios with DM masses below 800 GeV or mediator masses below 2 TeV. For Majorana DM and real vector DM, a similar exclusion pattern arises, except for isolated allowed regions featuring scenarios with $M_X \lesssim 4 \,\mathrm{GeV}$ and $100\,\mathrm{GeV} \lesssim M_X \lesssim 200\,\mathrm{GeV}$ respectively. In the latter cases, cosmic-ray antiproton data provide key constraints for $200 \,\mathrm{GeV} \lesssim M_X \lesssim 800 \,\mathrm{GeV}$, similar to the scalar DM case. Here, indirect detection bounds are particularly stringent for vector and Dirac DM, due to the S-wave nature of the annihilation process into quark pairs. For additional results on these models, we also refer to [339].

We now turn to simplified models in which the DM state interacts with the right-handed top quark t_R . In the left panel of Fig. 43, we show the viable parameter space of a topphilic model featuring a Majorana dark matter state and a coloured scalar mediator (namely the S3M_tR class of models), as obtained in [201]. Constraints on the parameter space are displayed in terms of the DM mass M_X and the mass splitting M_Y/M_X-1 between the mediator and the DM particle. At each point in the parameter space, the Yukawa coupling strength λ between the DM and the mediator is fixed to ensure that thermal freeze-out produces the observed DM relic density, $\Omega h^2 = 0.12$, the green contours being isolines of constant λ value.

The features at low DM masses arise due to resonant contributions from processes such as $XX \rightarrow h \rightarrow b\bar{b}$ via the loop-induced XXh coupling, as well as from the coannihilation channel $XY \rightarrow t \rightarrow Wb$. Additionally, $2 \rightarrow 3$ processes like $XX \rightarrow Wtb$, which are relevant below the $t\bar{t}$ threshold, and loop-induced processes like $XX \rightarrow gg$, are included in our calculations. For mediator annihilation $Y\bar{Y} \rightarrow gg$, Sommerfeld enhancement is accounted for, following [340], although bound-state effects are not included. In the lower-right region, below the black solid line, the DM relic density can only be explained via conversion-driven



Eur. Phys. J. C (2025) 85:975 Page 67 of 96 975




Fig. 42 Constraints on minimal simplified t-channel DM models from cosmological and astrophysical observables, as well as from the measured Z-boson visible decay width. The coloured region in the $(M_X, M_Y/M_X-1)$ plane represents scenarios that achieve $\Omega h^2 \simeq 0.12$, with the coupling value λ indicated by the grey-scale colour map. The left (right) panels correspond to models with self-conjugate (complex) DM, featuring scalar (top row), fermionic (middle row), and vector (bottom row) DM particles. The hatched regions indicate exclusions from gamma-ray searches ('ID gamma rays', including gamma-line searches from Fermi-LAT [353] and HESS [403] from the galactic cen-

tre and gamma-ray continuum searches in dSPhs by Fermi-LAT [348]), cosmic-ray antiproton searches ('ID antiprotons'), DM direct detection via spin-independent and spin-dependent interactions ('DD SI' and 'DD SD,' respectively, including limits from LZ [404], PICO-60 [377], CRESST-III [405] and DarkSide-50 [406]), and Z-boson visible decays ('Z decay', from [407]). The blank upper region corresponds to scenarios requiring non-perturbative couplings, while the white region at the bottom represents the CDFO regime (see Sect. 4.4.1.2). This figure is taken from [25], where further details can be found



975 Page 68 of 96 Eur. Phys. J. C (2025) 85:975





Fig. 43 Cosmologically viable region of the parameter space of models featuring a DM state coupling to the right-handed top quark t_R (with $\Omega h^2 = 0.12$), and constraints shown as functions of the DM mass M_X and the relative mass splitting $M_Y/M_X - 1$. The left panel corresponds to Majorana DM, while the right panel depicts a scenario with real scalar DM. These plots are based on the results from [201] and [27], respec-



tively, and the direction detection (DD) constraints on both plots are derived from the XENON1T bounds. The thick black line separates the WIMP region (above) from the CDFO region (see Sect. 4.4.1.2), and the shaded regions denote exclusions due to the various experimental and theoretical constraints discussed in the text

freeze-out, which will be discussed in Sect. 4.4.1.2. In addition, the dark grey region at large relative mass splittings is excluded due to DM overproduction, and the purpleshaded region is excluded by direct detection constraints from XENON1T [408]. Here, the loop-induced XXh and XXgg couplings originating from box diagrams play a critical role as they mediate DM-nucleon scattering through partonic processes like $Xg \rightarrow Xg$. The 'blind spot' in the direct detection constraints, visible at mass splittings of approximately 100 GeV in the left panel of Fig. 43, arises from destructive interference between these loop-induced contributions. Finally, indirect detection limits are relevant only within a very narrow band near the Higgs resonance at $M_X \simeq M_h/2$, and are thus not shown here. In this region, DM annihilation predominantly proceeds via the loop-induced DM-Higgs interaction, as detailed in [201].

In the right panel of Fig. 43, we move on with a study of the constraints that can be imposed on the parameter space of a simplified model with a real scalar DM candidate and a top-philic vector-like fermion mediator, as examined in [27]. In the white region above the black solid line, the measured DM abundance can be achieved with WIMP-like couplings ($\lambda \sim 10^{-2}$ to O(1)) through canonical freeze-out. Below this line, the DM relic density can instead be explained via the CDFO mechanism, which requires much weaker λ coupling values. In the WIMP region, constraints from both direct and indirect DM searches exclude substantial portions of the cosmologically viable parameter space, as visible by the parameter space regions excluded when combining direct detection constraints from XENON1T [408], indirect detection con-

straints from AMS-02 antiprotons [350], and the projected sensitivity of Fermi-LAT after 15 years of exposure [409]. Here, the related calculations necessitated a detailed treatment of radiative corrections, which was provided in [341], and that accounts for the non-negligible mass of the final-state SM particles in the relevant processes. NLO processes are particularly important during freeze-out as the LO annihilation cross section is D-wave suppressed [337]. In fact, for any t-channel DM models where the DM couples exclusively to the third generation, QCD corrections must be considered [27]. For instance, in all scenarios where the DM couples to the right-handed top quark t_R , loop-induced DM annihilation into gluons dominates and determines the relic density below the b-quark threshold [201].

Finally, we turn on the simplified models featuring universal couplings to all generations of SM fermions. In Fig. 44, we present the exclusion limits on the parameter space for the DM model S3M_uni with a SM-singlet Majorana DM candidate. This candidate couples universally to all SM uptype right-handed quarks via three mass-degenerate mediators, thus with a unique coupling strength λ . As above and following [302], the constraints on the parameter space are shown in the plane defined by the DM mass and the mass splitting between the mediator and the DM particle. At each point in the plane, λ is fixed to ensure that the observed DM relic density, $\Omega_{\rm DM} = 0.12 \pm 0.005$, is not exceeded. The relic density calculation takes into account LO annihilation and co-annihilation processes, Sommerfeld effects, and bound state formation. In the grey-shaded region, freeze-out underproduces DM, requiring alternative production mecha-



Eur. Phys. J. C (2025) 85:975 Page 69 of 96 975




Fig. 44 Constraints on a simplified DM model featuring three massdegenerate mediators universally coupling to a Majorana DM candidate and all generations of SM right-handed up quarks. The blue solid lines are isocontours with fixed λ , as labelled. The black-shaded region represents parameter values where avoiding DM overproduction would violate perturbative unitarity. The green and magenta areas show exclusions from spin-independent and spin-dependent direct detection experiments, respectively, and the grey-shaded region corresponds to a DM relic density explained through the CDFO mechanism. The results incorporate the effects of bound states, including their excitations, on freeze-out dynamics. This plot is adapted from [302]

nisms to account for the observed DM abundance. Although such mechanisms are not explicitly addressed for this model, their phenomenology is expected to align with the CDFO and freeze-in/superWIMP regimes discussed in Sects. 4.4.1.2 and 4.4.1.3.

The green-shaded region in Fig. 44 represents spinindependent direct detection constraints. For a Majorana DM candidate, all vector couplings to the SM neutral gauge bosons, including in particular those to the Z-boson that would otherwise lead to strong constraints, are identically zero. However, at the one-loop level, a DM-gluon coupling arises, yielding significant constraints at small mass splittings. These constraints are particularly sensitive for mass splitting values around the top quark mass $(M_Y M_X \sim M_t$), where a resonant behaviour from top quarks running in the loop amplifies the effect. The magentashaded region corresponds to spin-dependent direct detection constraints, which mainly arise from tree-level interactions. Since spin-dependent limits are typically much weaker than spin-independent ones, they generally play a subdominant role unless the mass splitting is relatively large or the DM mass is small. However, spin-dependent constraints could become more significant in the future, especially with improved experimental sensitivities such as those projected for DARWIN, where they may even dominate for large mass splittings. Lastly we emphasise that Fig. 44 provides enough information to complementarily combine collider and cosmology exclusions, by appropriately incorporating the results from the collider section. For instance, the bounds determined in Figs. 1 and 3 can be directly applied in the large λ regime where first-generation mediator production dominates the collider signal.

4.4.1.2 Minimal quark-philic models in the CDFO regime In all models discussed in Sect. 4.4.1.1, the cosmologically viable parameter space that explains the relic density via canonical freeze-out is limited to regions featuring not too small mass splittings (up to a certain DM mass well in the TeV range). In the small mass splitting regime, the DM density can be explained instead by the CDFO mechanism [34,35], as described in Sect. 4.2.2. This region is left blank in Fig. 42 and marked accordingly in Figs. 43 and 44. At the boundary between the canonical freeze-out and CDFO regimes, the required couplings decrease by several orders of magnitude, from a WIMP-like magnitude (10^{-2} to O(1)) above the boundary to approximately 10^{-6} deep in the CDFO region. Consequently, the lifetime of the mediator Y increases sharply, rendering it long-lived with striking implications for its signatures at colliders that we have discussed in Sect. 3.3. Since the CDFO scenario has been studied in detail in the literature for bottom-philic and top-philic models, the following discussion focuses solely on these cases. For mediators coupling to first-generation or second-generation quarks, the phenomenology is nevertheless expected to be very similar to the bottom-philic case.

For bottom-philic Majorana DM scenarios with a coloured scalar mediator (namely the S3M_bR class of models), the constraints on the parameter space are shown in Fig. 45. In the region below the thick black line, the observed relic density can be explained within the CDFO scenario. The required values for the Yukawa coupling λ , of the order of 10^{-7} , are indicated by green dashed contours. Isolines of constant decay length of the mediator are also shown, this time through grey dotted lines. In the bulk of the parameter space region above the kinematic threshold for the mediator's twobody decay, the decay length ranges from approximately 1 m (for small mass splittings) to 1 mm (near the upper boundary of the CDFO regime). Below this threshold, it becomes significantly larger, often exceeding the dimensions of typical LHC detectors. Unlike in the canonical freeze-out regime, the CDFO parameter space is not subject to strong constraints from direct or indirect detection due to the very weak DM-SM coupling, rendering the LHC searches the unique probes for this scenario. On the cosmological side, the dilution of dark sector particles is driven solely by mediator-pair annihilation. Thermal decoupling is a prolonged process, extending



975 Page 70 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 45 Parameter space regions of a bottom-philic Majorana fermion DM model [258] compatible with the observed relic density. The left panel shows the required couplings and the resulting decay length of the mediator, with the CDFO region lying below the thick black line. The results account for Sommerfeld enhancement and excited bound state effects. The right panel highlights the significance of bound state

effects in this scenario, showing the CDFO contour's boundary (solid lines) along with isolines of constant λ values in the canonical freezeout regime (dashed lines). Results including Sommerfeld enhancement and bound state effects are shown in blue, while those without these effects are shown in red

deeper into the non-relativistic regime than in the conventional freeze-out case. As a result, non-perturbative effects, such as Sommerfeld enhancement and bound state formation, play a particularly significant role [258]. These effects substantially extend the size of the CDFO regions in the parameter space, as illustrated in the right panel of Fig. 45. The red lines represent LO tree-level results, while the blue lines incorporate Sommerfeld enhancement and the effects of excited bound states, including states up to a principal quantum number of n = 15.

For top-philic models, the phenomenology exhibits distinctive features by virtue of the large top mass M_t . The regions of the parameter space relevant for the CDFO regime corresponds to the one below the solid black line in Fig. 43. Here, we focus on the Majorana DM case explicitly studied in [201], as the corresponding CDFO region for scenarios with real scalar DM solely extends to slightly higher masses and does not exhibit any different feature. A key difference from the bottom-philic scenario is that the two-body decay of the mediator is kinematically forbidden throughout the entire CDFO region. Since $M_Y - M_X < M_W + M_b$, the mediator's decay instead proceeds via very suppressed four-body channels. Consequently, the decay length of the mediator is large compared to the dimensions of the LHC detectors across the entire CDFO parameter space, despite the larger λ coupling values relevant for top-philic scenarios. However, the large lifetimes of the mediator also imply that certain regions of the parameter space are inconsistent with successful BBN. These regions are shaded in red in the figure. Contours of constant coupling strength are additionally shown in green, ranging from values of approximately 10^{-3} for $M_X \lesssim M_t$ to 10^{-6} for $M_X \gg M_t$. While these results include Sommerfeld enhancement, bound state effects are not accounted for in this analysis. As demonstrated in the bottom-philic case, such effects are expected to further enlarge the CDFO parameter space.

4.4.1.3 Minimal quark-philic models in the freeze-in and superWIMP regimes

For couplings smaller than those considered in the CDFO regime, DM does not thermalise in the early universe and we must rely on freeze-in and superWIMP DM production to explain the measured relic density, as detailed in Sects. 4.2.3 and 4.2.4. In the class of t-channel mediator models under consideration, both production mechanisms are present, though their relative importance can vary significantly depending on the specifications of the model and the benchmark point in the parameter space. The following discussion focuses on the case of top-philic fermionic DM. For mediators coupling to first-generation or second-generation quarks, the cosmologically viable parameter space is expected to be similar for large mediator masses $(M_Y \gg M_t)$ but differs for $M_Y \lesssim 1$ TeV. Moreover, scalar DM scenarios are expected to lead to qualitatively similar results [252].

In Fig. 46, we display the cosmologically viable regions of the parameter space associated with the top-philic Majorana DM setup considered. The results, based on [204], are shown in the $(M_X, M_Y - M_X)$ plane as a function of the coupling value λ . They assume no additional contributions to the



Eur. Phys. J. C (2025) 85:975 Page 71 of 96 975




Fig. 46 Regions of the top-philic Majorana DM model compatible with DM production through the freeze-in and superWIMP regime, and existing cosmological constraints as studied in [204]. The plot highlights the interplay between these two DM production processes across the parameter space (long dashed lines). Relevant constraints from structure formation (purple) and BBN (red) are also indicated. In addition, isolines of constant λ values and of constant mediator decay length are shown in green and grey, respectively

DM abundance prior to infrared freeze-in (e.g. from postinflationary reheating processes) and a reheating temperature above the mediator mass scale $T_{\rm RH} \gg M_Y$. The measured relic abundance is reproduced in the regions of the parameter space extending to the left of the solid black curve, whereas to the right of this curve, superWIMP production alone would exceed the observed relic density. In addition, long-dashed black lines indicate the relative contributions of freeze-in and superWIMP production to Ωh^2 , with the freeze-in contribution increasing as one moves away from the overproduction boundary, and thin green lines are isolines of constant λ values. For masses around the TeV scale that are typically relevant for collider searches, λ ranges between 10^{-8} and 10^{-12} . Those small values imply large mediator lifetimes, as shown through the light dashed lines demonstrating that TeV-scale mediators have macroscopic decay lengths.

Cosmological constraints are indicated by the different coloured regions. The purple area is excluded by structure formation constraints via Lyman- α observations, and the red area is inconsistent with successful BBN. Here, all the results include Sommerfeld enhancement and bound-state effects, although they only account for the impact of the ground bound state [204]. Higher bound-state excitations are expected to be significant as they reduce the superWIMP contribution, thereby expanding the cosmologically viable region of the parameter space towards larger DM masses

[261]. Finally, we should keep in mind that thermal effects, which are particularly relevant in the freeze-in regime, have only been studied recently in [284], as noted in Sect. 4.2. The full parameter space incorporating these effects has yet to be mapped.

The above results change if the assumption $T_{\rm RH}\gg M_Y$ is relaxed to consider lower reheating temperatures. For simplified t-channel DM models, such scenarios with low reheating temperatures have been investigated in [252,254,256]. These studies examine freeze-in production for both Majorana and scalar DM models with scalar and vector-like fermion mediators, respectively. If $T_{\rm RH} < M_Y$, meaning that DM freezes in during reheating, the entropy dilution that reduces the comoving number density leads to larger Yukawa couplings between DM, the mediator, and the SM particle with which they interact in order to reproduce the observed DM relic density. These larger couplings shorten the lifetime of the long-lived mediator, bringing it into a range testable via displaced-vertex searches, as discussed in Sect. 3.3.

4.4.1.4 Minimal quark-philic models: discussion and conclusion

We can summarise our findings and discussion with the following general remarks. In minimal models, the quark flavour to which the DM couples has only a minor effect on the relic density across all DM production mechanisms, provided that the DM mass and the DM-mediator mass splitting are significantly larger than the corresponding quark mass. This condition is typically satisfied for most flavour choices. However, in the top-philic case, parts of the parameter space exhibit unique features due to the top quark large mass and, in some cases, its substantial Yukawa coupling to the SM Higgs boson. In the canonical freeze-out regime, direct detection experiments impose the strongest constraints on DM, particularly for couplings to first-generation quarks. Here, these experiments exclude significant portions, or even all, of the cosmologically viable parameter space. In contrast, models featuring couplings to third-generation quarks are less constrained by direct detection. Astrophysical observations thus provide a valuable complement to collider searches for freezing-out DM, as they remain sensitive to very heavy DM candidates that are hard (or even impossible) to be efficiently produced at colliders. In the CDFO and freeze-in/superWIMP case, conventional DM searches through direct and indirect detection offer limited prospects due to the extremely weak coupling required. However, certain regions of the parameter space predicting a very longlived mediator are constrained by Big Bang Nucleosynthesis. These constraints apply to specific realisations of topphilic CDFO models and to portions of the parameter space related to the freeze-in/superWIMP regime that is also further constrained by cosmological structure formation, particularly Lyman- α forest observations. We additionally point




out that models where DM couples to second-generation quarks remain relatively under-explored. Nevertheless, their astroparticle phenomenology is expected to broadly resemble that of models involving first-generation quarks. This contrasts with collider physics, where charm quarks play a distinctive role due to factors such as charm tagging, uncertainties in parton distribution functions, and other collider-specific considerations.

In the *t*-channel mediator models considered in this work, DM is assumed to be a singlet under the SM gauge group. Relaxing this assumption introduces a broader class of models featuring t-channel mediators. Examples include in particular the Minimal Dark Matter framework [31], where DM lies in a non-trivial $SU(2)_L$ multiplet. Among the most studied realisations are the $SU(2)_L$ doublet ('higgsino' DM) and triplet ('wino' DM) scenarios. Detailed phenomenological studies can be found, for instance, in [410-414], and also in [415–418] in light of concurring excesses in related LHC searches. The DM observables and related constraints in the case of models with non-trivial $SU(2)_L$ multiplets differ significantly from the gauge-singlet case. For instance, the small mass splitting among states within a multiplet, typically in the 100 MeV range, often leads to LLP signatures driven by phase-space suppression like in higgsino and wino models. While these specific models are not discussed in detail here, Sect. 3.3 addresses analogous signatures in the considered t-channel mediator models, where long lifetimes are instead induced by the small couplings required in the CDFO, freezein or superWIMP regimes. To maintain focus, we have indeed restricted the previous discussion to gauge-singlet models, and we refer instead to [4] for a comprehensive overview of electroweak multiplet models. Specific non-minimal explorations are then conducted in the last part of this section, all highlighting interesting phenomenology not covered in the minimal framework.

4.4.1.5 Minimal leptophilic models

In this section, we explore leptophilic minimal models, focusing specifically on scenarios with a Majorana DM candidate and a scalar mediator with couplings to right-handed muons (i.e. the S1M_muR class of models). The phenomenology of models with couplings to right-handed electrons or taus is expected to be qualitatively similar, provided the DM and mediator masses satisfy M_X , $M_Y - M_X \gg M_\tau$. For a comprehensive overview of these models, we refer to [57,251,252,260,339,419–428].


The canonical freeze-out regime provides an interesting starting point to examine these models. The left panel of Fig. 47 illustrates the viable parameter space and projected constraints, as presented in [339], under the assumption that the DM relic abundance is achieved through thermal freeze-out. In the upper-right region (grey shading), DM is over-produced unless couplings become non-perturbatively large





For small mass splittings, the CDFO regime becomes relevant, requiring very weak DM couplings of $O(10^{-6})$. This region of the parameter space is represented in the lower-left corner of the left panel of Fig. 47, with the thick black line marking the transition from the WIMP regime to CDFO regime. The measured relic density is still reproduced, but the required coupling value drops by several orders of magnitude. In the model with a scalar mediator considered here, the mediator pair-annihilation cross section depends on electroweak contributions from γ/Z exchanges and potential Higgs-portal interactions, with their relative importance impacting the position of the boundary of the CDFO regime. While the Higgs portal contributions are taken vanishing in the left panel of the figure, introducing a sizeable Higgsportal coupling, such as $\lambda_H = 0.5$, significantly expands the viable CDFO parameter space. This is illustrated in the right panel of the figure, where we can consider CDFO scenarios with DM masses ranging up to above 1 TeV [260]. The green thick line in the figure denotes the new boundary of the CDFO regime, extending hence the relevant region of the parameter space significantly beyond 1 TeV. Additionally, blue thin lines represent isolines of constant DM-SMmediator coupling values (multiplied by 10⁶). It is important to note that non-perturbative effects, including Sommerfeld enhancement and bound state formation, could substantially further enlarge the region of the parameter space rele-



Eur. Phys. J. C (2025) 85:975 Page 73 of 96 975







shown in the lower left corner. Projected constraints from indirect and direct detection are shown through the dashed blue and purple areas, while LEP limits are given in green. The figure in the right panel is dedicated to the CDFO regime and is adapted from [260]. Here, the Higgs-portal coupling is set to $\lambda_H=0.5$, and the blue lines are isolines of constant $\lambda/10^{-6}$ values

vant for the CDFO regime by modifying the mediator pairannihilation cross section. While these effects have not been included in the results of Fig. 47, they are expected to follow the trends observed for quark-philic models [258]. Additionally, the small coupling in the CDFO regime ensures that direct and indirect detection constraints remain negligible. Testing this scenario at colliders, however, is possible via LLP signatures, as discussed in Sect. 3.3.

Finally, the freeze-in and superWIMP regimes arise for even smaller couplings, where DM production occurs out of equilibrium. These scenarios, characterised by extremely weak couplings, are not testable via direct or indirect detection experiments. Instead, the long-lived nature of the mediator *Y* leads to LLP signatures at colliders, as detailed in Sect. 3.3. Although the cosmologically viable regions of the parameter space of leptophilic models in the freeze-in/superWIMP regime have not been fully mapped, as for quark-philic models, they are expected to qualitatively resemble the latter case, albeit with different mass scales due to weaker mediator-SM interactions [252].

4.4.2 Non-minimal models

4.4.2.1 Flavoured dark matter

The presence of multiple DM flavours significantly influences early-universe cosmology and shapes the experimental constraints on these models. In flavoured DM frameworks, it is common to consider three generations of DM particles X_i with a mass hierarchy $M_{X_1} > M_{X_2} > M_{X_3}$. The lightest flavour X_3 is assumed to be stable and constitutes the

observed DM relic density, while the heavier flavours $X_{1,2}$, as well as the mediator Y, are unstable and decay into X_3 and SM particles. The cosmological evolution of such models depends on the representations of the X_i and Y states and on the structure of the coupling matrix λ linking the SM fermions to the dark sector. Furthermore, in the context of the DMFV models introduced in Sect. 2.2, the matrix λ not only governs the texture of the interactions but also determines the DM mass spectrum, as the mass matrix $M_{X,ij}$ can be expressed in terms of a spurion expansion in $\lambda^{\dagger}\lambda$.

Flavoured DM has been explored in the regimes of both canonical freeze-out and conversion-driven freeze-out. Early studies [26,73-76,78,79] of DMFV models have largely focused on two limiting scenarios of freeze-out. In the first called single-flavour freeze-out (SFF), the flavour X_3 is sufficiently separated in mass from the other states $X_{1,2}$ and Ysuch that co-annihilation effects are negligible. In the second that was named quasi-degenerate freeze-out (QDF), the X_i flavours are all nearly degenerate, and their combined annihilations contribute equally to the freeze-out process. More recent analyses, however, have incorporated general co-annihilation effects into the study of both quark-flavoured and lepton-flavoured DM [77,80]. Furthermore, in addition to canonical freeze-out, the case of CDFO has been studied in a flavoured Majorana DM model coupled to right-handed uptype quarks u_{Ri} [77]. The inclusion of heavier dark flavours $X_{1,2}$ and the additional coupling parameters λ_{ij} have been found to relax the constraints on the viable regions of the parameter space imposed by the relic abundance condition in comparison to unflavoured models. Notably, an inverse




975 Page 74 of 96 Eur. Phys. J. C (2025) 85:975

 10^{2}

 10^{-2}

 $M_Y/M_{X_3} - 1$ [%]




100 200 500 1000 2000 M_{X_3} [GeV] matter and the mediator Δm_{Y_3} . On the left panel, the (co-)annihilation freeze-out regime is considered, with QDF scenarios shown in green, SFF ones in yellow, and generic ones in blue. On the right panel, the CDFO regime is examined instead, with conversions between X_3/X_2 (yellow), X_3/Y (blue), and combined $X_3/X_2/Y$ (green)

Fig. 48 Viable regions of the parameter space of flavoured Majorana DM scenarios [77] in which the dark matter couples to right-handed up quarks. Constraints from the observed dark matter relic abundance, direct and indirect detection experiments, and flavour data are shown as a function of the DM mass m_{X_3} and the mass splitting between the dark

mass hierarchy within the dark sector (with the η parameter of (7) being greater than zero), where the lightest state X_3 has the weakest coupling to visible matter, offers a compelling scenario. Here, the relic density relies predominantly on the annihilation of the heavier DM flavours and/or the mediator, and this configuration naturally evades direct and indirect detection experiment bounds. The latter indeed probe only the interactions of the X_3 state, and DM is thus hidden from these searches.

Figure 48 displays the allowed regions of the parameter space for scenarios featuring flavoured Majorana DM coupled to the right-handed up quarks u_{Ri} . On the left panel of the figure, the standard (co-)annihilation freeze-out scenario is examined, while the right panel depicts various CDFO scenarios [77]. Accounting for co-annihilation effects (blue points) significantly broadens the size of the viable region of the parameter space compared to the SFF (yellow points) and QDF (green points) benchmark cases. However, small mass splittings between the X_3 state and the mediator Y remain excluded in the canonical freeze-out scenario. In the CDFO regime, where semi-efficient annihilation between X_3 and Yoccurs, this region of the parameter space becomes in contrast viable (green, blue). Moreover, for small splittings between X_3 and the heavier flavours, the conversion between these states can also become semi-efficient, enabling additional viable parameter space regions for CDFO scenarios involving the heavier dark flavours (yellow). Intriguingly, flavoured DM models with a CDFO realisation offer the possibility of



The typical constraints that could be imposed on flavoured DM arise from a combination of direct and indirect detection experiments, electroweak precision measurements, LHC searches, and flavour physics. In the latter case, the most stringent limits typically stem from neutral meson mixing observables for quark-flavoured DM [26,73-76] and from radiative decays such as $\ell_i \to \ell_i \gamma$ for lepton-flavoured DM [78–80]. However, these flavour observables constrain the structure of the coupling matrix λ rather than the overall mass or coupling scale. In addition, direct and indirect detection constraints are often relaxed relative to minimal non-flavoured models. This occurs due to the extra parametric freedom provided by the flavour structure of the coupling matrix λ , which can allow cancellations between tree-level and loop-level contributions, thereby suppressing the DM-nucleon scattering cross section [26,74]. Consequently, flavoured DM serves as a concrete realisation of the xenophobic DM paradigm [430]. Finally, constraints from indirect detection are found generally weaker than those from direct detection experiments.

4.4.2.2 Frustrated dark matter

The fDM framework [30] introduced in Sect. 2.4 describes a family of non-minimal models in which a fermionic dark matter state X couples via a Yukawa-like interaction to a pair of mediators $\{\varphi, \psi\}$. At least one of these mediators interacts with the Standard Model, and to satisfy gauge invariance,



Eur. Phys. J. C (2025) 85:975 Page 75 of 96 975

they carry the same quantum numbers. While in Sect. 3.2.3 we provided a detailed review of the collider implications the fDM framework, we now focus on its cosmological consequences. Frustrated DM indeed exhibits intriguing astrophysical phenomenology, combining compatibility with the observed relic density and promising prospects for detection via both direct and indirect searches for DM.

By construction, $2 \rightarrow 2$ interactions between the DM and the SM occur only at one-loop order. Consequently, for most DM masses, DM annihilation is dominated by XX scattering to at least four SM particles, which effectively reduce to t-channel XX annihilations to a pair of mediators when the mediators can be produced nearly on-shell. For perturbative Yukawa-like DM-mediator couplings $\lambda_X \lesssim \sqrt{4\pi}$, efficient DM annihilation requires these t-channel processes to be kinematically accessible; otherwise, DM tends to be overabundant under the assumption of a standard cosmological history. However, the fDM framework allows for a wide range of mediator masses and SM couplings, enabling the construction of numerous specific realisations with large experimentally viable regions in the model parameter space. The regions of the fDM parameter space that can produce a viable thermal relic are, in principle, testable via indirect searches for DM annihilation in the cosmos. While loopsuppressed $2 \rightarrow 2$ annihilation processes leading to photons or a γZ final state are possible, the dominant experimental signature of the models is typically an apparent excess in the continuum spectra of gamma or cosmic rays originating from DM annihilation into quarks and/or charged leptons. For fDM scenarios at or below the TeV scale, the strongest constraints thus come from Fermi-LAT searches for gamma-ray production in nearby dwarf spheroidal galaxies. On the other hand, direct detection is fully loop-suppressed since there are no tree-level diagrams for fDM scattering off SM particles. Nonetheless, constraints from direct detection experiments such as XENON1T can be quite stringent. In scenarios where mediators carry weak hypercharge, spin-independent DMnucleon scattering via off-shell photon exchange then provides the most robust constraints, that can also be interpreted as upper limits on the DM magnetic dipole moment [368]. In the absence of such processes, box diagrams involving mediators and SM fields can sometimes generate detectable direct detection signals [366] that may lead to competitive or even exceed limits from indirect detection. More conclusive quantitative statements require, however, specifying a particular fDM construction.

In Sect. 3.2.3, we proposed a benchmark scenario that was ideal for LHC exploration. In this scenario, the DM is a Dirac fermion, and the mediators are QCD colour sextets. The sole renormalisable model of this type features a colour-sextet scalar φ coupling to pairs of like-sign quarks, while the colour-sextet fermion ψ is sequestered from the SM, interacting only through usual gauge interactions. In the




Fig. 49 Limits from direct and indirect searches (XENON1T and Fermi-LAT) on a fDM realisation featuring Dirac DM and colour-sextet mediators. Here the scalar is taken to couple to pairs of like-sign up-type quarks, preferentially to the third generation

simplest case, this can be achieved by imposing a \mathbb{Z}_2 symmetry under which ψ and X are odd while the SM states are even. Moreover, to avoid scenarios with stable colourful fermions, we impose the hierarchy $M_{\psi} > M_X$. Furthermore, experimental constraints from direct searches and measurements of flavour-changing neutral currents in neutral meson transitions [92,93] motivate a scenario in which the scalar mediator preferentially couples to third-generation quarks. Specifically, we thus assume that both quarks to which the mediator couples are of up-type, with dominant φqq couplings to ut and tt systems. Current constraints favour TeV-scale mediators, which in turn suggest TeV-scale DM masses $(M_X \gtrsim M_{\varphi})$ to ensure efficient annihilation.

To illustrate the cosmological constraints that could be imposed on such an fDM setup, Fig. 49 shows contours of $\Omega h^2 = 0.1198$ in the (M_X, M_{ψ}) plane. We consider different values of M_{φ} , and a DM-mediator coupling $\lambda_X = 1.25$ that is relatively large but remains within the perturbative regime. The figure also includes Fermi-LAT Pass 8 (dashed lines) and XENON1T 1 ton-year (dotted lines) constraints. Here, the Fermi-LAT limits act as lower bounds on M_X , while the XENON1T limits correspond instead to a lower bound on M_{ψ} . In the chosen scenario, the indirect detection limits are derived primarily from $2 \rightarrow 4$ annihilation processes as calculated with MadDM, while the direct detection bounds stem from one-loop off-shell photon-mediated DM-nucleon scattering, an effective field theory analysis, and our own calculation of the DM magnetic dipole moment [368]. In this specific realisation, frustrated DM remains a viable thermal relic across certain ranges of M_X and M_{ψ} values for all the considered mediator masses M_{φ} . While heavier mediator and DM mass spectrum correspond to regions of the parameter space less constrained by indirect detection, these setups turn out



975 Page 76 of 96 Eur. Phys. J. C (2025) 85:975





Fig. 50 Constraints on the parameter space of a composite *t*-channel DM scenario featuring one DM state *X* and one mediator state *Y* that couples to the top quark with a strength λ_t . Results are given in the (M_X, M_Y) plane, and the value of the coupling λ_t , given through the colour map, is determined in order to reproduce the observed relic density $\Omega_{CDM}h^2 = 0.1186$. The black lines correspond to isolines of constant coupling values with $\lambda_t = 1, 2, 3, 4$ and 5

to be complementarily probed, or even excluded, by ongoing direct detection experiments. It is however important to emphasise that these limits do not directly apply to all fDM models. The mass scales, interaction strengths, and experimental constraints can shift depending on the mediator's SM charge assignments or other model specifics. For instance, if the DM is a Majorana fermion instead of a Dirac fermion, the limits from direct detection are significantly altered, as a Majorana fermion does not possess a magnetic (and an electric) dipole moment.

4.4.2.3 Composite dark matter

Another compelling possibility for exploring *t*-channel dark matter models beyond minimal frameworks arises from composite scenarios such as those introduced in Sect. 2.3, that lead to a new physics particle spectrum including both even and odd new states, one of the latter playing the role of the DM. In the present section, we consider two distinct realisations of such composite models, and we discuss their implications for dark matter observables in terms of the model parameters. The results shown below highlight an interesting and non-trivial interplay between the couplings and the mass parameters, offering insights into the underlying dynamics of composite models and their implications for DM phenomenology.

In the first case that we consider, the only non-vanishing free parameters of the model comprise a single coupling λ_t that appears in the Lagrangian (8), the DM mass M_X , and the mediator mass M_Y . To estimate the DM relic density including NLO effects, we employ MadDM with the model implementation documented in Sect. 2.3. We perform a parameter scan in which M_X ranges between 200 GeV and 3000 GeV, M_Y varies between 200 GeV and 6.5



We next focus on a second class of scenarios in which the model is extended to include the two couplings λ_t and λ_t' of the Lagrangian (8), the two mediator masses M_Y and $M_{Y'}$, and the DM mass M_X . We perform a parameter space scan in which M_X is varied between 200 GeV and 3000 GeV, λ_t and λ'_t between 0.1 and 6, and M_Y and $M_{Y'}$ between 200 GeV and 3500 GeV. Two specific mass hierarchies are considered, $M_{Y'} < M_X < M_Y$ and $M_X < M_{Y'}$, M_Y , and the results are given, in the left and right columns of Fig. 51 respectively, in the coupling plane (λ_t, λ_t') . The colour code represents the masses of the different states required to reproduce the relic density as observed by the Planck collaboration for any given coupling configuration. When $M_{Y'} < M_X < M_Y$, λ_t can reach values up to 2.1, while λ_t' remains below 0.6, regardless of the values of M_X , M_Y , and $M_{Y'}$. Parameter combinations show a concentration of viable scenarios featuring small λ_t values (around 0.5) and λ_t' values (between 0.1 and 0.3) when $M_X < 1000$ GeV, $M_Y < 1500$ GeV, and $M_{Y'}$ < 600 GeV. Spectra featuring larger DM masses $(M_X \sim 2000 \text{ GeV})$ and mediator masses $(M_Y \sim 3000 \text{ GeV})$ or $M_{Y'} \sim 1800$ GeV) are also viable, but in this case they require combinations of higher λ_t values (around 2) with smaller λ'_t values (below 0.4 or between 0.1 and 0.6). On the other hand, when $M_X < M_{Y'}, M_Y, \lambda'_t$ can reach larger values than λ_t , with λ_t' going up to 5.5 and λ_t up to 3.6, independent of the masses M_X , M_Y , and $M_{Y'}$. For $M_X < 400$ GeV, λ_t values cluster around 3 to 3.5, while λ'_t remains below 5.5. For mediator masses $M_Y > 2500 \text{ GeV}$ or $M_{Y'} > 2000 \text{ GeV}$, viable combinations include $\lambda_t > 2$ with λ_t' below 5.5 and lower values of both couplings. For these composite models a fully phenomenological investigation taking into account the relevant dark matter constraints from astroparticle searches is still missing, and thus deserves future studies.

4.4.2.4 Non-Abelian dark sector portal

In this section, we consider the simplest realisation of the FPVDM model described in Sect. 2.5, assuming that new vector-like fermions interact with only one SM flavour that we take to be the top quark, and that no mixing occurs between the SM Higgs boson h and the new scalar H. Moreover, the hierarchy in the fermion sector follows $m_t < m_{tD} \le m_T$, and H can have any mass value allowed by data. We test the above setup against multiple DM observables, importing our implementation within micromegas [431]. The relic



Eur. Phys. J. C (2025) 85:975 Page 77 of 96 975




Fig. 51 Constraints on the parameter space of a composite t-channel DM scenario featuring one DM state X, two mediator states Y (that is \mathbb{Z}_2 odd) and Y' (that is \mathbb{Z}_2 -even), and the two new physics couplings λ_t and λ_t' of the Lagrangian (8). Results are given in the (λ_t, λ_t') plane, and the value of the masses in GeV, represented through the colour code,

are determined so that the relic density matches observations. The top, central and bottom rows respectively address the masses M_X , M_Y and $M_{Y'}$, and we consider the mass hierarchies $M_{Y'} < M_X < M_Y$ (left) and $M_X < M_{Y'}$, M_Y (right)



975 Page 78 of 96 Eur. Phys. J. C (2025) 85:975




Fig. 52 Representative Feynman diagrams for *t*-channel and resonant contributions to DM annihilation and DM-mediator co-annihilation processes (top), and processes relevant for direct detection experiments (bottom)

density is determined by the interplay of annihilation and co-annihilation processes, some of which being represented in the top row of Fig. 52. Indirect detection constraints are tied to DM annihilation rates during the CMB epoch, excluding regions of parameter space where energy injection into the SM plasma in the early universe is inconsistent with data. In the procedure that we follow to extract bounds on the model, both relic density and indirect detection observable predictions are tested against Planck data [178]. Finally, direct detection constraints are also assessed, and in the considered FPVDM scenario they are associated with processes such as those in the bottom row of Fig. 52. Limits are this time determined by confronting our predictions against the rsults of the XENON1T collaboration [371].

The regions of the parameter space compatible within 5% with the relic density as measured by the Planck collaboration are shown by the green, cyan, and blue areas in Fig. 53. In this figure, the results of a comprehensive scan of the parameter space of the model are projected in the (m_{V_D}, g_D) plane to highlight their dependence on the dark gauge boson mass and coupling. These regions correspond to a relic density driven by DM annihilations dominated by tchannel diagrams, resonantly-enhanced H contributions and DM-t_D co-annihilations, respectively. Generic DM annihilations induced by the t-channel diagrams set a lower limit on the dark gauge coupling g_D as a function of m_{V_D} , while scenarios exhibiting an H-resonant enhancement allow g_D to be reduced by up to two orders of magnitude. Furthermore, the strong DM- t_D co-annihilation channel permits even lower values of g_D for moderately heavy DM. For $m_{V_D} > 2$ TeV, however, the co-annihilation mechanism saturates, while Hresonant annihilation requires larger g_D values for increasing DM masses to maintain the observed relic density. As a result, the region with $m_{V_D} \gtrsim 2$ TeV typically corresponds to overabundant DM (indicated by the dark red area in the




Fig. 53 Excluded and allowed parameter space regions of the considered FPVDM realisation, as obtained from a full five-dimensional scan of the parameters in (24). Results are projected in the (m_{V_D}, g_D) plane and include scenarios compatible with observations by virtue of t-channel annihilations (green), H-funnel resonant contributions (cyan) and co-annihilations (blue), as well as under-abundant (grey) and overabundant (red) DM setups. Constraints from indirect and direct detection are additionally displayed through the orange and magenta regions, while white areas represent a non-perturbative regime

figure), except for scenarios with large g_D values involving $V_D V_D \rightarrow V' V'$ or H-funnel annihilations. Finally, regions with $m_{V_D} \lesssim 2$ TeV are additionally partially excluded by direct and/or indirect detection constraints, as indicated by the magenta and orange points, respectively.

To facilitate comparison with the LHC bounds discussed in Sect. 3.2.4, we present in Fig. 54 the cosmological constraints projected onto the (m_{t_D}, m_{V_D}) plane for two benchmark scenarios. Both benchmarks share the parameter values $m_H = 1000$ GeV and $m_T = 1600$ GeV, but differ in the choice of the gauge coupling g_D , which is set to 0.05 and 0.3 in the left and right panel of the figure respectively.



Eur. Phys. J. C (2025) 85:975 Page 79 of 96 975




Fig. 54 Excluded regions of the considered FPVDM realisation by cosmological observables projected onto the (m_{tD}, m_{VD}) plane for $m_T = 1600$ GeV, $m_H = 1000$ GeV and $g_D = 0.05$ (left) or 0.3 (right). Non-perturbative and large kinetic-mixing regions are also shown as hatched areas

This projection clearly illustrates how smaller gauge couplings significantly restrict the allowed region of the parameter space, leaving only regions where dark matter annihilation is enhanced by H-resonance effects or co-annihilation processes. The correct relic density as observed by the Planck collaboration is achieved at the boundary of the overabundant regions.

Our results demonstrate that the simplest FPVDM realisation that connects a vector DM candidate to SM fermions via a non-Abelian $SU(2)_D$ gauge group without requiring a Higgs portal at tree level, thus involving dark sector interactions with a single SM fermion, has significant potential to explain DM phenomena. Furthermore, alternative realisations involving interactions with other SM fermions could address a large set of observed anomalies. For instance, if the vector-like fermion interacts with the SM leptonic sector, it might contribute to explaining the $(g-2)_{\mu}$ anomaly [432] and open novel opportunities for future e^+e^- colliders [433-436]. Moreover, non-minimal scenarios, incorporating scalar sector mixing, extra vector-like partners, or interactions within the same vector-like representation, could expand the scope of FPVDM, offering rich prospects for both phenomenological and experimental studies, as well as insights into the complementarity of collider and non-collider observables.

5 Conclusion

In this report that has been designed following work achieved in the context of the LHC Dark Matter Working Group,

we have explored the rich phenomenology of t-channel dark matter models. These range from minimal simplified scenarios, where the dark sector comprises a single dark matter candidate and one mediator interacting with a specific SM state, to complex and non-minimal constructions inspired by flavoured dark matter, compositeness, frustrated dark matter, and gauged dark sectors. We have examined dark matter production in the early universe, considering canonical freeze-out as well as the freeze-in, superWIMP, and conversion-driven freeze-out mechanisms, and studied the resulting cosmological and astrophysical implications. At the same time, we have investigated collider signatures, distinguishing between scenarios where new particles decay promptly (except for the stable dark matter candidate) and those featuring long-lived particles on collider scales. Our results highlight the theoretical diversity of t-channel models and emphasise the intricate interplay between collider, cosmological, and astrophysical studies. In addition, they demonstrate the need for complementary efforts to constrain or validate these theoretical frameworks, particularly in the event of a discovery.

Looking ahead, this whitepaper underscores the importance of a holistic approach to dark matter research, combining theoretical model building, high-precision Monte Carlo simulations, and detailed phenomenological studies leveraging data from both colliders and cosmology. Furthermore, we advocate for continued collaboration between theorists and experimentalists, not only to maximise the potential of existing data but also to develop innovative search strategies capable of probing both minimal and non-minimal models.




975 Page 80 of 96 Eur. Phys. J. C (2025) 85:975

For this purpose, this whitepaper has been designed to provide a comprehensive and up-to-date reference that can serve as a baseline for future theoretical and experimental investigations. Given the wide range of possible signals, we deliberately refrain from proposing specific benchmark scenarios. This choice is intended to avoid biasing future searches toward a limited set of cases while potentially overlooking other relevant possibilities. Instead, we focus on providing model-independent parametrisations that help identify key features of minimal scenarios and enable efficient parametric scans. This strategy has been applied, for instance, to analyse all collider signals emerging from the considered simplified t-channel models using a single set of Monte Carlo simulations, with datasets appropriately re-weighted to explore different configurations characterised by varying couplings. Numerical tools for collider and cosmological analyses, including recast efficiencies, cross section tables and simulated samples for individual contributions in these simplified scenarios, are available upon request.

All scenarios discussed in this work, including non-minimal ones, are presented using a consistent notation wherever possible. This uniform approach aims to facilitate future analyses, enable robust reinterpretations of experimental results, and ultimately pave the way for ground-breaking discoveries. While this report does not propose specific benchmarks, we recommend that practical implementations of these guidelines in concrete experimental analyses include a harmonised approach across LHC experiments when selecting scenarios to report search results.

Acknowledgements We thank Simone Tentori for providing illustrative Feynman diagrams, Marco Drewes for highlighting current and future developments relevant for freeze-in and thermal corrections calculations, and Sam Junius for assisting in adapting the results of [252]. We are also grateful to the CERN IT division for maintaining the computing resources that have enabled us to run the simulations necessary in this report. AB acknowledges support from the STFC Consolidated Grant ST/L000296/1, the Leverhulme Trust RPG-2022-057 Grant, as well as from the Royal Society International Exchanges Grant IES/R1/211138. AL is supported by FAPESP grants no. 2018/25225-9 and 2021/01089-1. AM is supported in part by the Strategic Research Program High-Energy Physics of the Research Council of the Vrije Universiteit Brussel and by the iBOF 'Unlocking the Dark Universe with Gravitational Wave Observations: from Quantum Optics to Quantum Gravity' of the Vlaamse Interuniversitaire Raad. BF, MDG and TM have been partly supported by Grant ANR-21-CE31-0013 (project DMwith-LLPatLHC) from the French Agence Nationale de la Recherche. CA has been partially supported by the F.R.S.-FNRS under the 'Excellence of Science' EOS be.h project no. 30820817. CP and WN are supported by an SA-CERN excellence bursary. DK thanks the generous support of the Wolfson Foundation and the Royal Society to allow him to spend his sabbatical year at the University of Glasgow. DT is supported by the U.S. Department of Energy, Office of Science. EC, JuHa and MaBe were supported by the Emmy Noether grant 'Baryogenesis, Dark Matter and Neutrinos: Comprehensive analyses and accurate methods in particle cosmology' (HA 8555/1-1, Project No. 400234416) funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) and Cluster of Excellence



Data Availability Statement This manuscript has no associated data. [Authors' comment: Data sharing not applicable to this article as no datasets were generated or analysed during the current study.]

Code Availability Statement This manuscript has associated code/software in a data repository. [Authors' comment: The code/software generated during and/or analysed during the current study is available in the repository: https://github.com/BFuks/DMSimpt. Software configuration cards can be obtained upon request.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.
Funded by SCOAP³.

Appendix A: Implementation of the ATLAS-EXOT-2018-06 search in MadAnalysis5

As sketched in Sect. 3.1.1, the ATLAS-EXOT-2018-06 analysis [133] is sensitive to a signal comprising a not too large



Eur. Phys. J. C (2025) 85:975 Page 81 of 96 975

number of jets, the leading one being very energetic (with a transverse momentum larger than 150 GeV), no leptons (electrons, muons, and taus) or photons, and a significant amount of missing transverse energy well separated from the jet activity. It exploits 139 fb⁻¹ of LHC collision data at a centre-of-mass energy of 13 TeV, recorded during the period 2015-2018 by the ATLAS collaboration, and hence updates previous analyses conducted with $3.2 \, \text{fb}^{-1}$ [437] and 36.1 fb⁻¹ [438] of data using less sophisticated signal selections. It is thus relevant for probing the t-channel models that we explore in this whitepaper. In this section, we report on the validation of its implementation in the MadAnalysis 5 framework, facilitated by the substantial additional data made available via HepData [439] by the ATLAS collaboration. This includes detailed cut-flow tables and exclusion curves for given benchmark scenarios, as well as digitised information on the figures.¹⁴

1. Description of the analysis

The signal topology exploits jets and missing energy while vetoing leptons. Jets are reconstructed by clustering particles with the anti- k_T jet algorithm [142] with a radius parameter of R = 0.4. Only jets with transverse momentum $p_T > 20$ GeV and pseudo-rapidity $|\eta|$ < 2.8 are considered. Moreover, the MV2 b-tagging algorithm [440] is used to identify *b*-jets, defined as jets with $p_T > 30$ GeV and $|\eta| < 2.5$ originating from b-quark fragmentation with an average efficiency of 60%. Electron candidates must satisfy $p_T > 7 \,\text{GeV}$, $|\eta|$ < 2.47, and 'Loose' track selection criteria [441], which requires their longitudinal impact parameter to be less than 0.5 mm. Overlaps between identified electrons and jets with $p_T > 30 \text{ GeV}$ are resolved by discarding non-b-tagged jets within $\Delta R < 0.2$ of any identified electron, and by removing electrons within $\Delta R < 0.4$ of any remaining jets. Muon candidates must pass a 'Medium' identification selection [442], and have $p_T > 7$ GeV, $|\eta| < 2.5$, and a longitudinal impact parameter smaller than 0.5 mm. Jets with $p_T > 30 \text{ GeV}$ and fewer than three associated tracks with $p_T > 500 \text{ MeV}$ are then discarded if they are within $\Delta R < 0.4$ of an identified muon. Hadronically-decaying tau leptons are reconstructed from jets with $p_T > 10$ GeV and $|\eta| < 2.5$, and they must satisfy 'Loose' identification requirements, have a transverse momentum larger than 20 GeV after energyscale corrections, and be associated with either one or three charged tracks [443,444]. Tau candidates within $\Delta R < 0.2$ of an electron or muon are removed, as are any jets within $\Delta R = 0.2$ of a reconstructed tau-lepton. Finally, the missing transverse momentum $\mathbf{p}_T^{\text{miss}}$ is reconstructed from the negative vector sum of the transverse momenta of all reconstructed objects with $p_T > 20$ GeV and $|\eta| < 4.5$.

Event preselection requires a significant amount of missing energy, $E_T^{\rm miss} > 200$ GeV, and an energetic leading jet with $p_T > 150$ GeV and $|\eta| < 2.4$. Moreover, up to three additional jets with $p_T > 30$ GeV and $|\eta| < 2.8$ are allowed. Additionally, the missing transverse momentum $\mathbf{p}_T^{\rm miss}$ must be well separated from the four leading jets by an angle of $\Delta\phi(j,\mathbf{p}_T^{\rm miss}) > 0.4$ for events with $E_T^{\rm miss} > 250$ GeV and $\Delta\phi(j,\mathbf{p}_T^{\rm miss}) > 0.6$ for events with $E_T^{\rm miss} \leq 250$ GeV, such a separation criterion helping to reduce the multijet background contributions.

Next, the ATLAS collaboration implements a twofold analysis strategy, including signal regions (SRs) with either an inclusive or an exclusive selection on the missing transverse energy. Inclusive E_T^{miss} signal regions (denoted by names starting with IM) are used for model-independent interpretations of the search results, while exclusive signal regions (denoted by names starting with EM) are used for model-dependent interpretations. In the first series of 13 signal regions (EMO, EM1, ..., EM12), the analysis considers an exclusive missing transverse energy selection defined by $E_{\min} < E_T^{\text{miss}} < E_{\max}$. The different thresholds range from 200 GeV to 1200 GeV, as shown in Table 9 which also includes all preselection cuts and an extra cut on the missing transverse momentum ($E_T^{\text{miss}} > 150 \text{ GeV}$) allowing us to be consistent with generator-level cuts biasing event generation to the phase space region of interest (as implemented in the ATLAS simulations). In the second set of 13 signal regions (IMO, IM1, ..., IM12), the analysis instead considers an inclusive missing transverse energy selection defined by $E_T^{\text{miss}} > E_{\text{threshold}}$, with the different thresholds again ranging from 200 GeV to 1200 GeV, as given in Table 9.

2. Validation of the implementation

The validation of our implementation of the ATLAS-EXOT-2018-06 search in MadAnalysis 5 has been achieved by focusing on top squark production and decay in the *R*-parity-conserving Minimal Supersymmetric Standard Model (MSSM). We have considered two different final states, corresponding to the processes

$$pp \to \tilde{t}_1 \tilde{t}_1^* \to (c \tilde{\chi}_1^0) (\bar{c} \tilde{\chi}_1^0) \quad \text{and}$$

$$pp \to \tilde{t}_1 \tilde{t}_1^* \to (bf \, \bar{f}' \tilde{\chi}_1^0) (\bar{b} f \, \bar{f}' \tilde{\chi}_1^0), \tag{A1}$$

and we have computed limits and cut-flows for different choices of the stop and neutralino masses, with all other superpartners being decoupled.

Hard-scattering signal event generation has been achieved with MadGraph5_aMC@NLO [50] (version 3.4.2), while the simulation of supersymmetric particle decays, parton showering, and hadronisation has been performed with Pythia [445] (version 8.2). This event generation procedure relies on the MSSM implementation in MadGraph5_aMC@NLO



¹⁴ See the webpage https://www.hepdata.net/record/ins1847779.

975 Page 82 of 96 Eur. Phys. J. C (2025) 85:975

Table 9 Selection cuts defining the different signal regions of the ATLAS-EXOT-2018-06 analysis [133]

Cuts	Exclusive SRs	Inclusive SRs
Preselection	$E_T^{\rm miss} > 150~{\rm GeV}$	
	Lepton veto	
	$N(j) \in [1, 4]$	
	$\Delta\Phi(j_i, \mathbf{p}_T^{\mathrm{miss}}) > 0.4 \ (\forall i)$	
	$p_T(j_1) > 150 \mathrm{GeV}$	
	$E_T^{ m miss} > 200~{ m GeV}$	
Bin 0	EM0: 200 GeV $< E_T^{ m miss} <$ 250 GeV	IM0: $E_T^{\rm miss} > 200~{ m GeV}$
Bin 1	EM1: 250 GeV $< E_T^{\text{miss}} < 300 \text{ GeV}$	IM1: $E_T^{\text{miss}} > 250 \text{ GeV}$
Bin 2	EM2: 300 GeV $< E_T^{\text{miss}} < 350 \text{ GeV}$	IM2: $E_T^{\text{miss}} > 300 \text{ GeV}$
Bin 3	EM3: 350 GeV $< E_T^{\text{miss}} < 400 \text{ GeV}$	IM3: $E_T^{\text{miss}} > 350 \text{ GeV}$
Bin 4	EM4: $400 \text{ GeV} < E_T^{\text{miss}} < 500 \text{ GeV}$	IM4: $E_T^{ m miss} > 400~{ m GeV}$
Bin 5	EM5: $500 \text{ GeV} < E_T^{\text{miss}} < 600 \text{ GeV}$	IM5: $E_T^{\rm miss} > 500~{\rm GeV}$
Bin 6	EM6: $600 \text{ GeV} < E_T^{\text{miss}} < 700 \text{ GeV}$	IM6: $E_T^{\rm miss} > 600~{ m GeV}$
Bin 7	EM7: 700 GeV $< E_T^{\text{miss}} < 800 \text{ GeV}$	IM7: $E_T^{\text{miss}} > 700 \text{ GeV}$
Bin 8	EM8: $800 \text{ GeV} < E_T^{\text{miss}} < 900 \text{ GeV}$	IM8: $E_T^{ m miss} > 800~{ m GeV}$
Bin 9	EM9: 900 GeV $< E_T^{\text{miss}} < 1000 \text{ GeV}$	IM9: $E_T^{\text{miss}} > 900 \text{ GeV}$
Bin 10	EM10: $1000 \text{ GeV} < E_T^{\text{miss}} < 1100 \text{ GeV}$	$IM10: E_T^{\text{miss}} > 1000 \text{ GeV}$
Bin 11	EM11: $1100 \text{ GeV} < E_T^{\text{miss}} < 1200 \text{ GeV}$	IM11: $E_T^{\text{miss}} > 1100 \text{ GeV}$
Bin 12	EM12: $E_T^{\text{miss}} > 1200 \text{ GeV}$	IM12: $E_T^{\text{miss}} > 1200 \text{ GeV}$

described in [446], which makes use of FeynRules [41,42] and its UFO interface [46,47]. Moreover, we have merged event samples including up to two additional hard partons in the final state, following the MLM prescription as implemented in MadGraph5_aMC@NLO [183,184] with a merging scale set to one quarter of the stop mass. To match the statistics of the reference cut-flows provided by the ATLAS collaboration on HepData, we simulated 100,000 events before merging, which resulted in samples of about 90,000 merged events.

The information available on HepData provides cut-flow information for several benchmark scenarios. We begin our validation by focusing on the second process of Eq. (A1), considering a scenario with a compressed spectrum featuring a stop mass of $m_{\tilde{t}_1}=450$ GeV and a neutralino mass of $m_{\tilde{\chi}_1^0}=443$ GeV. We have generated events as introduced above, and then analysed the produced sample by applying all the analysis preselection cuts and assessing how many signal events ($N_{\rm events}$) would populate the different exclusive bins in missing transverse energy. Our results are displayed, for each individual cut, in Table 10, that reports the number of events surviving each cut and the associated cumulative efficiency ε . We include both the predictions provided by ATLAS ($N_{\rm events}^{\rm ATLAS}$) and those predicted using our implementation in MadAnalysis 5 ($N_{\rm events}^{\rm MA5}$ and $\varepsilon_{\rm MA5}$),

along with the relative difference R between the two,

$$R = \left| \frac{\varepsilon_{\text{ATLAS}} - \varepsilon_{\text{MA5}}}{\varepsilon_{\text{ATLAS}}} \right|. \tag{A2}$$

We observe a good agreement between our predictions and the ATLAS ones, with *R* values ranging from 1% to 25%. The largest discrepancies are associated with signal regions in which the number of Monte Carlo events populating the bin is small, especially for the ATLAS predictions, indicating that a large numerical uncertainty must be accounted in the comparison.

To further validate our implementation, we also determine exclusion contours at the 95% confidence level (CL) for the two simplified models corresponding to the two processes of Eq. (A1). Our results are presented in the stop mass versus neutralino mass plane in Fig. 55 for the $pp \rightarrow \tilde{t}_1 \tilde{t}_1^* \rightarrow (c \tilde{\chi}_1^0) (\bar{c} \tilde{\chi}_1^0)$ process (left panel) and $pp \rightarrow \tilde{t}_1 \tilde{t}_1^* \rightarrow (bf \, \bar{f}' \tilde{\chi}_1^0) (\bar{b} \, f \, \bar{f}' \tilde{\chi}_1^0)$ process (right panel). We explore scenarios with stop and neutralino masses varying between 300 and 600 GeV, and superimpose the exclusion contours obtained with MadAnalysis 5 (green) with the official ones provided by the ATLAS Collaboration (black). An excellent degree of agreement is observed, with the excluded mass configurations agreeing at the level of a few percent. This therefore validates our implementation.



Eur. Phys. J. C (2025) 85:975 Page 83 of 96 975

Table 10 Cut-flow associated with the ATLAS-EXOT-2018-06 analysis, for the signal emerging from the process $pp \to \tilde{\iota}_1 \tilde{\iota}_1^* \to (bf \, \bar{f}' \tilde{\chi}_1^0) (\bar{b} \, f \, \bar{f}' \tilde{\chi}_1^0)$ and a spectrum with $(m_{\tilde{\iota}_1}, m_{\tilde{\chi}_1^0}) =$

(450, 443) GeV. We compare our predictions for the the number of generated events surviving each cut and with the associated efficiencies with the information provided by ATLAS on HepData

Cut	ATLAS		MadAnalysis 5		R[%]
	N _{events} ATLAS	$arepsilon_{ ext{ATLAS}}[\%]$	N _{events} ^{MA5}	$arepsilon_{ ext{MA5}}[\%]$	
$E_T^{\text{miss}} > 150 \text{ GeV}$	39598	100	89529	100	_
Lepton veto	37547	94.82	85417	95.41	0.62
$N(j) \in [1, 4]$	35412	89.43	76195	85.11	4.38
$\Delta\Phi(j_i, \mathbf{p}_T^{\text{miss}}) > 0.4 \ (\forall i)$	33319	84.14	69253	77.35	8.07
$p_T(j_1) > 150 \mathrm{GeV}$	23134	58.42	47157	52.67	9.84
$E_T^{\rm miss} > 200~{ m GeV}$	18801	47.48	39183	43.77	7.81
EM0	4488	11.34	8509	9.50	16.23
EM1	3789	9.57	7946	8.88	7.21
EM2	2857	7.21	6226	6.95	3.61
EM3	2111	5.33	4621	5.16	3.19
EM4	2618	6.61	5847	6.53	1.21
EM5	1352	3.41	2895	3.23	5.28
EM6	712	1.80	1501	1.67	7.22
EM7	393	0.99	719	0.80	19.19
EM8	204	0.52	408	0.46	11.54
EM9	122	0.31	207	0.23	25.80
EM10	58	0.15	124	0.14	6.67
EM11	42	0.11	77	0.09	18.18
EM12	55	0.14	103	0.11	21.43





Fig. 55 Excluded region at the 95% CL, displayed in the $(m_{\tilde{t}_1}, m_{\tilde{\chi}_1^0})$ mass plane for the $pp \to \tilde{t}_1 \tilde{t}_1^* \to (c \tilde{\chi}_1^0) (\bar{c} \tilde{\chi}_1^0)$ (left) and $pp \to \tilde{t}_1 \tilde{t}_1^* \to (bf \bar{f}' \tilde{\chi}_1^0) (\bar{b} f \bar{f}' \tilde{\chi}_1^0)$ (right) processes. The green crosses and red dots respectively correspond to scenarios allowed and excluded



when using our implementation of the ATLAS-EXOT-2018-06 analysis in MadAnalysis 5. The excluded parameter space regions are thus delineated by the green contours, that could be compared to the official ATLAS exclusions (black)



975 Page 84 of 96 Eur. Phys. J. C (2025) 85:975

References

- G. Bertone, D. Hooper, J. Silk, Particle dark matter: evidence, candidates and constraints. Phys. Rep. Ser. 405, 279 (2005). https://doi.org/10.1016/j.physrep.2004.08.031. arXiv:hep-ph/0404175
- J. Silk et al., Particle Dark Matter: Observations, Models and Searches (Cambridge Univ. Press, Cambridge, 2010). https://doi. org/10.1017/CBO9780511770739
- G. Bertone, D. Hooper, History of dark matter. Rev. Mod. Phys. Ser. 90, 045002 (2018). https://doi.org/10.1103/RevModPhys.90. 045002. arXiv:1605.04909
- 4. M. Cirelli, A. Strumia, J. Zupan, Dark matter. arXiv:2406.01705
- J. Alwall, P. Schuster, N. Toro, Simplified models for a first characterization of new physics at the LHC. Phys. Rev. D Ser. 79, 075020 (2009). https://doi.org/10.1103/PhysRevD.79. 075020. arXiv:0810.3921
- shape LHC New Physics Working Group collaboration, Simplified models for LHC new physics searches. J. Phys. G Ser. 39, 105005 (2012). https://doi.org/10.1088/0954-3899/39/10/105005. arXiv:1105.2838
- P.J. Fox, C. Williams, Next-to-leading order predictions for dark matter production at hadron colliders. Phys. Rev. D Ser. 87, 054030 (2013). https://doi.org/10.1103/PhysRevD.87. 054030. arXiv:1211.6390
- U. Haisch, F. Kahlhoefer, E. Re, QCD effects in mono-jet searches for dark matter. JHEP Ser. 12, 007 (2013). https://doi.org/10. 1007/JHEP12(2013)007. arXiv:1310.4491
- M. Backović, M. Krämer, F. Maltoni, A. Martini, K. Mawatari, M. Pellen, Higher-order QCD predictions for dark matter production at the LHC in simplified models with s-channel mediators. Eur. Phys. J. C Ser. 75, 482 (2015). https://doi.org/10.1140/epjc/s10052-015-3700-6. arXiv:1508.05327
- D. Abercrombie et al., Dark matter benchmark models for early LHC Run-2 searches: report of the ATLAS/CMS Dark Matter Forum. Phys. Dark Univ. Ser. 27, 100371 (2020). https://doi.org/ 10.1016/j.dark.2019.100371. arXiv:1507.00966
- A. Boveia et al., Recommendations on presenting LHC searches for missing transverse energy signals using simplified s-channel models of dark matter. Phys. Dark Univ. Ser. 27, 100365 (2020). https://doi.org/10.1016/j.dark.2019.100365. arXiv:1603.04156
- A. Albert et al., Recommendations of the LHC Dark Matter Working Group: comparing LHC searches for dark matter mediators in visible and invisible decay channels and calculations of the thermal relic density. Phys. Dark Univ. Ser. 26, 100377 (2019). https://doi.org/10.1016/j.dark.2019.100377. arXiv:1703.05703
- A. Albert et al., Displaying dark matter constraints from colliders with varying simplified model parameters. arXiv:2203.12035
- M. Garny, A. Ibarra, S. Vogl, Dark matter annihilations into two light fermions and one gauge boson: general analysis and antiproton constraints. JCAP Ser. 04, 033 (2012). https://doi.org/10. 1088/1475-7516/2012/04/033. arXiv:1112.5155
- M. Garny, A. Ibarra, M. Pato, S. Vogl, Internal bremsstrahlung signatures in light of direct dark matter searches. JCAP Ser. 12, 046 (2013). https://doi.org/10.1088/1475-7516/2013/12/046. arXiv:1306.6342
- Y. Bai, J. Berger, Fermion portal dark matter. JHEP Ser. 11, 171 (2013). https://doi.org/10.1007/JHEP11(2013)171. arXiv:1308.0612
- A. DiFranzo, K.I. Nagao, A. Rajaraman, T.M.P. Tait, Simplified models for dark matter interacting with quarks. JHEP Ser. 11, 014 (2013). https://doi.org/10.1007/JHEP11(2013)014. arXiv:1308.2679
- H. An, L.-T. Wang, H. Zhang, Dark matter with t-channel mediator: a simple step beyond contact interaction. Phys. Rev. D

- Ser. **89**, 115014 (2014). https://doi.org/10.1103/PhysRevD.89. 115014. arXiv:1308.0592
- M. Papucci, A. Vichi, K.M. Zurek, Monojet versus the rest of the world I: t-channel models. JHEP Ser. 11, 024 (2014). https://doi. org/10.1007/JHEP11(2014)024. arXiv:1402.2285
- S. Chang, R. Edezhath, J. Hutchinson, M. Luty, Effective WIMPs. Phys. Rev. D Ser. 89, 015011 (2014). https://doi.org/10.1103/ PhysRevD.89.015011. arXiv:1307.8120
- F. Giacchino, L. Lopez-Honorez, M.H.G. Tytgat, Bremsstrahlung and gamma ray lines in 3 scenarios of dark matter annihilation. JCAP Ser. 08, 046 (2014). https://doi.org/10.1088/1475-7516/ 2014/08/046. arXiv:1405.6921
- J. Hisano, R. Nagai, N. Nagata, Effective theories for dark matter nucleon scattering. JHEP Ser. 05, 037 (2015). https://doi.org/10. 1007/JHEP05(2015)037. arXiv:1502.02244
- S. El Hedri, A. Kaminska, M. de Vries, J. Zurita, Simplified phenomenology for colored dark sectors. JHEP Ser. 04, 118 (2017). https://doi.org/10.1007/JHEP04(2017)118. arXiv:1703.00452
- C. Arina, B. Fuks, L. Mantani, H. Mies, L. Panizzi, J. Salko, Closing in on *t*-channel simplified dark matter models. Phys. Lett. B Ser. 813, 136038 (2021). https://doi.org/10.1016/j.physletb.2020. 136038. arXiv:2010.07559
- C. Arina, B. Fuks, J. Heisig, M. Krämer, L. Mantani, L. Panizzi, Comprehensive exploration of t-channel simplified models of dark matter. Phys. Rev. D Ser. 108, 115007 (2023). https://doi.org/10. 1103/PhysRevD.108.115007. arXiv:2307.10367
- P. Agrawal, M. Blanke, K. Gemmler, Flavored dark matter beyond minimal flavor violation. JHEP Ser. 10, 072 (2014). https://doi. org/10.1007/JHEP10(2014)072. arXiv:1405.6709
- S. Colucci, B. Fuks, F. Giacchino, L. Lopez Honorez, M.H.G. Tytgat, J. Vandecasteele, Top-philic vector-like portal to scalar dark matter. Phys. Rev. D 98, 035002 (2018). https://doi.org/10. 1103/PhysRevD.98.035002. arXiv:1804.05068
- A.S. Cornell, A. Deandrea, T. Flacke, B. Fuks, L. Mason, Contact interactions and top-philic scalar dark matter. JHEP Ser. 07, 026 (2021). https://doi.org/10.1007/JHEP07(2021)026. arXiv:2104.12795
- A.S. Cornell, A. Deandrea, T. Flacke, B. Fuks, L. Mason, Top partners and scalar dark matter: a nonminimal reappraisal. Phys. Rev. D Ser. 107, 075004 (2023). https://doi.org/10.1103/PhysRevD. 107.075004. arXiv:2209.13093
- L.M. Carpenter, T. Murphy, T.M.P. Tait, Distinctive signals of frustrated dark matter. JHEP Ser. 09, 175 (2022). https://doi.org/ 10.1007/JHEP09(2022)175. arXiv:2205.06824
- M. Cirelli, N. Fornengo, A. Strumia, Minimal dark matter. Nucl. Phys. B Ser. 753, 178 (2006). https://doi.org/10.1016/j.nuclphysb. 2006.07.012. arXiv:hep-ph/0512090
- A. Belyaev, G. Cacciapaglia, D. Locke, A. Pukhov, Minimal consistent dark matter models for systematic experimental characterisation: fermion dark matter. JHEP Ser. 10, 014 (2022). https://doi.org/10.1007/JHEP10(2022)014. arXiv:2203.03660
- A. Belyaev, A. Deandrea, S. Moretti, L. Panizzi, D.A. Ross, N. Thongyoi, Fermionic portal to vector dark matter from a new gauge sector. Phys. Rev. D Ser. 108, 095001 (2023). https://doi.org/10.1103/PhysRevD.108.095001. arXiv:2204.03510
- M. Garny, J. Heisig, B. Lülf, S. Vogl, Coannihilation without chemical equilibrium. Phys. Rev. D Ser. 96, 103521 (2017). https://doi.org/10.1103/PhysRevD.96.103521. arXiv:1705.09292
- R.T. D'Agnolo, D. Pappadopulo, J.T. Ruderman, Fourth exception in the calculation of relic abundances. Phys. Rev. Lett. Ser. 119, 061102 (2017). https://doi.org/10.1103/PhysRevLett.119. 061102. arXiv:1705.08450
- J. McDonald, Thermally generated gauge singlet scalars as selfinteracting dark matter. Phys. Rev. Lett. Ser. 88,



Eur. Phys. J. C (2025) 85:975 Page 85 of 96 975

091304 (2002). https://doi.org/10.1103/PhysRevLett.88.091304. arXiv:hep-ph/0106249

- L.J. Hall, K. Jedamzik, J. March-Russell, S.M. West, Freeze-in production of FIMP dark matter. JHEP Ser. 03, 080 (2010). https:// doi.org/10.1007/JHEP03(2010)080. arXiv:0911.1120
- L. Covi, J.E. Kim, L. Roszkowski, Axinos as cold dark matter. Phys. Rev. Lett. Ser. 82, 4180 (1999). https://doi.org/10.1103/ PhysRevLett.82.4180. arXiv:hep-ph/9905212
- J.L. Feng, A. Rajaraman, F. Takayama, SuperWIMP dark matter signals from the early universe. Phys. Rev. D Ser. 68, 063504 (2003). https://doi.org/10.1103/PhysRevD.68.063504. arXiv:hep-ph/0306024
- C. Arina, B. Fuks, L. Mantani, A universal framework for t-channel dark matter models. Eur. Phys. J. C Ser. 80, 409 (2020). https://doi.org/10.1140/epjc/s10052-020-7933-7. arXiv:2001.05024
- 41. N.D. Christensen et al., A comprehensive approach to new physics simulations. Eur. Phys. J. Ser. C71, 1541 (2011). https://doi.org/10.1140/epjc/s10052-011-1541-5. arXiv:0906.2474
- A. Alloul, N.D. Christensen, C. Degrande, C. Duhr, B. Fuks, Feyn-Rules 2.0 a complete toolbox for tree-level phenomenology. Comput. Phys. Commun. 185, 2250 (2014). https://doi.org/10.1016/j.cpc.2014.04.012. arXiv:1310.1921
- S. Frixione, B. Fuks, V. Hirschi, K. Mawatari, H.-S. Shao, P.A. Sunder et al., Automated simulations beyond the Standard Model: supersymmetry. JHEP Ser. 12, 008 (2019). https://doi.org/10.1007/JHEP12(2019)008. arXiv:1907.04898
- C. Degrande, Automatic evaluation of UV and R2 terms for beyond the Standard Model Lagrangians: a proof-of-principle. Comput. Phys. Commun. Ser. 197, 239 (2015). https://doi.org/ 10.1016/j.cpc.2015.08.015. arXiv:1406.3030
- T. Hahn, Generating Feynman diagrams and amplitudes with FeynArts 3. Comput. Phys. Commun. Ser. 140, 418 (2001). https://doi.org/10.1016/S0010-4655(01)00290-9. arXiv:hep-ph/0012260
- C. Degrande, C. Duhr, B. Fuks, D. Grellscheid, O. Mattelaer, T. Reiter, UFO the universal FeynRules output. Comput. Phys. Commun. Ser. 183, 1201 (2012). https://doi.org/10.1016/j.cpc. 2012.01.022. arXiv:1108.2040
- L. Darmé et al., UFO 2.0: the 'Universal Feynman Output' format. Eur. Phys. J. C 83, 631 (2023). https://doi.org/10.1140/epjc/s10052-023-11780-9. arXiv:2304.09883
- PARTICLE DATA GROUP Collaboration, Review of particle physics. PTEP 2022, 083C01 (2022). https://doi.org/10.1093/ ptep/ptac097
- P.Z. Skands et al., SUSY Les Houches accord: interfacing SUSY spectrum calculators, decay packages, and event generators. JHEP Ser. 07, 036 (2004). https://doi.org/10.1088/1126-6708/2004/07/036. arXiv:hep-ph/0311123
- J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer et al., The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations. JHEP Ser. 07, 079 (2014). https://doi.org/10.1007/JHEP07(2014)079. arXiv:1405.0301
- A. Belyaev, N.D. Christensen, A. Pukhov, CalcHEP 3.4 for collider physics within and beyond the Standard Model. Comput. Phys. Commun. 184, 1729 (2013). https://doi.org/10.1016/j.cpc. 2013.01.014. arXiv:1207.6082
- G. Bélanger, F. Boudjema, A. Goudelis, A. Pukhov, B. Zaldivar, micrOMEGAs5.0: freeze-in. Comput. Phys. Commun. 231, 173 (2018). https://doi.org/10.1016/j.cpc.2018.04.027. arXiv:1801.03509
- G. Alguero, G. Belanger, F. Boudjema, S. Chakraborti, A. Goudelis, S. Kraml et al., micrOMEGAs 6.0: N-component dark matter. Comput. Phys. Commun. 299, 109133 (2024). https://doi.org/10.1016/j.cpc.2024.109133. arXiv:2312.14894

- 54. F. Ambrogi, C. Arina, M. Backovic, J. Heisig, F. Maltoni, L. Mantani et al., MadDM v.3.0: a comprehensive tool for dark matter studies. Phys. Dark Univ. 24, 100249 (2019). https://doi.org/10.1016/j.dark.2018.11.009. arXiv:1804.00044
- C. Arina, J. Heisig, F. Maltoni, L. Mantani, D. Massaro, O. Mattelaer et al., Studying dark matter with MadDM 3.1: a short user guide. PoS TOOLS2020, 009 (2021). https://doi.org/10.22323/1.392.0009. arXiv:2012.09016
- C. Arina, J. Heisig, F. Maltoni, D. Massaro, O. Mattelaer, Indirect dark-matter detection with MadDM v3.2 lines and loops.
 Eur. Phys. J. C 83, 241 (2023). https://doi.org/10.1140/epjc/s10052-023-11377-2. arXiv:2107.04598
- M.J. Baker, A. Thamm, Leptonic WIMP coannihilation and the current dark matter search strategy. JHEP Ser. 10, 187 (2018). https://doi.org/10.1007/JHEP10(2018)187. arXiv:1806.07896
- J. Kile, A. Soni, Flavored dark matter in direct detection experiments and at LHC. Phys. Rev. D Ser. 84, 035016 (2011). https://doi.org/10.1103/PhysRevD.84.035016. arXiv:1104.5239
- J.F. Kamenik, J. Zupan, Discovering dark matter through flavor violation at the LHC. Phys. Rev. D Ser. 84, 111502 (2011). https:// doi.org/10.1103/PhysRevD.84.111502. arXiv:1107.0623
- B. Batell, J. Pradler, M. Spannowsky, Dark matter from minimal flavor violation. JHEP Ser. 08, 038 (2011). https://doi.org/10.1007/JHEP08(2011)038. arXiv:1105.1781
- P. Agrawal, S. Blanchet, Z. Chacko, C. Kilic, Flavored dark matter, and its implications for direct detection and colliders. Phys. Rev. D Ser. 86, 055002 (2012). https://doi.org/10.1103/PhysRevD.86. 055002. arXiv:1109.3516
- B. Batell, T. Lin, L.-T. Wang, Flavored dark matter and R-parity violation. JHEP Ser. 01, 075 (2014). https://doi.org/10.1007/ JHEP01(2014)075. arXiv:1309.4462
- J. Kile, Flavored dark matter: a review. Mod. Phys. Lett. A Ser. 28, 1330031 (2013). https://doi.org/10.1142/S0217732313300310. arXiv:1308.0584
- J. Kile, A. Kobach, A. Soni, Lepton-flavored dark matter. Phys. Lett. B Ser. 744, 330 (2015). https://doi.org/10.1016/j.physletb. 2015.04.005, arXiv:1411.1407
- L. Lopez-Honorez, L. Merlo, Dark matter within the minimal flavour violation ansatz. Phys. Lett. B Ser. 722, 135 (2013). https:// doi.org/10.1016/j.physletb.2013.04.015. arXiv:1303.1087
- R.S. Chivukula, H. Georgi, Composite technicolor Standard Model. Phys. Lett. B Ser. 188, 99 (1987). https://doi.org/10.1016/ 0370-2693(87)90713-1
- L.J. Hall, L. Randall, Weak scale effective supersymmetry. Phys. Rev. Lett. Ser. 65, 2939 (1990). https://doi.org/10.1103/ PhysRevLett.65.2939
- A.J. Buras, P. Gambino, M. Gorbahn, S. Jager, L. Silvestrini, Universal unitarity triangle and physics beyond the standard model. Phys. Lett. B Ser. 500, 161 (2001). https://doi.org/10. 1016/S0370-2693(01)00061-2. arXiv:hep-ph/0007085
- G. D'Ambrosio, G.F. Giudice, G. Isidori, A. Strumia, Minimal flavor violation: an effective field theory approach. Nucl. Phys. B Ser. 645, 155 (2002). https://doi.org/10.1016/S0550-3213(02)00836-2. arXiv:hep-ph/0207036
- A.J. Buras, Minimal flavor violation. Acta Phys. Polon. B Ser. 34, 5615 (2003). arXiv:hep-ph/0310208
- V. Cirigliano, B. Grinstein, G. Isidori, M.B. Wise, Minimal flavor violation in the lepton sector. Nucl. Phys. B Ser. 728, 121 (2005). https://doi.org/10.1016/j.nuclphysb.2005.08. 037. arXiv:hep-ph/0507001
- P. Agrawal, Z. Chacko, E.C.F.S. Fortes, C. Kilic, Skew-flavored dark matter. Phys. Rev. D Ser. 93, 103510 (2016). https://doi.org/ 10.1103/PhysRevD.93.103510. arXiv:1511.06293
- T. Jubb, M. Kirk, A. Lenz, Charming dark matter. JHEP Ser. 12, 010 (2017). https://doi.org/10.1007/JHEP12(2017)010. arXiv:1709.01930



975 Page 86 of 96 Eur. Phys. J. C (2025) 85:975

 M. Blanke, S. Kast, Top-flavoured dark matter in dark minimal flavour violation. JHEP Ser. 05, 162 (2017). https://doi.org/10. 1007/JHEP05(2017)162. arXiv:1702.08457

- M. Blanke, S. Das, S. Kast, Flavoured dark matter moving left. JHEP Ser. 02, 105 (2018). https://doi.org/10.1007/ JHEP02(2018)105. arXiv:1711.10493
- H. Acaroglu, M. Blanke, Tasting flavoured Majorana dark matter. JHEP Ser. 05, 086 (2022). https://doi.org/10.1007/ JHEP05(2022)086. arXiv:2109.10357
- H. Acaroğlu, M. Blanke, J. Heisig, M. Krämer, L. Rathmann, Flavoured Majorana dark matter then and now: from freeze-out scenarios to LHC signatures. JHEP Ser. 06, 179 (2024). https:// doi.org/10.1007/JHEP06(2024)179. arXiv:2312.09274
- M.-C. Chen, J. Huang, V. Takhistov, Beyond minimal lepton flavored dark matter. JHEP Ser. 02, 060 (2016). https://doi.org/10.1007/JHEP02(2016)060. arXiv:1510.04694
- H. Acaroğlu, P. Agrawal, M. Blanke, Lepton-flavoured scalar dark matter in dark minimal flavour violation. JHEP Ser. 05, 106 (2023). https://doi.org/10.1007/JHEP05(2023)106. arXiv:2211.03809
- H. Acaroğlu, M. Blanke, M. Tabet, Opening the Higgs portal to lepton-flavoured dark matter. JHEP Ser. 11, 079 (2023). https:// doi.org/10.1007/JHEP11(2023)079. arXiv:2309.10700
- 81. H. Acaroğlu, P. Agrawal, M. Blanke, Flavoured $(g-2)_{\mu}$ with dark lepton seasoning. SciPost Phys. **15**, 176 (2023). https://doi.org/10.21468/SciPostPhys.15.4.176. arXiv:2212.08142
- 82. J. Heisig, Conversion-driven leptogenesis: a testable theory of dark matter and baryogenesis at the electroweak scale. Phys. Rev. Lett. Ser. 133, 191803 (2024). https://doi.org/10.1103/ PhysRevLett.133.191803. arXiv:2404.12428
- 83. G. Panico, A. Wulzer, *The Composite Nambu–Goldstone Higgs*, vol. 913 (Springer, 2016). https://doi.org/10.1007/978-3-319-22617-0. arXiv:1506.01961
- G. Cacciapaglia, C. Pica, F. Sannino, Fundamental composite dynamics: a review. Phys. Rep. Ser. 877, 1 (2020). https://doi. org/10.1016/j.physrep.2020.07.002. arXiv:2002.04914
- 85. G. Cacciapaglia, A. Deandrea, K. Sridhar, Review of fundamental composite dynamics. Eur. Phys. J. ST Ser. **231**, 1221 (2022). https://doi.org/10.1140/epjs/s11734-022-00549-y
- D.B. Kaplan, Flavor at SSC energies: a new mechanism for dynamically generated fermion masses. Nucl. Phys. B Ser. 365, 259 (1991). https://doi.org/10.1016/S0550-3213(05)80021-5
- A. Belyaev, R.S. Chivukula, B. Fuks, E.H. Simmons, X. Wang, Vectorlike top quark production via a chromomagnetic moment at the LHC. Phys. Rev. D Ser. 104, 095024 (2021). https://doi. org/10.1103/PhysRevD.104.095024. arXiv:2107.12402
- L.M. Carpenter, T. Murphy, T.M.P. Tait, Phenomenological cornucopia of SU(3) exotica. Phys. Rev. D Ser. 105, 035014 (2022). https://doi.org/10.1103/PhysRevD.105.035014. arXiv:2110.11359
- J. Shu, T.M.P. Tait, K. Wang, Explorations of the top quark forward-backward asymmetry at the tevatron. Phys. Rev. D Ser. 81, 034012 (2010). https://doi.org/10.1103/PhysRevD.81. 034012. arXiv:0911.3237
- T. Han, I. Lewis, T. McElmurry, QCD corrections to scalar diquark production at hadron colliders. JHEP Ser. 01, 123 (2010). https://doi.org/10.1007/JHEP01(2010)123. arXiv:0909.2666
- T. Han, I. Lewis, Z. Liu, Colored resonant signals at the LHC: largest rate and simplest topology. JHEP Ser. 12, 085 (2010). https://doi.org/10.1007/JHEP12(2010)085. arXiv:1010.4309
- K.S. Babu, P.S. Bhupal Dev, R.N. Mohapatra, Neutrino mass hierarchy, neutron–anti-neutron oscillation from baryogenesis. Phys. Rev. D 79, 015017 (2009). https://doi.org/10.1103/PhysRevD.79.015017. arXiv:0811.3411
- K.S. Babu, P.S. Bhupal Dev, E.C.F.S. Fortes, R.N. Mohapatra, Post-Sphaleron baryogenesis and an upper limit on the neutron—

- antineutron oscillation time. Phys. Rev. D **87**, 115019 (2013). https://doi.org/10.1103/PhysRevD.87.115019. arXiv:1303.6918
- J. Hubisz, P. Meade, Phenomenology of the littlest Higgs with T-parity. Phys. Rev. D Ser. 71, 035016 (2005). https://doi.org/10. 1103/PhysRevD.71.035016. arXiv:hep-ph/0411264
- T. Hambye, Hidden vector dark matter. JHEP Ser. 01, 028 (2009). https://doi.org/10.1088/1126-6708/2009/01/028. arXiv:0811.0172
- F. Chen, J.M. Cline, A.R. Frey, Nonabelian dark matter: models and constraints. Phys. Rev. D Ser. 80, 083516 (2009). https://doi. org/10.1103/PhysRevD.80.083516. arXiv:0907.4746
- J.L. Diaz-Cruz, E. Ma, Neutral SU(2) gauge extension of the standard model and a vector-boson dark-matter candidate. Phys. Lett. B Ser. 695, 264 (2011). https://doi.org/10.1016/j.physletb.2010. 11.039. arXiv:1007.2631
- S. Bhattacharya, J.L. Diaz-Cruz, E. Ma, D. Wegman, Dark vector-gauge-boson model. Phys. Rev. D Ser. 85, 055008 (2012). https://doi.org/10.1103/PhysRevD.85.055008. arXiv:1107.2093
- O. Lebedev, H.M. Lee, Y. Mambrini, Vector Higgs-portal dark matter and the invisible Higgs. Phys. Lett. B Ser. 707, 570 (2012). https://doi.org/10.1016/j.physletb.2012.01.029. arXiv:1111.4482
- Y. Farzan, A.R. Akbarieh, VDM: a model for vector dark matter. JCAP Ser. 10, 026 (2012). https://doi.org/10.1088/1475-7516/ 2012/10/026. arXiv:1207.4272
- S. Baek, P. Ko, W.-I. Park, E. Senaha, Higgs portal vector dark matter: revisited. JHEP Ser. 05, 036 (2013). https://doi.org/10. 1007/JHEP05(2013)036. arXiv:1212.2131
- 102. E. Koorambas, Vector gauge boson dark matter for the SU(N) gauge group model. Int. J. Theor. Phys. Ser. **52**, 4374 (2013). https://doi.org/10.1007/s10773-013-1756-3
- S. Fraser, E. Ma, M. Zakeri, SU(2)_N model of vector dark matter with a leptonic connection. Int. J. Mod. Phys. A Ser. 30, 1550018 (2015). https://doi.org/10.1142/S0217751X15500189. arXiv:1409.1162
- 104. P. Ko, W.-I. Park, Y. Tang, Higgs portal vector dark matter for GeV scale γ -ray excess from galactic center. JCAP Ser. **09**, 013 (2014). https://doi.org/10.1088/1475-7516/2014/09/013. arXiv:1404.5257
- W.-C. Huang, Y.-L.S. Tsai, T.-C. Yuan, G2HDM: gauged two Higgs doublet model. JHEP Ser. 04, 019 (2016). https://doi.org/ 10.1007/JHEP04(2016)019. arXiv:1512.00229
- C. Gross, O. Lebedev, Y. Mambrini, Non-Abelian gauge fields as dark matter. JHEP Ser. 08, 158 (2015). https://doi.org/10.1007/ JHEP08(2015)158. arXiv:1505.07480
- A. DiFranzo, P.J. Fox, T.M.P. Tait, Vector dark matter through a radiative Higgs portal. JHEP Ser. 04, 135 (2016). https://doi.org/ 10.1007/JHEP04(2016)135. arXiv:1512.06853
- P. Ko, Y. Tang, Residual non-Abelian dark matter and dark radiation. Phys. Lett. B Ser. 768, 12 (2017). https://doi.org/10.1016/j.physletb.2017.02.033. arXiv:1609.02307
- 109. B. Barman, S. Bhattacharya, S.K. Patra, J. Chakrabortty, Non-Abelian vector boson dark matter, its unified route and signatures at the LHC. JCAP Ser. 12, 021 (2017). https://doi.org/10.1088/1475-7516/2017/12/021. arXiv:1704.04945
- 110. W.-C. Huang, H. Ishida, C.-T. Lu, Y.-L.S. Tsai, T.-C. Yuan, Signals of new gauge bosons in gauged two Higgs doublet model. Eur. Phys. J. C Ser. 78, 613 (2018). https://doi.org/10.1140/epjc/s10052-018-6067-7. arXiv:1708.02355
- 111. B. Barman, S. Bhattacharya, M. Zakeri, Multipartite dark matter in $SU(2)_N$ extension of Standard Model and signatures at the LHC. JCAP Ser. **09**, 023 (2018). https://doi.org/10.1088/1475-7516/2018/09/023. arXiv:1806.01129
- B. Barman, S. Bhattacharya, M. Zakeri, Non-Abelian vector boson as FIMP dark matter. JCAP Ser. 02, 029 (2020). https:// doi.org/10.1088/1475-7516/2020/02/029. arXiv:1905.07236



Eur. Phys. J. C (2025) 85:975 Page 87 of 96 975

 D. Buttazzo, L. Di Luzio, P. Ghorbani, C. Gross, G. Landini, A. Strumia et al., Scalar gauge dynamics and dark matter. JHEP Ser. 01, 130 (2020). https://doi.org/10.1007/JHEP01(2020)130. arXiv:1911.04502

- 114. T. Abe, M. Fujiwara, J. Hisano, K. Matsushita, A model of electroweakly interacting non-abelian vector dark matter. JHEP Ser. 07, 136 (2020). https://doi.org/10.1007/JHEP07(2020)136. arXiv:2004.00884
- C. Gross, S. Karamitsos, G. Landini, A. Strumia, Gravitational vector dark matter. JHEP Ser. 03, 174 (2021). https://doi.org/10. 1007/JHEP03(2021)174. arXiv:2012.12087
- T.A. Chowdhury, S. Saad, Non-Abelian vector dark matter and lepton g-2. JCAP Ser. 10, 014 (2021). https://doi.org/10.1088/ 1475-7516/2021/10/014. arXiv:2107.11863
- 117. N. Baouche, A. Ahriche, G. Faisel, S. Nasri, Phenomenology of the hidden SU(2) vector dark matter model. Phys. Rev. D Ser. 104, 075022 (2021). https://doi.org/10.1103/PhysRevD.104.075022. arXiv:2105.14387
- Z. Hu, C. Cai, Y.-L. Tang, Z.-H. Yu, H.-H. Zhang, Vector dark matter from split SU(2) gauge bosons. JHEP Ser. 07, 089 (2021). https://doi.org/10.1007/JHEP07(2021)089. arXiv:2103.00220
- K.S. Babu, S. Jana, A. Thapa, Vector boson dark matter from trinification. JHEP Ser. 02, 051 (2022). https://doi.org/10.1007/ JHEP02(2022)051. arXiv:2112.12771
- G. Arcadi, A. Djouadi, M. Kado, The Higgs-portal for vector dark matter and the effective field theory approach: a reappraisal. Phys. Lett. B Ser. 805, 135427 (2020). https://doi.org/10.1016/j. physletb.2020.135427. arXiv:2001.10750
- S. Baek, P. Ko, P. Wu, Heavy quark-philic scalar dark matter with a vector-like fermion portal. JCAP Ser. 07, 008 (2018). https://doi.org/10.1088/1475-7516/2018/07/008. arXiv:1709.00697
- 122. J. Hisano, A. Ibarra, R. Nagai, Direct detection of vector dark matter through electromagnetic multipoles. JCAP Ser. 10, 015 (2020). https://doi.org/10.1088/1475-7516/2020/10/015. arXiv:2007.03216
- A. Semenov, LanHEP: a package for the automatic generation of Feynman rules in field theory. Version 3.0. Comput. Phys. Commun. 180, 431 (2009). https://doi.org/10.1016/j.cpc.2008.10.012. arXiv:0805.0555
- 124. G. Brooijmans et al., Les Houches 2011: physics at TeV colliders New Physics Working Group Report. in 7th Les Houches Workshop on Physics at TeV Colliders, vol. 3 (2012), pp. 221–463. arXiv:1203.1488
- D. Berdine, N. Kauer, D. Rainwater, Breakdown of the narrow width approximation for new physics. Phys. Rev. Lett. Ser. 99, 111601 (2007). https://doi.org/10.1103/PhysRevLett.99.111601. arXiv:hep-ph/0703058
- NNPDF Collaboration, The path to proton structure at 1% accuracy. Eur. Phys. J. C Ser. 82, 428 (2022). https://doi.org/10.1140/epjc/s10052-022-10328-7. arXiv:2109.02653
- 127. A. Buckley, J. Ferrando, S. Lloyd, K. Nordström, B. Page, M. Rüfenacht et al., LHAPDF6: parton density access in the LHC precision era. Eur. Phys. J. C Ser. 75, 132 (2015). https://doi.org/10.1140/epjc/s10052-015-3318-8. arXiv:1412.7420
- P. Artoisenet, R. Frederix, O. Mattelaer, R. Rietkerk, Automatic spin-entangled decays of heavy resonances in Monte Carlo simulations. JHEP Ser. 03, 015 (2013). https://doi.org/10.1007/JHEP03(2013)015. arXiv:1212.3460
- 129. J. Alwall, C. Duhr, B. Fuks, O. Mattelaer, D.G. Öztürk, C.-H. Shen, Computing decay rates for new physics theories with Feyn-Rules and MadGraph 5_aMC@NLO. Comput. Phys. Commun. Ser. 197, 312 (2015). https://doi.org/10.1016/j.cpc.2015.08.031. arXiv:1402.1178
- C. Bierlich et al., A comprehensive guide to the physics and usage of PYTHIA 8.3. SciPost Phys. Codeb. 2022, 8 (2022). https://doi. org/10.21468/SciPostPhysCodeb.8. arXiv:2203.11601

- S. Frixione, B.R. Webber, Matching NLO QCD computations and parton shower simulations. JHEP Ser. 06, 029 (2002). https://doi. org/10.1088/1126-6708/2002/06/029. arXiv:hep-ph/0204244
- C. Borschensky, B. Fuks, A. Kulesza, D. Schwartländer, Scalar leptoquark pair production at the LHC: precision predictions in the era of flavour anomalies. JHEP Ser. 02, 157 (2022). https:// doi.org/10.1007/JHEP02(2022)157. arXiv:2108.11404
- 133. ATLAS Collaboration, Search for new phenomena in events with an energetic jet and missing transverse momentum in pp collisions at \sqrt{s} =13 TeV with the ATLAS detector. Phys. Rev. D Ser. **103**, 112006 (2021). https://doi.org/10.1103/PhysRevD.103. 112006. arXiv:2102.10874
- 134. ATLAS Collaboration, Search for new phenomena in final states with large jet multiplicities and missing transverse momentum using $\sqrt{s}=13$ TeV proton-proton collisions recorded by ATLAS in Run 2 of the LHC. JHEP Ser. **10**, 062 (2020). https://doi.org/10.1007/JHEP10(2020)062. arXiv:2008.06032
- 135. ATLAS Collaboration, Search for squarks and gluinos in final states with jets and missing transverse momentum using 139 fb⁻¹ of \sqrt{s} =13 TeV pp collision data with the ATLAS detector
- CMS Collaboration, Search for supersymmetry in proton-proton collisions at 13 TeV in final states with jets and missing transverse momentum. JHEP Ser. 10, 244 (2019). https://doi.org/10.1007/ JHEP10(2019)244. arXiv:1908.04722
- 137. CMS Collaboration, Search for new particles in events with energetic jets and large missing transverse momentum in protonproton collisions at $\sqrt{s} = 13$ TeV. JHEP Ser. **11**, 153 (2021). https://doi.org/10.1007/JHEP11(2021)153. arXiv:2107.13021
- E. Conte, B. Fuks, G. Serret, MadAnalysis 5, a user-friendly framework for collider phenomenology. Comput. Phys. Commun. Ser. 184, 222 (2013). https://doi.org/10.1016/j.cpc.2012.09.009. arXiv:1206.1599
- E. Conte, B. Dumont, B. Fuks, C. Wymant, Designing and recasting LHC analyses with MadAnalysis 5. Eur. Phys. J. C Ser. 74, 3103 (2014). https://doi.org/10.1140/epjc/s10052-014-3103-0.arXiv:1405.3982
- E. Conte, B. Fuks, Confronting new physics theories to LHC data with MADANALYSIS 5. Int. J. Mod. Phys. A Ser. 33, 1830027 (2018). https://doi.org/10.1142/S0217751X18300272. arXiv:1808.00480
- M. Cacciari, G.P. Salam, G. Soyez, FastJet user manual. Eur. Phys. J. C Ser. 72, 1896 (2012). https://doi.org/10.1140/epjc/ s10052-012-1896-2. arXiv:1111.6097
- 142. M. Cacciari, G.P. Salam, G. Soyez, The anti-k_t jet clustering algorithm. JHEP Ser. **04**, 063 (2008). https://doi.org/10.1088/1126-6708/2008/04/063. arXiv:0802.1189
- DELPHES 3 Collaboration, DELPHES 3, A modular framework for fast simulation of a generic collider experiment. JHEP 02 057 (2014). https://doi.org/10.1007/JHEP02(2014)057. arXiv:1307.6346
- 144. J.Y. Araz, B. Fuks, G. Polykratis, Simplified fast detector simulation in MADANALYSIS 5. Eur. Phys. J. C Ser. 81, 329 (2021). https://doi.org/10.1140/epjc/s10052-021-09052-5. arXiv:2006.09387
- 145. B. Dumont, B. Fuks, S. Kraml, S. Bein, G. Chalons, E. Conte et al., Toward a public analysis database for LHC new physics searches using MADANALYSIS 5. Eur. Phys. J. C Ser. 75, 56 (2015). https://doi.org/10.1140/epjc/s10052-014-3242-3. arXiv:1407.3278
- 146. J.Y. Araz, M. Frank, B. Fuks, Reinterpreting the results of the LHC with MadAnalysis 5: uncertainties and higher-luminosity estimates. Eur. Phys. J. C Ser. 80, 531 (2020). https://doi.org/10. 1140/epjc/s10052-020-8076-6. arXiv:1910.11418
- 147. J. Kim, T. Lee, J. Kim, H. Jang, Implementation of the ATLAS-SUSY-2018-06 analysis in the MadAnalysis 5 framework (electroweakinos with Jigsaw variables; 139 fb⁻¹). Mod.



975 Page 88 of 96 Eur. Phys. J. C (2025) 85:975

- Phys. Lett. A Ser. **36**, 2141004 (2021). https://doi.org/10.1142/ S0217732321410042
- 148. M. Mrowietz, S. Bein, J. Sonneveld, Implementation of the CMS-SUS-19-006 analysis in the MadAnalysis 5 framework (supersymmetry with large hadronic activity and missing transverse energy; 137 fb⁻¹). Mod. Phys. Lett. A Ser. 36, 2141007 (2021). https://doi.org/10.1142/S0217732321410078
- B. Fuks et al., Proceedings of the second MadAnalysis 5 workshop on LHC recasting in Korea. Mod. Phys. Lett. A Ser. 36, 2102001 (2021). https://doi.org/10.1142/S0217732321020016. arXiv:2101.02245
- 150. L.M. Carpenter, T. Murphy, M.J. Smylie, $t\bar{t}t\bar{t}$ signatures through the lens of color-octet scalars. JHEP Ser. **01**, 047 (2022). https://doi.org/10.1007/JHEP01(2022)047. arXiv:2107.13565
- F. Ambrogi, Implementation of a search for squarks and gluinos in the multi-jet + missing energy channel (139 fb-1; 13 TeV; ATLAS-CONF-2019-040). https://doi.org/10.14428/DVN/NW3NPG
- 152. D. Agin, Implementation of a search for new physics with jets and missing transverse energy (139/fb; 13 TeV; ATLAS-EXOT-2018-06). https://doi.org/10.14428/DVN/REPAMM
- M. Malte, S. Bein, J. Sonneveld, Re-implementation of a search for supersymmetry in the HT/missing HT channel (137 fb-1; CMS-SUSY-19-006). https://doi.org/10.14428/DVN/4DEJQM
- 154. A. Albert, Implementation of a search for new phenomena in events featuring energetic jets and missing transverse energy (137 fb-1; 13 TeV; CMS-EXO-20-004). https://doi.org/10.14428/ DVN/IRF7ZL
- 155. T. Murphy, Implementation of a search for supersymmetry in the multijet (at least 8 jets) + missing energy channel (139/fb; 13 TeV; ATLAS-SUSY-2018-17). https://doi.org/10.14428/DVN/ 12CZWU
- J.M. Butterworth, D. Grellscheid, M. Krämer, B. Sarrazin, D. Yallup, Constraining new physics with collider measurements of Standard Model signatures. JHEP Ser. 03, 078 (2017). https://doi.org/10.1007/JHEP03(2017)078. arXiv:1606.05296
- 157. A. Buckley et al., Testing new physics models with global comparisons to collider measurements: the Contur toolkit. SciPost Phys. Core 4, 013 (2021). https://doi.org/10.21468/SciPostPhysCore.4. 2.013. arXiv:2102.04377
- 158. C. Bierlich, A. Buckley, J.M. Butterworth, C. Gutschow, L. Lonnblad, T. Procter et al., Robust independent validation of experiment and theory: Rivet version 4 release note. SciPost Phys. Codeb. 36, 1 (2024). https://doi.org/10.21468/ SciPostPhysCodeb.36. arXiv:2404.15984
- A. Verbytskyi, A. Buckley, D. Grellscheid, D. Konstantinov, J. WilliamMonk, L. Lönnblad et al., HepMC3 Event Record Library for Monte Carlo event generators. J. Phys: Conf. Ser. 1525, 012017 (2020). https://doi.org/10.1088/1742-6596/1525/ 1/012017
- 160. ATLAS Collaboration, Measurement of the $Z(\rightarrow \ell^+\ell^-)\gamma$ production cross-section in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector. JHEP Ser. **03**, 054 (2020). https://doi.org/10. 1007/JHEP03(2020)054. arXiv:1911.04813
- 161. ATLAS Collaboration, Differential cross-sections for events with missing transverse momentum and jets measured with the ATLAS detector in 13 TeV proton-proton collisions. JHEP Ser. 08, 223 (2024). https://doi.org/10.1007/JHEP08(2024)223. arXiv:2403.02793
- 162. ATLAS Collaboration, Measurement of detector-corrected observables sensitive to the anomalous production of events with jets and large missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector. Eur. Phys. J. C Ser. 77, 765 (2017). https://doi.org/10.1140/epjc/s10052-017-5315-6. arXiv:1707.03263
- 163. ATLAS Collaboration, Differential $t\bar{t}$ cross-section measurements using boosted top quarks in the all-hadronic final state with

- 139 fb⁻¹ of ATLAS data. JHEP Ser. **04**, 080 (2023). https://doi. org/10.1007/JHEP04(2023)080. arXiv:2205.02817
- 164. CMS Collaboration, Measurement of differential cross sections for the production of top quark pairs and of additional jets in lepton+jets events from pp collisions at $\sqrt{s} = 13$ TeV. Phys. Rev. D Ser. 97, 112003 (2018). https://doi.org/10.1103/PhysRevD.97. 112003. arXiv:1803.08856
- 165. CMS Collaboration, Measurement of differential cross sections for top quark pair production using the lepton+jets final state in proton-proton collisions at 13 TeV. Phys. Rev. D Ser. 95, 092001 (2017). https://doi.org/10.1103/PhysRevD.95. 092001. arXiv:1610.04191
- J.Y. Araz, A. Buckley, B. Fuks, H. Reyes-Gonzalez, W. Waltenberger, S.L. Williamson et al., Strength in numbers: optimal and scalable combination of LHC new-physics searches. SciPost Phys. 14, 077 (2023). https://doi.org/10.21468/SciPostPhys.14.4. 077, arXiv:2209.00025
- A. Feike, J. Fiaschi, B. Fuks, M. Klasen, A. Puck Neuwirth, Combination and reinterpretation of LHC SUSY searches. JHEP 07, 122 (2024). https://doi.org/10.1007/JHEP07(2024)122. arXiv:2403.11715
- M. Mahdi Altakach, S. Kraml, A. Lessa, S. Narasimha, T. Pascal, W. Waltenberger, SModelS v2.3: enabling global likelihood analyses. SciPost Phys. 15, 185 (2023). https://doi.org/10.21468/ SciPostPhys.15.5.185
- 169. M.M. Altakach, S. Kraml, A. Lessa, S. Narasimha, T. Reymermier et al., Global LHC constraints on electroweak-inos with SModelS v2.3. SciPost Phys. 16, 101 (2024). https://doi.org/10.21468/ SciPostPhys.16.4.101. arXiv:2312.16635
- B. Fuks, H.-S. Shao, QCD next-to-leading-order predictions matched to parton showers for vector-like quark models. Eur. Phys. J. C Ser. 77, 135 (2017). https://doi.org/10.1140/epjc/ s10052-017-4686-z. arXiv:1610.04622
- B. Fuks (ed.), Proceedings of the First MadAnalysis 5 Workshop on LHC Recasting in Korea, vol. 6 (2018)
- 172. J.Y. Araz, B. Fuks, M.D. Goodsell, M. Utsch, Recasting LHC searches for long-lived particles with MadAnalysis 5. Eur. Phys. J. C Ser. 82, 597 (2022). https://doi.org/10.1140/epjc/s10052-022-10511-w. arXiv:2112.05163
- 173. ATLAS Collaboration, Search for electroweak production of charginos and sleptons decaying into final states with two leptons and missing transverse momentum in $\sqrt{s}=13$ TeV pp collisions using the ATLAS detector. Eur. Phys. J. C Ser. 80, 123 (2020). https://doi.org/10.1140/epjc/s10052-019-7594-6. arXiv:1908.08215
- 174. ATLAS Collaboration, Searches for electroweak production of supersymmetric particles with compressed mass spectra in \sqrt{s} = 13 TeV pp collisions with the ATLAS detector. Phys. Rev. D Ser. **101**, 052005 (2020). https://doi.org/10.1103/PhysRevD.101. 052005. arXiv:1911.12606
- 175. shape ATLAS collaboration, Search for direct production of charginos, neutralinos and sleptons in final states with two leptons and missing transverse momentum in pp collisions at $\sqrt{s} = 8 \text{ TeV}$ with the ATLAS detector. JHEP Ser. **05**, 071 (2014). https://doi.org/10.1007/JHEP05(2014)071. arXiv:1403.5294
- 176. M. Blanke, P. Pani, G. Polesello, G. Rovelli, Single-top final states as a probe of top-flavoured dark matter models at the LHC. JHEP Ser. 01, 194 (2021). https://doi.org/10.1007/JHEP01(2021)194. arXiv:2010.10530
- J. Andrea, B. Fuks, F. Maltoni, Monotops at the LHC. Phys. Rev. D Ser. 84, 074025 (2011). https://doi.org/10.1103/PhysRevD.84. 074025. arXiv:1106.6199
- PLANCK Collaboration, Planck 2018 results. VI. Cosmological parameters. Astron. Astrophys. 641, A6 (2020). https://doi.org/ 10.1051/0004-6361/201833910. arXiv:1807.06209



Eur. Phys. J. C (2025) 85:975 Page 89 of 96 975

179. ATLAS Collaboration, Search for same-sign top-quark production and fourth-generation down-type quarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. JHEP Ser. **04**, 069 (2012). https://doi.org/10.1007/JHEP04(2012)069. arXiv:1202.5520

- R.D. Ball et al., Parton distributions with LHC data. Nucl. Phys. B Ser. 867, 244 (2013). https://doi.org/10.1016/j.nuclphysb.2012. 10.003. arXiv:1207.1303
- 181. CMS Collaboration, Search for pair-produced resonances decaying to quark pairs in proton-proton collisions at $\sqrt{s}=13$ TeV. Phys. Rev. D Ser. **98**, 112014 (2018). https://doi.org/10.1103/PhysRevD.98.112014. arXiv:1808.03124
- 182. J.M. Campbell, R.K. Ellis, tīW⁺⁻ production and decay at NLO. JHEP Ser. **07**, 052 (2012). https://doi.org/10.1007/JHEP07(2012)052. arXiv:1204.5678
- 183. M.L. Mangano, M. Moretti, F. Piccinini, M. Treccani, Matching matrix elements and shower evolution for top-quark production in hadronic collisions. JHEP Ser. 01, 013 (2007). https://doi.org/ 10.1088/1126-6708/2007/01/013. arXiv:hep-ph/0611129
- 184. J. Alwall, S. de Visscher, F. Maltoni, QCD radiation in the production of heavy colored particles at the LHC. JHEP Ser. 02, 017 (2009). https://doi.org/10.1088/1126-6708/2009/02/017. arXiv:0810.5350
- 185. J.Y. Araz, A. Buckley, B. Fuks, Searches for new physics with boosted top quarks in the MadAnalysis 5 and Rivet frameworks. Eur. Phys. J. C Ser. 83, 664 (2023). https://doi.org/10.1140/epjc/ s10052-023-11779-2. arXiv:2303.03427
- T. Plehn, M. Spannowsky, M. Takeuchi, D. Zerwas, Stop reconstruction with tagged tops. JHEP Ser. 10, 078 (2010). https://doi.org/10.1007/JHEP10(2010)078. arXiv:1006.2833
- 187. G. Kasieczka, T. Plehn, T. Schell, T. Strebler, G.P. Salam, Resonance searches with an updated top tagger. JHEP Ser. 06, 203 (2015). https://doi.org/10.1007/JHEP06(2015)203. arXiv:1503.05921
- Y.L. Dokshitzer, G.D. Leder, S. Moretti, B.R. Webber, Better jet clustering algorithms. JHEP Ser. 08, 001 (1997). https://doi.org/ 10.1088/1126-6708/1997/08/001. arXiv:hep-ph/9707323
- S. Bentvelsen, I. Meyer, The Cambridge jet algorithm: features and applications. Eur. Phys. J. C Ser. 4, 623 (1998). https://doi. org/10.1007/s100520050232. arXiv:hep-ph/9803322
- M. Wobisch, T. Wengler, Hadronization corrections to jet crosssections in deep inelastic scattering. in Workshop on Monte Carlo Generators for HERA Physics (Plenary Starting Meeting), vol. 4 (1998), pp. 270–279. arXiv:hep-ph/9907280
- G. Cowan, K. Cranmer, E. Gross, O. Vitells, Asymptotic formulae for likelihood-based tests of new physics. Eur. Phys. J. C Ser. 71, 1554 (2011). https://doi.org/10.1140/epjc/s10052-011-1554-0. arXiv:1007.1727
- 192. L. Darmé, B. Fuks, F. Maltoni, Top-philic heavy resonances in four-top final states and their EFT interpretation. JHEP Ser. 09, 143 (2021). https://doi.org/10.1007/JHEP09(2021)143. arXiv:2104.09512
- 193. L. Darmé, B. Fuks, H.-L. Li, M. Maltoni, O. Mattelaer, J. Touchèque, Boosting beyond: a novel approach to probing top-philic resonances at the LHC. arXiv:2404.14482
- 194. NNPDF Collaboration, Parton distributions for the LHC Run II. JHEP Ser. 04, 040 (2015). https://doi.org/10.1007/ JHEP04(2015)040. arXiv:1410.8849
- 195. T. Hahn, S. Paßehr, C. Schappacher, FormCalc 9 and extensions. PoS LL2016 068 (2016). https://doi.org/10.1088/1742-6596/762/1/012065. arXiv:1604.04611
- 196. CMS Collaboration, Search for top squarks and dark matter particles in opposite-charge dilepton final states at $\sqrt{s}=13$ TeV. Phys. Rev. D Ser. 97, 032009 (2018). https://doi.org/10.1103/PhysRevD.97.032009. arXiv:1711.00752
- 197. ATLAS Collaboration, Search for pair-production of vector-like quarks in pp collision events at $\sqrt{s} = 13$ TeV with at least one

- leptonically-decaying Z boson and a third-generation quark with the ATLAS detector
- 198. CMS Collaboration, Search for pair production of vector-like quarks in leptonic final states in proton-proton collisions at \sqrt{s} = 13 TeV. JHEP Ser. **07**, 020 (2023). https://doi.org/10.1007/JHEP07(2023)020. arXiv:2209.07327
- 199. J. Alimena et al., Searching for long-lived particles beyond the Standard Model at the large hadron collider. J. Phys. G Ser. 47, 090501 (2020). https://doi.org/10.1088/1361-6471/ab4574. arXiv:1903.04497
- T. Asaka, K. Ishiwata, T. Moroi, Right-handed sneutrino as cold dark matter. Phys. Rev. D Ser. 73, 051301 (2006). https://doi.org/ 10.1103/PhysRevD.73.051301. arXiv:hep-ph/0512118
- M. Garny, J. Heisig, M. Hufnagel, B. Lülf, Top-philic dark matter within and beyond the WIMP paradigm. Phys. Rev. D Ser. 97, 075002 (2018). https://doi.org/10.1103/PhysRevD.97. 075002. arXiv:1802.00814
- M. Garny, J. Heisig, Interplay of super-WIMP and freezein production of dark matter. Phys. Rev. D Ser. 98, 095031 (2018). https://doi.org/10.1103/PhysRevD.98.095031. arXiv:1809.10135
- 203. M. Deshpande, J. Hamann, D. Sengupta, M. White, A.G. Williams, Y.Y.Y. Wong, Revisiting cosmological constraints on supersymmetric SuperWIMPs. Eur. Phys. J. C Ser. 84, 667 (2024). https://doi.org/10.1140/epjc/s10052-024-12992-3. arXiv:2309.05709
- 204. Q. Decant, J. Heisig, D.C. Hooper, L. Lopez-Honorez, Lyman-α constraints on freeze-in and superWIMPs. JCAP Ser. 03, 041 (2022). https://doi.org/10.1088/1475-7516/2022/03/041. arXiv:2111.09321
- 205. G. Ballesteros, M.A.G. Garcia, M. Pierre, How warm are non-thermal relics? Lyman-α bounds on out-of-equilibrium dark matter. JCAP Ser. 03, 101 (2021). https://doi.org/10.1088/1475-7516/2021/03/101. arXiv:2011.13458
- F. D'Eramo, A. Lenoci, Lower mass bounds on FIMP dark matter produced via freeze-in. JCAP Ser. 10, 045 (2021). https://doi.org/ 10.1088/1475-7516/2021/10/045. arXiv:2012.01446
- 207. ATLAS Collaboration, Search for heavy, long-lived, charged particles with large ionisation energy loss in pp collisions at $\sqrt{s}=13$ TeV using the ATLAS experiment and the full Run 2 dataset. JHEP Ser. **2306**, 158 (2023). https://doi.org/10.1007/JHEP06(2023)158. arXiv:2205.06013
- 208. CMS Collaboration, Search for heavy long-lived charged particles with large ionization energy loss in proton-proton collisions at \sqrt{s} = 13 TeV. arXiv:2410.09164
- 209. CMS Collaboration, Search for supersymmetry in final states with disappearing tracks in proton-proton collisions at s=13 TeV. Phys. Rev. D Ser. 109, 072007 (2024). https://doi.org/10.1103/PhysRevD.109.072007. arXiv:2309.16823
- 210. ATLAS Collaboration, Search for long-lived charginos based on a disappearing-track signature using 136 fb⁻¹ of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. Eur. Phys. J. C Ser. 82, 606 (2022). https://doi.org/10.1140/epjc/s10052-022-10489-5. arXiv:2201.02472
- 211. CMS Collaboration, Search for long-lived particles using displaced vertices and missing transverse momentum in proton-proton collisions at s=13 TeV. Phys. Rev. D Ser. 109, 112005 (2024). https://doi.org/10.1103/PhysRevD.109.112005. arXiv:2402.15804
- 212. ATLAS Collaboration, Search for long-lived, massive particles in events with displaced vertices and missing transverse momentum in $\sqrt{s} = 13 \text{ TeV } pp$ collisions with the ATLAS detector. Phys. Rev. D Ser. 97, 052012 (2018). https://doi.org/10.1103/PhysRevD.97. 052012. arXiv:1710.04901
- ATLAS Collaboration, Search for nearly mass-degenerate Higgsinos using low-momentum mildly displaced tracks in pp col-



975 Page 90 of 96 Eur. Phys. J. C (2025) 85:975

- lisions at s=13 TeV with the ATLAS detector. Phys. Rev. Lett. Ser. **132**, 221801 (2024). https://doi.org/10.1103/PhysRevLett. 132.221801. arXiv:2401.14046
- 214. ATLAS Collaboration, Search for displaced leptons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector. Phys. Rev. Lett. Ser. **127**, 051802 (2021). https://doi.org/10.1103/PhysRevLett. 127.051802. arXiv:2011.07812
- 215. CMS Collaboration, Search for long-lived particles decaying to leptons with large impact parameter in proton-proton collisions at $\sqrt{s} = 13$ TeV. Eur. Phys. J. C Ser. **82**, 153 (2022). https://doi.org/10.1140/epjc/s10052-022-10027-3. arXiv:2110.04809
- 216. ATLAS Collaboration, Search for neutral long-lived particles in pp collisions at $\sqrt{s} = 13$ TeV that decay into displaced hadronic jets in the ATLAS calorimeter. JHEP Ser. **06**, 005 (2022). https://doi.org/10.1007/JHEP06(2022)005. arXiv:2203.01009
- 217. CMS Collaboration, Search for long-lived particles decaying in the CMS muon detectors in proton-proton collisions at s=13 TeV. Phys. Rev. D Ser. 110, 032007 (2024). https://doi.org/10.1103/ PhysRevD.110.032007. arXiv:2402.01898
- 218. ATLAS Collaboration, Search for events with a pair of displaced vertices from long-lived neutral particles decaying into hadronic jets in the ATLAS muon spectrometer in pp collisions at √s=13 TeV. Phys. Rev. D Ser. 106, 032005 (2022). https://doi.org/10.1103/PhysRevD.106.032005. arXiv:2203.00587
- 219. CMS Collaboration, Constraints on the pMSSM, AMSB model and on other models from the search for long-lived charged particles in proton-proton collisions at sqrt(s) = 8 TeV. Eur. Phys. J. C Ser. 75, 325 (2015). https://doi.org/10.1140/epjc/ s10052-015-3533-3. arXiv:1502.02522
- J. Heisig, J. Kersten, Long-lived staus from strong production in a simplified model approach. Phys. Rev. D Ser. 86, 055020 (2012). https://doi.org/10.1103/PhysRevD.86.055020. arXiv:1203.1581
- J. Heisig, A. Lessa, L. Quertenmont, Simplified models for exotic BSM searches. JHEP Ser. 12, 087 (2015). https://doi.org/10.1007/ JHEP12(2015)087. arXiv:1509.00473
- G.F. Giudice, M. McCullough, D. Teresi, dE/dx from boosted long-lived particles. JHEP Ser. 08, 012 (2022). https://doi.org/10. 1007/JHEP08(2022)012. arXiv:2205.04473
- 223. CMS Collaboration, Search for disappearing tracks in proton-proton collisions at $\sqrt{s}=13$ TeV. Phys. Lett. B Ser. **806**, 135502 (2020). https://doi.org/10.1016/j.physletb.2020.135502. arXiv:2004.05153
- 224. ATLAS Collaboration, Search for pairs of muons with small displacements in pp collisions at s=13 TeV with the ATLAS detector. Phys. Lett. B Ser. **846**, 138172 (2023). https://doi.org/10.1016/j.physletb.2023.138172. arXiv:2305.02005
- 225. ATLAS Collaboration, Search for displaced leptons in $\sqrt{s} = 13$ TeV and 13.6 TeV pp collisions with the ATLAS detector. arXiv:2410.16835
- 226. CMS Collaboration, Search for long-lived particles decaying to a pair of muons in proton-proton collisions at $\sqrt{s} = 13$ TeV. JHEP Ser. **05**, 228 (2023). https://doi.org/10.1007/JHEP05(2023)228. arXiv:2205.08582
- 227. shape CMS collaboration, Search for long-lived heavy neutral leptons with displaced vertices in proton-proton collisions at \sqrt{s} =13 TeV. JHEP Ser. **07**, 081 (2022). https://doi.org/10.1007/JHEP07(2022)081. arXiv:2201.05578
- 228. CMS Collaboration, Search for long-lived particles decaying to final states with a pair of muons in proton-proton collisions at $\sqrt{s} = 13.6$ TeV. JHEP **05** 047 (2024). https://doi.org/10.1007/JHEP05(2024)047. arXiv:2402.14491
- 229. CMS Collaboration, Search for long-lived particles decaying into muon pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV collected with a dedicated high-rate data stream. JHEP Ser. **04**, 062 (2022). https://doi.org/10.1007/JHEP04(2022)062. arXiv:2112.13769

- ATLAS Collaboration, Performance of the reconstruction of large impact parameter tracks in the inner detector of ATLAS. Eur. Phys. J. C Ser. 83, 1081 (2023). https://doi.org/10.1140/epjc/ s10052-023-12024-6. arXiv:2304.12867
- CMS Collaboration, Enriching the physics program of the CMS Experiment via data scouting and data parking. arXiv:2403.16134
- 232. ATLAS Collaboration, Search for long-lived, massive particles in events with displaced vertices and multiple jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. JHEP Ser. **06**, 200 (2023). https://doi.org/10.1007/JHEP06(2023)200. arXiv:2301.13866
- 233. ATLAS Collaboration, Search for neutral long-lived particles that decay into displaced jets in the ATLAS calorimeter in association with leptons or jets using pp collisions at $\sqrt{s} = 13$ TeV. JHEP Ser. 11, 036 (2024). https://doi.org/10.1007/JHEP11(2024)036. arXiv:2407.09183
- 234. ATLAS Collaboration, Search for light long-lived particles in pp collisions at s=13 TeV using displaced vertices in the ATLAS inner detector. Phys. Rev. Lett. Ser. 133, 161803 (2024). https://doi.org/10.1103/PhysRevLett.133.161803. arXiv:2403.15332
- 235. CMS Collaboration, Search for long-lived particles using displaced jets in proton-proton collisions at $\sqrt{s}=13$ TeV. Phys. Rev. D Ser. **104**, 012015 (2021). https://doi.org/10.1103/PhysRevD. 104.012015. arXiv:2012.01581
- 236. CMS Collaboration, Search for long-lived particles decaying to jets with displaced vertices in proton-proton collisions at \sqrt{s} = 13 TeV. Phys. Rev. D Ser. **104**, 052011 (2021). https://doi.org/10. 1103/PhysRevD.104.052011. arXiv:2104.13474
- 237. CMS Collaboration, Search for long-lived particles using non-prompt jets and missing transverse momentum with proton-proton collisions at $\sqrt{s}=13$ TeV. Phys. Lett. B Ser. **797**, 134876 (2019). https://doi.org/10.1016/j.physletb.2019.134876. arXiv:1906.06441
- 238. CMS Collaboration, Search for long-lived particles decaying into displaced jets in proton-proton collisions at $\sqrt{s}=13$ TeV. Phys. Rev. D Ser. **99**, 032011 (2019). https://doi.org/10.1103/PhysRevD.99.032011. arXiv:1811.07991
- 239. CMS Collaboration, Search for new particles decaying to a jet and an emerging jet. JHEP Ser. **02**, 179 (2019). https://doi.org/10.1007/JHEP02(2019)179. arXiv:1810.10069
- 240. CMS Collaboration, Search for heavy stable charged particles with 12.9 ${\rm fb^{-1}}$ of 2016 data
- 241. ATLAS Collaboration, Search for heavy charged long-lived particles in the ATLAS detector in 36.1 fb⁻¹ of proton-proton collision data at $\sqrt{s} = 13$ TeV. Phys. Rev. D **99**, 092007 (2019). https://doi.org/10.1103/PhysRevD.99.092007. arXiv:1902.01636
- 242. ATLAS Collaboration, Search for long-lived charginos based on a disappearing-track signature in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector. JHEP Ser. **06**, 022 (2018). https://doi.org/10.1007/JHEP06(2018)022. arXiv:1712.02118
- 243. CMS Collaboration, Search for disappearing tracks as a signature of new long-lived particles in proton-proton collisions at $\sqrt{s}=13$ TeV. JHEP Ser. **08**, 016 (2018). https://doi.org/10.1007/JHEP08(2018)016. arXiv:1804.07321
- 244. ATLAS Collaboration, Search for long-lived, massive particles in events with a displaced vertex and a muon with large impact parameter in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. Phys. Rev. D Ser. **102**, 032006 (2020). https://doi.org/10.1103/PhysRevD.102.032006. arXiv:2003.11956
- 245. "LLP Recasting Repository." https://github.com/llprecasting/ recastingCodes
- 246. CheckMATE Collaboration, Constraining electroweak and strongly charged long-lived particles with CheckMATE. Eur. Phys. J. C Ser. 81, 968 (2021). https://doi.org/10.1140/epjc/ s10052-021-09727-z. arXiv:2104.04542
- J. Heisig, S. Kraml, A. Lessa, Constraining new physics with searches for long-lived particles: implementation into SMod-



Eur. Phys. J. C (2025) 85:975 Page 91 of 96 975

elS. Phys. Lett. B Ser. **788**, 87 (2019). https://doi.org/10.1016/j.physletb.2018.10.049. arXiv:1808.05229

- F. Ambrogi et al., SModelS v1.2: long-lived particles, combination of signal regions, and other novelties. Comput. Phys. Commun. 251, 106848 (2020). https://doi.org/10.1016/j.cpc.2019.07.013. arXiv:1811.10624
- G. Alguero, J. Heisig, C.K. Khosa, S. Kraml, S. Kulkarni, A. Lessa et al., Constraining new physics with SModelS version 2. JHEP Ser. 08, 068 (2022). https://doi.org/10.1007/JHEP08(2022)068. arXiv:2112.00769
- 250. M.M. Altakach, S. Kraml, A. Lessa, S. Narasimha, T. Pascal, C. Ramos et al., SModelS v3: going beyond Z₂ topologies. JHEP Ser. 11, 074 (2024). https://doi.org/10.1007/JHEP11(2024)074. arXiv:2409.12942
- G. Bélanger et al., LHC-friendly minimal freeze-in models. JHEP Ser. 02, 186 (2019). https://doi.org/10.1007/JHEP02(2019)186. arXiv:1811.05478
- L. Calibbi, F. D'Eramo, S. Junius, L. Lopez-Honorez, A. Mariotti, Displaced new physics at colliders and the early universe before its first second. JHEP Ser. 05, 234 (2021). https://doi.org/10.1007/ JHEP05(2021)234. arXiv:2102.06221
- S. Junius, Searching for feebly interacting dark matter in colliders and beam-dump experiments. Ph.D. thesis, U. Brussels, U. Brussels (main) (2022). arXiv:2212.09432
- M. Becker, E. Copello, J. Harz, J. Lang, Y. Xu, Confronting dark matter freeze-in during reheating with constraints from inflation. JCAP Ser. 01, 053 (2024). https://doi.org/10.1088/1475-7516/ 2024/01/053. arXiv:2306.17238
- R.T. Co, F. D'Eramo, L.J. Hall, D. Pappadopulo, Freeze-in dark matter with displaced signatures at colliders. JCAP Ser. 12, 024 (2015). https://doi.org/10.1088/1475-7516/2015/12/024. arXiv:1506.07532
- 256. G. Brooijmans et al., Les Houches 2019 physics at TeV colliders: New Physics Working Group Report. in 11th Les Houches Workshop on Physics at TeV Colliders: PhysTeV Les Houches, vol. 2 (2020), arXiv:2002.12220
- 257. F. D'Eramo, N. Fernandez, S. Profumo, Dark matter freezein production in fast-expanding universes. JCAP Ser. 02, 046 (2018). https://doi.org/10.1088/1475-7516/2018/02/046. arXiv:1712.07453
- 258. M. Garny, J. Heisig, Bound-state effects on dark matter coannihilation: pushing the boundaries of conversion-driven freeze-out. Phys. Rev. D Ser. 105, 055004 (2022). https://doi.org/10.1103/PhysRevD.105.055004. arXiv:2112.01499
- J. Heisig, A. Lessa, L.M.D. Ramos, Probing conversiondriven freeze-out at the LHC. Phys. Rev. D Ser. 110, 015031 (2024). https://doi.org/10.1103/PhysRevD.110.015031. arXiv:2404.16086
- S. Junius, L. Lopez-Honorez, A. Mariotti, A feeble window on leptophilic dark matter. JHEP Ser. 07, 136 (2019). https://doi.org/ 10.1007/JHEP07(2019)136. arXiv:1904.07513
- T. Binder, M. Garny, J. Heisig, S. Lederer, K. Urban, Excited bound states and their role in dark matter production. Phys. Rev. D Ser. 108, 095030 (2023). https://doi.org/10.1103/PhysRevD. 108.095030. arXiv:2308.01336
- 262. CMS Collaboration, Search for inelastic dark matter in events with two displaced muons and missing transverse momentum in proton-proton collisions at s=13 TeV. Phys. Rev. Lett. Ser. 132, 041802 (2024). https://doi.org/10.1103/PhysRevLett.132. 041802. arXiv:2305.11649
- 263. G. Belanger et al., Leptoquark Manoeuvres in the dark: a simultaneous solution of the dark matter problem and the $R_{D^{(*)}}$ anomalies. JHEP Ser. **02**, 042 (2022). https://doi.org/10.1007/JHEP02(2022)042. arXiv:2111.08027
- 264. D. Curtin et al., Long-lived particles at the energy frontier: the MATHUSLA physics case. Rep. Prog. Phys. Ser.

- 82, 116201 (2019). https://doi.org/10.1088/1361-6633/ab28d6. arXiv:1806.07396
- 265. LHC REINTERPRETATION FORUM Collaboration, Reinterpretation of LHC results for new physics: status and recommendations after Run 2. SciPost Phys. 9 022 (2020). https://doi.org/10.21468/SciPostPhys.9.2.022. arXiv:2003.07868
- J.Y. Araz et al., Les Houches guide to reusable ML models in LHC analyses. arXiv:2312.14575
- K. Cranmer et al., Publishing statistical models: getting the most out of particle physics experiments. SciPost Phys. 12, 037 (2022). https://doi.org/10.21468/SciPostPhys.12.1.037. arXiv:2109.04981
- 268. ATLAS Collaboration, Reproducing searches for new physics with the ATLAS experiment through publication of full statistical likelihoods
- CMS Collaboration, The CMS statistical analysis and combination tool: combine. Comput. Softw. Big Sci. Ser. 8, 19 (2024). https://doi.org/10.1007/s41781-024-00121-4. arXiv:2404.06614
- C. Balazs, T. Bringmann, F. Kahlhoefer, M. White, A primer on dark matter. arXiv:2411.05062
- P. Gondolo, G. Gelmini, Cosmic abundances of stable particles: improved analysis. Nucl. Phys. B Ser. 360, 145 (1991). https://doi.org/10.1016/0550-3213(91)90438-4
- K. Griest, D. Seckel, Three exceptions in the calculation of relic abundances. Phys. Rev. D Ser. 43, 3191 (1991). https://doi.org/ 10.1103/PhysRevD.43.3191
- 273. D.J.H. Chung, E.W. Kolb, A. Riotto, Production of massive particles during reheating. Phys. Rev. D Ser. 60, 063504 (1999). https://doi.org/10.1103/PhysRevD.60.063504. arXiv:hep-ph/9809453
- 274. G.F. Giudice, E.W. Kolb, A. Riotto, Largest temperature of the radiation era and its cosmological implications. Phys. Rev. D Ser. 64, 023508 (2001). https://doi.org/10.1103/PhysRevD.64. 023508. arXiv:hep-ph/0005123
- 275. R. Allahverdi et al., The first three seconds: a review of possible expansion histories of the early universe. arXiv:2006.16182
- M. Drees, F. Hajkarim, Dark matter production in an early matter dominated era. JCAP Ser. 02, 057 (2018). https://doi.org/10.1088/ 1475-7516/2018/02/057. arXiv:1711.05007
- N. Bernal, Y. Xu, WIMPs during reheating. JCAP Ser. 12,017 (2022). https://doi.org/10.1088/1475-7516/2022/12/017. arXiv:2209.07546
- J. Edsjo, P. Gondolo, Neutralino relic density including coannihilations. Phys. Rev. D Ser. 56, 1879 (1997). https://doi.org/10.1103/PhysRevD.56.1879. arXiv:hep-ph/9704361
- T. Bringmann, S. Hofmann, Thermal decoupling of WIMPs from first principles. JCAP Ser. 04, 016 (2007). https://doi.org/10.1088/ 1475-7516/2007/04/016. arXiv:hep-ph/0612238
- 280. T. Binder, T. Bringmann, M. Gustafsson, A. Hryczuk, Early kinetic decoupling of dark matter: when the standard way of calculating the thermal relic density fails. Phys. Rev. D Ser. 96, 115010 (2017). https://doi.org/10.1103/PhysRevD.96. 115010. arXiv:1706.07433
- M. Beneke, F. Dighera, A. Hryczuk, Relic density computations at NLO: infrared finiteness and thermal correction. JHEP Ser. 10, 045 (2014). https://doi.org/10.1007/JHEP10(2014)045. arXiv:1409.3049
- M. Drewes, J.U. Kang, Sterile neutrino dark matter production from scalar decay in a thermal bath. JHEP Ser. 05, 051 (2016). https://doi.org/10.1007/JHEP05(2016)051. arXiv:1510.05646
- T. Binder, L. Covi, K. Mukaida, Dark matter Sommerfeldenhanced annihilation and bound-state decay at finite temperature. Phys. Rev. D Ser. 98, 115023 (2018). https://doi.org/10. 1103/PhysRevD.98.115023. arXiv:1808.06472
- M. Becker, E. Copello, J. Harz, C. Tamarit, Dark matter freezein from non-equilibrium QFT: towards a consistent treatment of thermal effects. arXiv:2312.17246



975 Page 92 of 96 Eur. Phys. J. C (2025) 85:975

- W.-Y. Ai, A. Beniwal, A. Maggi, D.J.E. Marsh, From QFT to Boltzmann: freeze-in in the presence of oscillating condensates. JHEP Ser. 02, 122 (2024). https://doi.org/10.1007/JHEP02(2024)122. arXiv:2310.08272
- K. Kainulainen, S. Nurmi, O. Väisänen, Tachyonic production of dark relics: classical lattice vs. quantum 2PI in Hartree truncation. JHEP 10, 009 (2024). https://doi.org/10.1007/JHEP10(2024)009. arXiv:2406.17468
- M. Becker, J. Harz, E. Morgante, C. Puchades-Ibáñez, P. Schwaller, ALP Production from Abelian gauge bosons: beyond hard thermal loops. arXiv:2502.01729
- S. Kim, M. Laine, On thermal corrections to near-threshold annihilation. JCAP Ser. 01, 013 (2017). https://doi.org/10.1088/ 1475-7516/2017/01/013. arXiv:1609.00474
- S. Biondini, J. Ghiglieri, Freeze-in produced dark matter in the ultra-relativistic regime. JCAP Ser. 03, 075 (2021). https://doi. org/10.1088/1475-7516/2021/03/075. arXiv:2012.09083
- F. D'Eramo, F. Hajkarim, S. Yun, Thermal QCD axions across thresholds. JHEP Ser. 10, 224 (2021). https://doi.org/10.1007/ JHEP10(2021)224. arXiv:2108.05371
- K. Bouzoud, J. Ghiglieri, Thermal axion production at hard and soft momenta. JHEP Ser. 01, 163 (2025). https://doi.org/10.1007/ JHEP01(2025)163. arXiv:2404.06113
- 292. M. Beneke, A. Bharucha, A. Hryczuk, S. Recksiegel, P. Ruiz-Femenia, The last refuge of mixed wino-Higgsino dark matter. JHEP Ser. 01, 002 (2017). https://doi.org/10.1007/JHEP01(2017)002. arXiv:1611.00804
- B. von Harling, K. Petraki, Bound-state formation for thermal relic dark matter and unitarity. JCAP Ser. 12, 033 (2014). https:// doi.org/10.1088/1475-7516/2014/12/033. arXiv:1407.7874
- T. Binder, B. Blobel, J. Harz, K. Mukaida, Dark matter boundstate formation at higher order: a non-equilibrium quantum field theory approach. JHEP Ser. 09, 086 (2020). https://doi.org/10. 1007/JHEP09(2020)086. arXiv:2002.07145
- T. Binder, K. Mukaida, K. Petraki, Rapid bound-state formation of dark matter in the early universe. Phys. Rev. Lett. Ser. 124, 161102 (2020). https://doi.org/10.1103/PhysRevLett.124. 161102. arXiv:1910.11288
- S. Biondini, N. Brambilla, G. Qerimi, A. Vairo, Effective field theories for dark matter pairs in the early universe: cross sections and widths. JHEP Ser. 07, 006 (2023). https://doi.org/10.1007/ JHEP07(2023)006. arXiv:2304.00113
- S. Biondini, N. Brambilla, G. Qerimi, A. Vairo, Effective field theories for dark matter pairs in the early universe: center-of-mass recoil effects. JHEP Ser. 07, 021 (2024). https://doi.org/10.1007/ JHEP07(2024)021. arXiv:2402.12787
- S. Biondini, N. Brambilla, A. Dashko, G. Qerimi, A. Vairo, Effective field theories for dark matter pairs in the early universe: Debye mass effects. arXiv:2501.03327
- J. Heisig, Dark matter and long-lived particles at the LHC. in 53rd Rencontres de Moriond on Electroweak Interactions and Unified Theories (2018), pp. 213–220. arXiv:1805.07361
- L. Lopez-Honorez, Q. Decant, S. Junius, M.H.G. Tytgat, Dark matter from WIMP to FIMP over three regimes: cosmology versus colliders. in *Strong Interactions from QCD to New Strong Dynamics at LHC and Future Colliders*, vol. 12 (2022), pp. 22– 27. arXiv:2212.12377
- J. Harz, K. Petraki, Radiative bound-state formation in unbroken perturbative non-Abelian theories and implications for dark matter. JHEP Ser. 07, 096 (2018). https://doi.org/10.1007/ JHEP07(2018)096. arXiv:1805.01200
- M. Becker, E. Copello, J. Harz, K.A. Mohan, D. Sengupta, Impact of Sommerfeld effect and bound state formation in simplified tchannel dark matter models. JHEP Ser. 08, 145 (2022). https:// doi.org/10.1007/JHEP08(2022)145. arXiv:2203.04326

- J. Harz, K. Petraki, Higgs enhancement for the dark matter relic density. Phys. Rev. D Ser. 97, 075041 (2018). https://doi.org/10. 1103/PhysRevD.97.075041. arXiv:1711.03552
- J. Harz, K. Petraki, Higgs-mediated bound states in dark-matter models. JHEP Ser. 04, 130 (2019). https://doi.org/10.1007/ JHEP04(2019)130. arXiv:1901.10030
- F. Elahi, C. Kolda, J. Unwin, UltraViolet freeze-in. JHEP Ser. 03, 048 (2015). https://doi.org/10.1007/JHEP03(2015)048. arXiv:1410.6157
- M. Becker, M.J.F. Lozano, J. Harz, C. Tamarit, Multiple soft scatterings in scalar dark matter freeze-in. arXiv:2506.11185
- J. Hisano, S. Matsumoto, M.M. Nojiri, Explosive dark matter annihilation. Phys. Rev. Lett. Ser. 92, 031303 (2004). https://doi. org/10.1103/PhysRevLett.92.031303. arXiv:hep-ph/0307216
- 308. J. Hisano, S. Matsumoto, M.M. Nojiri, O. Saito, Non-perturbative effect on dark matter annihilation and gamma ray signature from galactic center. Phys. Rev. D Ser. 71, 063528 (2005). https://doi. org/10.1103/PhysRevD.71.063528. arXiv:hep-ph/0412403
- M. Beneke, C. Hellmann, P. Ruiz-Femenia, Non-relativistic pair annihilation of nearly mass degenerate neutralinos and charginos I. General framework and S-wave annihilation. JHEP Ser. 03, 148 (2013). https://doi.org/10.1007/JHEP03(2013)148. arXiv:1210.7928
- K. Petraki, M. Postma, M. Wiechers, Dark-matter bound states from Feynman diagrams. JHEP Ser. 06, 128 (2015). https://doi. org/10.1007/JHEP06(2015)128. arXiv:1505.00109
- S. Cassel, Sommerfeld factor for arbitrary partial wave processes. J. Phys. G Ser. 37, 105009 (2010). https://doi.org/10.1088/0954-3899/37/10/105009. arXiv:0903.5307
- R. Iengo, Sommerfeld enhancement: general results from field theory diagrams. JHEP Ser. 05, 024 (2009). https://doi.org/10. 1088/1126-6708/2009/05/024. arXiv:0902.0688
- J. Ellis, F. Luo, K.A. Olive, Gluino coannihilation revisited. JHEP Ser. 09, 127 (2015). https://doi.org/10.1007/JHEP09(2015)127. arXiv:1503.07142
- 314. T. Binder, A. Filimonova, K. Petraki, G. White, Saha equilibrium for metastable bound states and dark matter freeze-out. Phys. Lett. B Ser. 833, 137323 (2022). https://doi.org/10.1016/j.physletb.2022.137323. arXiv:2112.00042
- M.M. Flores, K. Petraki, Unitarity in the non-relativistic regime and implications for dark matter. Phys. Lett. B Ser. 858, 139022 (2024). https://doi.org/10.1016/j.physletb.2024.139022. arXiv:2405.02222
- M. Beneke, T. Binder, L. de Ros, M. Garny, S. Lederer, Perturbative unitarity violation in radiative capture transitions to dark matter bound states. arXiv:2411.08737
- K. Petraki, A. Socha, C. Vasilaki, The role of unitarisation on darkmatter freeze-out via metastable bound states. arXiv:2505.20443
- A. Mitridate, M. Redi, J. Smirnov, A. Strumia, Cosmological implications of dark matter bound states. JCAP Ser. 05, 006 (2017). https://doi.org/10.1088/1475-7516/2017/05/006. arXiv:1702.01141
- M. Kawasaki, K. Kohri, N. Sugiyama, MeV scale reheating temperature and thermalization of neutrino background. Phys. Rev. D Ser. 62, 023506 (2000). https://doi.org/10.1103/PhysRevD.62.023506. arXiv:astro-ph/0002127
- T. Moroi, L. Randall, Wino cold dark matter from anomaly mediated SUSY breaking. Nucl. Phys. B Ser. 570, 455 (2000). https://doi.org/10.1016/S0550-3213(99)00748-8. arXiv:hep-ph/9906527
- G.D. Coughlan, W. Fischler, E.W. Kolb, S. Raby, G.G. Ross, Cosmological problems for the Polonyi potential. Phys. Lett. B Ser. 131, 59 (1983). https://doi.org/10.1016/0370-2693(83)91091-2
- J.D. Barrow, Massive particles as a probe of the early universe.
 Nucl. Phys. B Ser. 208, 501 (1982). https://doi.org/10.1016/ 0550-3213(82)90233-4



Eur. Phys. J. C (2025) 85:975 Page 93 of 96 975

 L.H. Ford, Gravitational particle creation and inflation. Phys. Rev. D Ser. 35, 2955 (1987). https://doi.org/10.1103/PhysRevD.35. 2955

- 324. A.A. Starobinsky, A new type of isotropic cosmological models without singularity. Phys. Lett. B Ser. **91**, 99 (1980). https://doi.org/10.1016/0370-2693(80)90670-X
- R. Kallosh, A. Linde, Non-minimal inflationary attractors. JCAP Ser. 10, 033 (2013). https://doi.org/10.1088/1475-7516/2013/10/ 033. arXiv:1307.7938
- R. Kallosh, A. Linde, Universality class in conformal inflation.
 JCAP Ser. 07, 002 (2013). https://doi.org/10.1088/1475-7516/2013/07/002. arXiv:1306.5220
- PLANCK Collaboration, Planck 2018 results. X. Constraints on inflation. Astron. Astrophys. 641 A10 (2020). https://doi.org/10. 1051/0004-6361/201833887. arXiv:1807.06211
- J. Silva-Malpartida, N. Bernal, J. Jones-Pérez, R.A. Lineros, From WIMPs to FIMPs with low reheating temperatures. JCAP Ser. 09, 015 (2023). https://doi.org/10.1088/1475-7516/2023/09/015. arXiv:2306.14943
- J. Silva-Malpartida, N. Bernal, J. Jones-Pérez, R.A. Lineros, From WIMPs to FIMPs: impact of early matter domination. JCAP Ser. 03, 003 (2025). https://doi.org/10.1088/1475-7516/2025/03/003. arXiv:2408.08950
- T. Bringmann, J. Edsjö, DarkSUSY 6.3 Freeze-in, out-of-equilibrium freeze-out, cosmic-ray upscattering and further new features. PoS CompTools2021 038 (2022). https://doi.org/10.22323/1.409.0038. arXiv:2203.07439
- 331. GAMBIT Dark Matter Workgroup Collaboration, DarkBit: a GAMBIT module for computing dark matter observables and likelihoods. Eur. Phys. J. C Ser. 77, 831 (2017). https://doi.org/ 10.1140/epjc/s10052-017-5155-4. arXiv:1705.07920
- T. Bringmann, L. Bergstrom, J. Edsjo, New gamma-ray contributions to supersymmetric dark matter annihilation. JHEP Ser. 01, 049 (2008). https://doi.org/10.1088/1126-6708/2008/01/049.arXiv:0710.3169
- T. Bringmann, F. Calore, G. Vertongen, C. Weniger, On the relevance of sharp gamma-ray features for indirect dark matter searches. Phys. Rev. D Ser. 84, 103525 (2011). https://doi.org/10.1103/PhysRevD.84.103525. arXiv:1106.1874
- 334. V. Barger, W.-Y. Keung, D. Marfatia, Bremsstrahlung in dark matter annihilation. Phys. Lett. B Ser. **707**, 385 (2012). https://doi.org/10.1016/j.physletb.2012.01.001. arXiv:1111.4523
- 335. T. Toma, Internal Bremsstrahlung signature of real scalar dark matter and consistency with thermal relic density. Phys. Rev. Lett. Ser. 111, 091301 (2013). https://doi.org/10.1103/PhysRevLett. 111.091301. arXiv:1307.6181
- M. Garny, A. Ibarra, S. Rydbeck, S. Vogl, Majorana dark matter with a coloured mediator: collider vs direct and indirect searches. JHEP Ser. 06, 169 (2014). https://doi.org/10.1007/JHEP06(2014)169. arXiv:1403.4634
- F. Giacchino, L. Lopez-Honorez, M.H.G. Tytgat, Scalar dark matter models with significant internal Bremsstrahlung. JCAP Ser. 10, 025 (2013). https://doi.org/10.1088/1475-7516/2013/10/025. arXiv:1307.6480
- 338. F. Giacchino, A. Ibarra, L. Lopez Honorez, M.H.G. Tytgat, S. Wild, Signatures from scalar dark matter with a vector-like quark mediator. JCAP 02, 002 (2016). https://doi.org/10.1088/ 1475-7516/2016/02/002. arXiv:1511.04452
- M. Garny, A. Ibarra, S. Vogl, Signatures of Majorana dark matter with t-channel mediators. Int. J. Mod. Phys. D Ser. 24, 1530019 (2015). https://doi.org/10.1142/S0218271815300190. arXiv:1503.01500
- A. Ibarra, A. Pierce, N.R. Shah, S. Vogl, Anatomy of coannihilation with a scalar top partner. Phys. Rev. D Ser. 91, 095018 (2015). https://doi.org/10.1103/PhysRevD.91.095018. arXiv:1501.03164

- S. Colucci, F. Giacchino, M.H.G. Tytgat, J. Vandecasteele, Radiative corrections to vectorlike portal dark matter. Phys. Rev. D Ser. 98, 115029 (2018). https://doi.org/10.1103/PhysRevD.98. 115029. arXiv:1805.10173
- 342. A. Bouquet, P. Salati, J. Silk, γ-ray lines as a probe for a cold dark matter halo. Phys. Rev. D Ser. **40**, 3168 (1989). https://doi.org/10.1103/PhysRevD.40.3168
- L. Bergstrom, Radiative processes in dark matter photino annihilation. Phys. Lett. B Ser. 225, 372 (1989). https://doi.org/10.1016/0370-2693(89)90585-6
- 344. S. Rudaz, On the annihilation of heavy neutral fermion pairs into monochromatic gamma-rays and its astrophysical implications. Phys. Rev. D Ser. 39, 3549 (1989). https://doi.org/10.1103/ PhysRevD.39.3549
- 345. L. Bergstrom, P. Ullio, Full one loop calculation of neutralino annihilation into two photons. Nucl. Phys. B Ser. **504**, 27 (1997). https://doi.org/10.1016/S0550-3213(97)00530-0. arXiv:hep-ph/9706232
- Z. Bern, P. Gondolo, M. Perelstein, Neutralino annihilation into two photons. Phys. Lett. B Ser. 411, 86 (1997). https://doi.org/10. 1016/S0370-2693(97)00990-8. arXiv:hep-ph/9706538
- 347. G. Bertone, C.B. Jackson, G. Shaughnessy, T.M.P. Tait, A. Vallinotto, The WIMP Forest: indirect detection of a chiral square. Phys. Rev. D Ser. **80**, 023512 (2009). https://doi.org/10.1103/PhysRevD.80.023512. arXiv:0904.1442
- 348. FERMI- LAT, DES Collaboration, Searching for dark matter annihilation in recently discovered Milky Way satellites with Fermi-LAT. Astrophys. J. 834 110 (2017). https://doi.org/10. 3847/1538-4357/834/2/110. arXiv:1611.03184
- 349. A. McDaniel, M. Ajello, C.M. Karwin, M. Di Mauro, A. Drlica-Wagner, M.A. Sánchez-Conde, Legacy analysis of dark matter annihilation from the Milky Way dwarf spheroidal galaxies with 14 years of Fermi-LAT data. Phys. Rev. D Ser. 109, 063024 (2024). https://doi.org/10.1103/PhysRevD.109.063024. arXiv:2311.04982
- A. Cuoco, J. Heisig, M. Korsmeier, M. Krämer, Constraining heavy dark matter with cosmic-ray antiprotons. JCAP Ser. 04, 004 (2018). https://doi.org/10.1088/1475-7516/2018/04/004. arXiv:1711.05274
- F. Calore, M. Cirelli, L. Derome, Y. Genolini, D. Maurin, P. Salati et al., AMS-02 antiprotons and dark matter: trimmed hints and robust bounds. SciPost Phys. 12, 163 (2022). https://doi.org/10.21468/SciPostPhys.12.5.163. arXiv:2202.03076
- 352. P. De la Torre Luque, M.W. Winkler, T. Linden, Antiproton bounds on dark matter annihilation from a combined analysis using the DRAGON2 code. JCAP 05 104 (2024). https://doi.org/10.1088/ 1475-7516/2024/05/104. arXiv:2401.10329
- 353. Fermi-LAT Collaboration, Updated search for spectral lines from galactic dark matter interactions with pass 8 data from the Fermi Large Area Telescope. Phys. Rev. D Ser. 91, 122002 (2015). https://doi.org/10.1103/PhysRevD.91.122002. arXiv:1506.00013
- F.S. Queiroz, C.E. Yaguna, C. Weniger, Gamma-ray limits on neutrino lines. JCAP Ser. 05, 050 (2016). https://doi.org/10.1088/ 1475-7516/2016/05/050. arXiv:1602.05966
- 355. M. Garny, A. Ibarra, S. Vogl, Antiproton constraints on dark matter annihilations from internal electroweak bremsstrahlung. JCAP Ser. 07, 028 (2011). https://doi.org/10.1088/1475-7516/2011/07/028. arXiv:1105.5367
- ICECUBE- GEN2 Collaboration, IceCube-Gen2: the window to the extreme Universe. J. Phys. G 48 060501 (2021). https://doi.org/ 10.1088/1361-6471/abbd48. arXiv:2008.04323
- 357. KM3NeT Collaboration, Letter of intent for KM3NeT 2.0. J. Phys. G 43, 084001 (2016). https://doi.org/10.1088/0954-3899/43/8/084001. arXiv:1601.07459



975 Page 94 of 96 Eur. Phys. J. C (2025) 85:975

 C.A. Argüelles, A. Diaz, A. Kheirandish, A. Olivares-Del-Campo,
 I. Safa, A.C. Vincent, Dark matter annihilation to neutrinos.
 Rev. Mod. Phys. Ser. 93, 035007 (2021). https://doi.org/10.1103/ RevModPhys.93.035007. arXiv:1912.09486

- 359. S. Basegmez Du Pree, C. Arina, A. Cheek, A. Dekker, M. Chianese, S. Ando, Robust limits from upcoming neutrino telescopes and implications on minimal dark matter models. JCAP Ser. 05, 054 (2021). https://doi.org/10.1088/1475-7516/2021/05/054.arXiv:2103.01237
- L.S. Miranda, S. Basegmez du Pree, K.C.Y. Ng, A. Cheek, C. Arina, Towards detecting super-GeV dark matter via annihilation to neutrinos. JCAP Ser. 08, 006 (2023). https://doi.org/10.1088/1475-7516/2023/08/006. arXiv:2211.12235
- 361. H.E.S.S. Collaboration, Search for photon-linelike signatures from dark matter annihilations with H.E.S.S. Phys. Rev. Lett. 110 041301 (2013). https://doi.org/10.1103/PhysRevLett.110. 041301. arXiv:1301.1173
- 362. H.E.S.S. Collaboration, H.E.S.S. limits on Linelike dark matter signatures in the 100 GeV to 2 TeV energy range close to the galactic center. Phys. Rev. Lett. 117, 151302 (2016). https://doi. org/10.1103/PhysRevLett.117.151302. arXiv:1609.08091
- 363. MAGIC Collaboration, Search for gamma-ray spectral lines from dark matter annihilation up to 100 TeV toward the galactic center with MAGIC. Phys. Rev. Lett. Ser. 130, 061002 (2023). https:// doi.org/10.1103/PhysRevLett.130.061002. arXiv:2212.10527
- G. Arcadi, D. Cabo-Almeida, F. Mescia, J. Virto, Dark matter direct detection in *t*-channel mediator models. JCAP Ser. 02, 005 (2024). https://doi.org/10.1088/1475-7516/2024/02/005. arXiv:2309.07896
- R.J. Hill, M.P. Solon, Standard Model anatomy of WIMP dark matter direct detection II: QCD analysis and hadronic matrix elements. Phys. Rev. D Ser. 91, 043505 (2015). https://doi.org/10. 1103/PhysRevD.91.043505. arXiv:1409.8290
- 366. K.A. Mohan, D. Sengupta, T.M.P. Tait, B. Yan, C.P. Yuan, Direct detection and LHC constraints on a *t*-channel simplified model of Majorana dark matter at one loop. JHEP Ser. 05, 115 (2019). https://doi.org/10.1007/JHEP05(2019)115. arXiv:1903.05650
- M. Drees, M. Nojiri, Neutralino nucleon scattering revisited. Phys. Rev. D Ser. 48, 3483 (1993). https://doi.org/10.1103/ PhysRevD.48.3483. arXiv:hep-ph/9307208
- C. Arina, A. Cheek, K. Mimasu, L. Pagani, Light and darkness: consistently coupling dark matter to photons via effective operators. Eur. Phys. J. C Ser. 81, 223 (2021). https://doi.org/10.1140/ epjc/s10052-021-09010-1. arXiv:2005.12789
- A. Kumar, S. Tulin, Top-flavored dark matter and the forward-backward asymmetry. Phys. Rev. D Ser. 87, 095006 (2013). https://doi.org/10.1103/PhysRevD.87.095006. arXiv:1303.0332
- 370. F. D'Eramo, B.J. Kavanagh, P. Panci, You can hide but you have to run: direct detection with vector mediators. JHEP Ser. 08, 111 (2016). https://doi.org/10.1007/JHEP08(2016)111. arXiv:1605.04917
- XENON Collaboration, Dark matter search results from a one ton-year exposure of XENON1T. Phys. Rev. Lett. Ser. 121, 111302 (2018). https://doi.org/10.1103/PhysRevLett.121. 111302. arXiv:1805.12562
- 372. XENON Collaboration, First dark matter search with nuclear recoils from the XENONnT Experiment. Phys. Rev. Lett. Ser. 131, 041003 (2023). https://doi.org/10.1103/PhysRevLett.131. 041003. arXiv:2303.14729
- 373. LZ Collaboration, First dark matter search results from the LUX-ZEPLIN (LZ) Experiment. Phys. Rev. Lett. Ser. 131, 041002 (2023). https://doi.org/10.1103/PhysRevLett.131. 041002. arXiv:2207.03764
- 374. L. Baudis, DARWIN/XLZD: a future xenon observatory for dark matter and other rare interactions. Nucl. Phys. B Ser.

- **1003**, 116473 (2024). https://doi.org/10.1016/j.nuclphysb.2024. 116473. arXiv:2404.19524
- 375. J. Aalbers et al., A next-generation liquid xenon observatory for dark matter and neutrino physics. J. Phys. G Ser. 50, 013001 (2023). https://doi.org/10.1088/1361-6471/ac841a. arXiv:2203.02309
- J. Billard, L. Strigari, E. Figueroa-Feliciano, Implication of neutrino backgrounds on the reach of next generation dark matter direct detection experiments. Phys. Rev. D Ser. 89, 023524 (2014). https://doi.org/10.1103/PhysRevD.89.023524. arXiv:1307.5458
- 377. PICO Collaboration, Dark matter search results from the PICO-60 C₃F₈ bubble chamber. Phys. Rev. Lett. Ser. **118**, 251301 (2017). https://doi.org/10.1103/PhysRevLett.118. 251301, arXiv:1702.07666
- 378. PANDAX Collaboration, Dark matter search results from 1.54 Tonne. Year exposure of PandaX-4T. Phys. Rev. Lett. 134, 011805 (2025). https://doi.org/10.1103/PhysRevLett.134. 011805. arXiv:2408.00664
- LZ Collaboration, Dark matter search results from 4.2 Tonneyears of exposure of the LUX-ZEPLIN (LZ) Experiment. Phys. Rev. Lett. 135, 011802 (2025). https://doi.org/10.1103/ 4dyc-z8zf. arXiv:2410.17036
- R. Essig, J. Mardon, T. Volansky, Direct detection of sub-GeV dark matter. Phys. Rev. D Ser. 85, 076007 (2012). https://doi.org/ 10.1103/PhysRevD.85.076007. arXiv:1108.5383
- T. Lin, Dark matter models and direct detection. PoS 333, 009 (2019). https://doi.org/10.22323/1.333.0009. arXiv:1904.07915
- M. Battaglieri et al., US cosmic visions: new ideas in dark matter 2017: Community Report. in *U.S. Cosmic Visions: New Ideas in Dark Matter*, vol. 7 (2017). arXiv:1707.04591
- 383. SuperCDMS Collaboration, Constraints on low-mass, relic dark matter candidates from a surface-operated SuperCDMS single-charge sensitive detector. Phys. Rev. D Ser. 102, 091101 (2020). https://doi.org/10.1103/PhysRevD.102.091101. arXiv:2005.14067
- 384. DAMIC Collaboration, Constraints on light dark matter particles interacting with electrons from DAMIC at SNOLAB. Phys. Rev. Lett. Ser. 123, 181802 (2019). https://doi.org/10.1103/PhysRevLett.123.181802. arXiv:1907.12628
- 385. SENSEI Collaboration, SENSEI: direct-detection results on sub-GeV dark matter from a new skipper-CCD. Phys. Rev. Lett. Ser. 125, 171802 (2020). https://doi.org/10.1103/PhysRevLett. 125.171802. arXiv:2004.11378
- PandaX-II Collaboration, Search for light dark matter-electron scatterings in the PandaX-II Experiment. Phys. Rev. Lett. Ser. 126, 211803 (2021). https://doi.org/10.1103/PhysRevLett.126. 211803. arXiv:2101.07479
- DarkSide Collaboration, Search for dark matter particle interactions with electron final states with DarkSide-50. Phys. Rev. Lett. Ser. 130, 101002 (2023). https://doi.org/10.1103/PhysRevLett. 130.101002. arXiv:2207.11968
- 388. XENON Collaboration, Light dark matter search with ionization signals in XENON1T. Phys. Rev. Lett. Ser. 123, 251801 (2019). https://doi.org/10.1103/PhysRevLett.123.251801.arXiv:1907.11485
- 389. T.R. Slatyer, C.-L. Wu, General constraints on dark matter decay from the cosmic microwave background. Phys. Rev. D Ser. 95, 023010 (2017). https://doi.org/10.1103/PhysRevD.95. 023010. arXiv:1610.06933
- 390. M. Lucca, N. Schöneberg, D.C. Hooper, J. Lesgourgues, J. Chluba, The synergy between CMB spectral distortions and anisotropies. JCAP Ser. 02, 026 (2020). https://doi.org/10.1088/1475-7516/2020/02/026. arXiv:1910.04619
- D.J. Fixsen, E.S. Cheng, J.M. Gales, J.C. Mather, R.A. Shafer,
 E.L. Wright, The cosmic microwave background spectrum from



Eur. Phys. J. C (2025) 85:975 Page 95 of 96 975

the full COBE FIRAS data set. Astrophys. J. Ser. **473**, 576 (1996). https://doi.org/10.1086/178173. arXiv:astro-ph/9605054

- J.C. Mather, D.J. Fixsen, R.A. Shafer, C. Mosier, D.T. Wilkinson, Calibrator design for the COBE far infrared absolute spectrophotometer (FIRAS). Astrophys. J. Ser. 512, 511 (1999). https://doi.org/10.1086/306805. arXiv:astro-ph/9810373
- 393. M. Hufnagel, K. Schmidt-Hoberg, S. Wild, BBN constraints on MeV-scale dark sectors Part I. Sterile decays. JCAP 02, 044 (2018). https://doi.org/10.1088/1475-7516/2018/02/044. arXiv:1712.03972
- 394. M. Kawasaki, K. Kohri, T. Moroi, Big-Bang nucleosynthesis and hadronic decay of long-lived massive particles. Phys. Rev. D Ser. 71, 083502 (2005). https://doi.org/10.1103/PhysRevD.71. 083502. arXiv:astro-ph/0408426
- 395. K. Jedamzik, Bounds on long-lived charged massive particles from Big Bang nucleosynthesis. JCAP Ser. 03, 008 (2008). https:// doi.org/10.1088/1475-7516/2008/03/008. arXiv:0710.5153
- 396. V. Iršič et al., New constraints on the free-streaming of warm dark matter from intermediate and small scale Lyman-α forest data. Phys. Rev. D Ser. **96**, 023522 (2017). https://doi.org/10.1103/PhysRevD.96.023522. arXiv:1702.01764
- A. Schneider, Astrophysical constraints on resonantly produced sterile neutrino dark matter. JCAP Ser. 04, 059 (2016). https://doi. org/10.1088/1475-7516/2016/04/059. arXiv:1601.07553
- R. Murgia, A. Merle, M. Viel, M. Totzauer, A. Schneider, "Non-cold" dark matter at small scales: a general approach. JCAP Ser. 11, 046 (2017). https://doi.org/10.1088/1475-7516/2017/11/046. arXiv:1704.07838
- 399. K.J. Bae, A. Kamada, S.P. Liew, K. Yanagi, Light axinos from freeze-in: production processes, phase space distributions, and Ly-α forest constraints. JCAP Ser. 01, 054 (2018). https://doi. org/10.1088/1475-7516/2018/01/054. arXiv:1707.06418
- P. Bode, J.P. Ostriker, N. Turok, Halo formation in warm dark matter models. Astrophys. J. Ser. 556, 93 (2001). https://doi.org/ 10.1086/321541. arXiv:astro-ph/0010389
- J. Heeck, D. Teresi, Cold keV dark matter from decays and scatterings. Phys. Rev. D Ser. 96, 035018 (2017). https://doi.org/10.1103/PhysRevD.96.035018. arXiv:1706.09909
- J. Lesgourgues, T. Tram, The Cosmic Linear Anisotropy Solving System (CLASS) IV: efficient implementation of non-cold relics. JCAP Ser. 09, 032 (2011). https://doi.org/10.1088/1475-7516/ 2011/09/032. arXiv:1104.2935
- 403. HESS Collaboration, Search for γ -ray line signals from dark matter annihilations in the inner galactic halo from 10 years of observations with H.E.S.S. Phys. Rev. Lett. **120** 201101 (2018). https://doi.org/10.1103/PhysRevLett.120.201101. arXiv:1805.05741
- LZ Collaboration, First dark matter search results from the LUX-ZEPLIN (LZ) Experiment. arXiv:2207.03764
- 405. CRESST Collaboration, First results from the CRESST-III low-mass dark matter program. Phys. Rev. D Ser. 100, 102002 (2019). https://doi.org/10.1103/PhysRevD.100.102002. arXiv:1904.00498
- 406. DarkSide Collaboration, Low-mass dark matter search with the DarkSide-50 Experiment. Phys. Rev. Lett. Ser. 121, 081307 (2018). https://doi.org/10.1103/PhysRevLett.121.081307. arXiv:1802.06994
- 407. PARTICLE DATA GROUP Collaboration, Review of particle physics. PTEP 2020 083C01 (2020). https://doi.org/10.1093/ ptep/ptaa104
- 408. XENON Collaboration, First dark matter search results from the XENON1T Experiment. Phys. Rev. Lett. Ser. 119, 181301 (2017). https://doi.org/10.1103/PhysRevLett.119. 181301. arXiv:1705.06655
- Fermi-LAT Collaboration, Sensitivity projections for dark matter searches with the Fermi Large Area Telescope. Phys. Rep.

- Ser. **636**, 1 (2016). https://doi.org/10.1016/j.physrep.2016.05. 001. arXiv:1605.02016
- N. Nagata, S. Shirai, Higgsino dark matter in high-scale supersymmetry. JHEP Ser. 01, 029 (2015). https://doi.org/10.1007/ JHEP01(2015)029. arXiv:1410.4549
- 411. C. Dessert, J.W. Foster, Y. Park, B.R. Safdi, W.L. Xu, Higgsino dark matter confronts 14 years of Fermi γ-ray data. Phys. Rev. Lett. Ser. 130, 201001 (2023). https://doi.org/10.1103/PhysRevLett.130.201001. arXiv:2207.10090
- 412. C. Arina, V. Martin-Lozano, G. Nardini, Dark matter versus $h \rightarrow \gamma \gamma$ and $h \rightarrow \gamma Z$ with supersymmetric triplets. JHEP Ser. **08**, 015 (2014). https://doi.org/10.1007/JHEP08(2014)015.
- 413. M. Cirelli, F. Sala, M. Taoso, Wino-like minimal dark matter and future colliders. JHEP Ser. 10, 033 (2014). https://doi.org/ 10.1007/JHEP01(2015)041. arXiv:1407.7058
- 414. M. Beneke, R. Szafron, K. Urban, Wino potential and Sommerfeld effect at NLO. Phys. Lett. B Ser. 800, 135112 (2020). https://doi. org/10.1016/j.physletb.2019.135112. arXiv:1909.04584
- 415. D. Agin, B. Fuks, M.D. Goodsell, T. Murphy, Monojets reveal overlapping excesses for light compressed Higgsinos. Phys. Lett. B Ser. 853, 138597 (2024). https://doi.org/10.1016/j.physletb. 2024.138597. arXiv:2311.17149
- D. Agin, B. Fuks, M.D. Goodsell, T. Murphy, Seeking a coherent explanation of LHC excesses for compressed spectra. Eur. Phys. J. C Ser. 84, 1218 (2024). https://doi.org/10.1140/epjc/s10052-024-13594-9. arXiv:2404.12423
- 417. M. Chakraborti, S. Heinemeyer, I. Saha, Consistent excesses in the search for $\tilde{\chi}_2^0 \tilde{\chi}_1^{\pm}$: wino/bino vs. Higgsino dark matter. Eur. Phys. J. C **84**, 812 (2024). https://doi.org/10.1140/epjc/s10052-024-13180-z. arXiv:2403.14759
- 418. S.P. Martin, The curtain lowers on directly detectable Higgsino dark matter. arXiv:2412.08958
- 419. Y. Bai, J. Berger, Lepton portal dark matter. JHEP Ser. 08, 153 (2014). https://doi.org/10.1007/JHEP08(2014)153. arXiv:1402.6696
- S. Chang, R. Edezhath, J. Hutchinson, M. Luty, Leptophilic effective WIMPs. Phys. Rev. D Ser. 90, 015011 (2014). https://doi.org/10.1103/PhysRevD.90.015011. arXiv:1402.7358
- P. Agrawal, Z. Chacko, C.B. Verhaaren, Leptophilic dark matter and the anomalous magnetic moment of the muon. JHEP Ser. 08, 147 (2014). https://doi.org/10.1007/JHEP08(2014)147. arXiv:1402.7369
- D. Barducci, A. Deandrea, S. Moretti, L. Panizzi, H. Prager, Characterizing dark matter interacting with extra charged leptons. Phys. Rev. D Ser. 97, 075006 (2018). https://doi.org/10. 1103/PhysRevD.97.075006. arXiv:1801.02707
- 423. J. Liu, X.-P. Wang, K.-P. Xie, Searching for lepton portal dark matter with colliders and gravitational waves. JHEP Ser. 06, 149 (2021). https://doi.org/10.1007/JHEP06(2021)149. arXiv:2104.06421
- 424. S. Biondini, P. Schicho, T.V.I. Tenkanen, Strong electroweak phase transition in t-channel simplified dark matter models. JCAP Ser. 10, 044 (2022). https://doi.org/10.1088/1475-7516/2022/10/ 044. arXiv:2207.12207
- 425. A. Jueid, S. Nasri, Lepton portal dark matter at muon colliders: total rates and generic features for phenomenologically viable scenarios. Phys. Rev. D Ser. 107, 115027 (2023). https://doi.org/10.1103/PhysRevD.107.115027. arXiv:2301.12524
- 426. A. Jueid, S. Nasri, R. Soualah, Searching for GeV-scale Majorana dark matter: inter spem et metum. JHEP Ser. 04, 012 (2021). https://doi.org/10.1007/JHEP04(2021)012. arXiv:2006.01348
- S.-I. Horigome, T. Katayose, S. Matsumoto, I. Saha, Leptophilic fermion WIMP: role of future lepton colliders. Phys. Rev. D Ser. 104, 055001 (2021). https://doi.org/10.1103/PhysRevD.104. 055001. arXiv:2102.08645



975 Page 96 of 96 Eur. Phys. J. C (2025) 85:975

428. J. Liu, B. Shuve, N. Weiner, I. Yavin, Looking for new charged states at the LHC: signatures of magnetic and Rayleigh dark matter. JHEP Ser. 07, 144 (2013). https://doi.org/10.1007/ JHEP07(2013)144. arXiv:1303.4404

- 429. M. Cermeño, C. Degrande, L. Mantani, Signatures of leptophilic t-channel dark matter from active galactic nuclei. Phys. Rev. D Ser. 105, 083019 (2022). https://doi.org/10.1103/PhysRevD.105. 083019. arXiv:2201.07247
- J.L. Feng, J. Kumar, D. Sanford, Xenophobic dark matter. Phys. Rev. D Ser. 88, 015021 (2013). https://doi.org/10.1103/ PhysRevD.88.015021. arXiv:1306.2315
- 431. G. Belanger, A. Mjallal, A. Pukhov, Recasting direct detection limits within micrOMEGAs and implication for non-standard dark matter scenarios. Eur. Phys. J. C Ser. 81, 239 (2021). https://doi.org/10.1140/epjc/s10052-021-09012-z. arXiv:2003.08621
- 432. MUON G-2 Collaboration, Measurement of the positive muon anomalous magnetic moment to 0.46 ppm. Phys. Rev. Lett. 126, 141801 (2021). https://doi.org/10.1103/PhysRevLett.126. 141801. arXiv:2104.03281
- 433. M. Aicheler, P. Burrows, M. Draper, T. Garvey, P. Lebrun, K. Peach et al. (eds.), A multi-TeV linear collider based on CLIC Technology: CLIC Conceptual Design Report
- ILC Collaboration, H. Baer et al. (eds.), The International Linear Collider Technical Design Report Volume 2: Physics. arXiv:1306.6352
- 435. F. An et al., Precision Higgs physics at the CEPC. Chin. Phys. C Ser. 43, 043002 (2019). https://doi.org/10.1088/1674-1137/43/4/ 043002. arXiv:1810.09037
- FCC Collaboration, FCC-ee: the lepton collider: future circular collider conceptual design report volume 2. Eur. Phys. J. ST Ser. 228, 261 (2019). https://doi.org/10.1140/epjst/e2019-900045-4
- 437. ATLAS Collaboration, Search for new phenomena in final states with an energetic jet and large missing transverse momentum in pp collisions at $\sqrt{s}=13$ TeV using the ATLAS detector. Phys. Rev. D Ser. **94**, 032005 (2016). https://doi.org/10.1103/PhysRevD.94.032005. arXiv:1604.07773
- 438. ATLAS Collaboration, Search for dark matter and other new phenomena in events with an energetic jet and large missing transverse momentum using the ATLAS detector. JHEP Ser. 01, 126 (2018). https://doi.org/10.1007/JHEP01(2018)126. arXiv:1711.03301
- 439. ATLAS Collaboration, Search for new phenomena in events with an energetic jet and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector
- ATLAS Collaboration, Optimisation of the ATLAS b-tagging performance for the 2016 LHC Run

- 441. ATLAS Collaboration, Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton-proton collision data. JINST Ser. 14, P12006 (2019). https://doi.org/10.1088/1748-0221/14/12/P12006. arXiv:1908.00005
- 442. ATLAS Collaboration, Muon reconstruction performance of the ATLAS detector in proton-proton collision data at √s =13 TeV. Eur. Phys. J. C Ser. 76, 292 (2016). https://doi.org/10.1140/epjc/s10052-016-4120-y. arXiv:1603.05598
- 443. ATLAS Collaboration, Identification and energy calibration of hadronically decaying tau leptons with the ATLAS experiment in pp collisions at \sqrt{s} =8 TeV. Eur. Phys. J. C Ser. **75**, 303 (2015). https://doi.org/10.1140/epjc/s10052-015-3500-z. arXiv:1412.7086
- 444. ATLAS Collaboration, Reconstruction, energy calibration, and identification of hadronically decaying tau leptons in the ATLAS experiment for Run-2 of the LHC
- 445. T. Sjöstrand, S. Ask, J.R. Christiansen, R. Corke, P. Ilten et al., An introduction to PYTHIA 8.2. Comput. Phys. Commun. 191, 159 (2015). https://doi.org/10.1016/j.cpc.2015.01.024. arXiv:1410.3012
- 446. C. Duhr, B. Fuks, A superspace module for the FeynRules package. Comput. Phys. Commun. Ser. **182**, 2404 (2011). https://doi.org/10.1016/j.cpc.2011.06.009. arXiv:1102.4191

