

Symmetry of minimizers for some nonlocal variational problems

Orlando Lopes^a, Mihai Mariş^{b,*}

^a *IMEUSP, Rua do Matao, 1010, Caixa postal 66281, CEP 05315-970, Sao Paulo, SP, Brazil*

^b *Université de Franche-Comté, Département de Mathématiques UMR 6623, 16, Route de Gray, 25030 Besançon Cedex, France*

Received 25 May 2007; accepted 13 October 2007

Available online 26 November 2007

Communicated by J.-M. Coron

Abstract

We present a new approach to study the symmetry of minimizers for a large class of nonlocal variational problems. This approach which generalizes the Reflection method is based on the existence of some integral identities. We study the identities that lead to symmetry results, the functionals that can be considered and the function spaces that can be used. Then we use our method to prove the symmetry of minimizers for a class of variational problems involving the fractional powers of Laplacian, for the generalized Choquard functional and for the standing waves of the Davey–Stewartson equation.

© 2007 Elsevier Inc. All rights reserved.

Keywords: Symmetry of minimizers; Nonlocal functional; Minimization under constraints; Fractional powers of Laplacian; Choquard functional; Davey–Stewartson equation

1. Introduction

Many important partial differential equations arising in physics are Euler–Lagrange equations of variational problems. Among the solutions of these equations those that correspond to a minimum of the associated functional (e.g. the “energy”) subject to some constraint are of particular interest. For example in many situations the set of such solutions is orbitally stable (see [9]).

* Corresponding author.

E-mail addresses: olopes@ime.usp.br (O. Lopes), mihai.maris@univ-fcomte.fr (M. Mariş).

In this paper we address the general question of whether, or not, the fact that the underlying problem has some symmetries is reflected on the minimizers. Namely if a problem is invariant under the action of a group of transformations, is it true that the corresponding minimizers are also invariant under the action of this group (or, perhaps, a subgroup of it)? As it is shown in [14], this may not be the case.

A classical approach to radial symmetry of minimizers is Schwarz symmetrization (or spherical decreasing rearrangement, see [16]). For a nonnegative function $u \in H^1(\mathbf{R}^N)$ its symmetrization u^* is a radially-decreasing function from $\mathbf{R}^N$ into $\mathbf{R}$ which has the property that $\text{meas}(\{x \in \mathbf{R}^N \mid u(x) > \lambda\}) = \text{meas}(\{x \in \mathbf{R}^N \mid u^*(x) > \lambda\})$ for any $\lambda > 0$. It is well known that u^* satisfies (among others) the following properties:

$$\begin{aligned} \int_{\mathbf{R}^N} |\nabla u^*(x)|^2 dx &\leq \int_{\mathbf{R}^N} |\nabla u(x)|^2 dx \quad \text{and} \\ \int_{\mathbf{R}^N} F(u^*(x)) dx &= \int_{\mathbf{R}^N} F(u(x)) dx, \end{aligned} \quad (1.1)$$

where F is, say, a smooth function from $\mathbf{R}$ into itself such that $F(u) \in L^1(\mathbf{R}^N)$ (see [16]). As a simple application of symmetrization, consider the problem of minimizing

$$E(u) = \frac{1}{2} \int_{\mathbf{R}^N} |\nabla u(x)|^2 dx + \int_{\mathbf{R}^N} F(u(x)) dx$$

subject to the constraint

$$\int_{\mathbf{R}^N} G(u(x)) dx = \lambda,$$

where $F, G \in C^1(\mathbf{R}, \mathbf{R})$ have the property that $F(u), G(u) \in L^1(\mathbf{R}^N)$ whenever $u \in H^1(\mathbf{R}^N)$. If $u \in H^1(\mathbf{R}^N)$ is a nonnegative minimizer, then it follows from (1.1) that u^* also satisfies the constraint and $E(u^*) \leq E(u)$; therefore, u^* is also a minimizer. To show that $u \equiv u^*$ except for translation is a more delicate question and this follows from a result in [6] and the Unique Continuation Principle.

In the case of vector-valued minimizers $u: \mathbf{R}^N \rightarrow \mathbf{R}^k$, symmetrization can also be used provided that each component of the minimizer is nonnegative, the function $F: \mathbf{R}^k \rightarrow \mathbf{R}$ satisfies a cooperative condition $F_{x_i x_j} \leq 0$ for $i \neq j$ and the constraint is of the form $\int_{\mathbf{R}^N} G_1(u_1) + G_2(u_2) + \cdots + G_k(u_k) dx = \text{constant}$. Notice that the function defining the constraint must have a special form because we want the value of the constraint to be preserved by symmetrization.

Another tool to prove radial symmetry of minimizers is the result by Gidas, Ni and Nirenberg [11] about the radial symmetry of positive solutions of the semilinear elliptic equation

$$-\Delta u + f(u) = 0.$$

In the case of systems, an extension of that result has been proved in [7,25] assuming a cooperative condition for the nonlinearity. In [11] as well as in its generalizations the nonlinearities are also allowed to depend on the space variable in a radial and monotonic way.

As we can see, in the vector case, besides the need to know in advance that the components of the minimizer are positive, both methods described above require the nonlinearity to satisfy a cooperative condition and the function defining the constraint to have a special form. To avoid these two restrictions, the Reflection method has been developed in [18,19]. We now briefly describe this method.

Consider the problem of minimizing

$$E(u, v) = \frac{1}{2} \int_{\mathbf{R}^N} (|\nabla u(x)|^2 + |\nabla v(x)|^2) dx + \int_{\mathbf{R}^N} F(u(x), v(x)) dx$$

subject to

$$\int_{\mathbf{R}^N} G(u(x), v(x)) dx = \lambda \neq 0.$$

To show that any minimizer (u, v) is symmetric with respect to x_1 (except possibly for a translation), we first make a translation in the x_1 variable in such a way that

$$\int_{\{x_1 < 0\}} G(u(x), v(x)) dx = \int_{\{x_1 > 0\}} G(u(x), v(x)) dx = \frac{\lambda}{2}. \quad (1.2)$$

Next, setting $x = (x_1, x')$, where $x' \in \mathbf{R}^{N-1}$, we define the functions u_1 and u_2 by

$$u_1(x) = u_1(x_1, x') = \begin{cases} u(x_1, x') & \text{if } x_1 < 0, \\ u(-x_1, x') & \text{if } x_1 \geq 0 \end{cases} \quad \text{and} \\ u_2(x) = \begin{cases} u(-x_1, x') & \text{if } x_1 < 0, \\ u(x_1, x') & \text{if } x_1 \geq 0. \end{cases}$$

In a similar way we define v_1 and v_2 . According to (1.2), the pairs (u_1, v_1) and (u_2, v_2) also satisfy the constraint (i.e. they are admissible). Moreover, a change of variables shows that

$$E(u_1, v_1) + E(u_2, v_2) = 2E(u, v). \quad (1.3)$$

Thus (u_1, v_1) and (u_2, v_2) are also minimizers. This shows that there exist minimizers which are symmetric with respect to x_1 . In fact, by using the Euler–Lagrange equations and the Unique Continuation Principle we can show that necessarily $(u_1, v_1) = (u, v) = (u_2, v_2)$. Clearly, this implies that any minimizer (u, v) is symmetric with respect to the first variable. Replacing the x_1 -direction by any other direction in $\mathbf{R}^N$ and repeating the same argument, we can show that (u, v) is radially symmetric except for translation (details will be given later). Notice that to use this argument there is no need to know the sign of components of the minimizers.

The main point of this paper is to extend the Reflection method to a class of nonlocal functionals. To be more specific, consider the problem of minimizing

$$E(u) = \int_{\mathbf{R}^N} m(\xi) |\widehat{u}(\xi)|^2 d\xi + \int_{\mathbf{R}^N} F(u) dx \quad (1.4)$$

subject to the constraint

$$Q(u) = \int_{\mathbf{R}^N} G(u) dx = \lambda \neq 0. \quad (1.5)$$

Defining $W(u) = \int_{\mathbf{R}^N} m(\xi) |\widehat{u}(\xi)|^2 d\xi$ and u_1, u_2 as above, instead of (1.3) we have

$$E(u_1) + E(u_2) - 2E(u) = W(u_1) + W(u_2) - 2W(u).$$

Therefore, to show that u_1 and u_2 are also minimizers we need to know that

$$W(u_1) + W(u_2) - 2W(u) \leq 0. \quad (1.6)$$

The key to the method developed here is to show that inequality (1.6) holds true (see Theorem 2.7). In this article we will use this extended Reflection method to prove the symmetry of all minimizers of the following functionals:

- the Hamiltonian of a coupled system between a multidimensional Korteweg–de Vries equation and a Benjamin–Ono equation. Here minimizers correspond to solitary waves;
- the generalized Choquard functional. In this case the minimizers give rise to standing waves for the generalized Hartree equation;
- the Hamiltonian of the generalized Davey–Stewartson equation. Here again, minimizers correspond to standing waves.

The existence of minimizers for these problems can be proved by using the concentration–compactness method [17] or the alternative method presented in [20] and will not be discussed here.

Notice that the symmetrization approach, in general, does not apply to the problems above. Indeed, in the first two examples, symmetrization cannot be used to prove the existence of a radially symmetric minimizer under the general assumptions on the nonlinearities made in this paper. Furthermore, with the tools available at the present time, it is not clear how to prove the radial symmetry of *all* minimizers, even in the cases where symmetrization can be used to prove the existence of a radially symmetric minimizer. Finally, in the last example, symmetrization cannot be used because one term of the Hamiltonian of the Davey–Stewartson equation is a singular integral operator whose kernel changes sign.

This paper is organized as follows: in the next section we present some integral identities for functionals of the form $W(u) = \int_{\mathbf{R}^N} m(\xi) |\widehat{u}(\xi)|^2 d\xi$. These identities are first proved for functions $u \in C_c^\infty$ and are crucial for our approach to symmetry. It will also appear clearly what kind of symbols $m(\xi)$ we may consider. In Section 3 we search for appropriate function spaces on which our method can be applied. It will be proved that we may work on $H^s(\mathbf{R}^N)$ or on $\dot{H}^s(\mathbf{R}^N)$ if $-\frac{1}{2} < s < \frac{3}{2}$. We will extend the integral identities obtained in Section 2 to these function spaces. In Section 4 we apply our results to the concrete problems presented above. We end this article with some open problems.

2. Some identities

In what follows, $x = (x_1, x_2, \dots, x_N) = (x_1, x')$ denotes a point of $\mathbf{R}^N$, $x' = (x_2, \dots, x_N) \in \mathbf{R}^{N-1}$, $\xi = (\xi_1, \xi_2, \dots, \xi_N) = (\xi_1, \xi') \in \mathbf{R}^N$ with $\xi' = (\xi_2, \dots, \xi_N) \in \mathbf{R}^{N-1}$. We denote the Fourier transform either by $\widehat{\cdot}$ or by $\mathcal{F}$.

The aim of this section is to prove an identity for some functionals of the type

$$W(u) = \int_{\mathbf{R}^N} m(\xi) |\widehat{u}(\xi)|^2 d\xi \quad (2.1)$$

which play a very important role in proving symmetries.

Consider a function $u \in C_c^\infty(\mathbf{R}^N)$. We define the reflected functions u_1 and u_2 as follows:

$$\begin{aligned} u_1(x) = u_1(x_1, x') &= \begin{cases} u(x_1, x') & \text{if } x_1 < 0, \\ u(-x_1, x') & \text{if } x_1 \geq 0 \end{cases} \quad \text{and} \\ u_2(x) &= \begin{cases} u(-x_1, x') & \text{if } x_1 < 0, \\ u(x_1, x') & \text{if } x_1 \geq 0. \end{cases} \end{aligned} \quad (2.2)$$

We also define

$$g(x) = \frac{1}{2}(u(x_1, x') + u(-x_1, x')) \quad \text{and} \quad f(x) = \frac{1}{2}(u(x_1, x') - u(-x_1, x')). \quad (2.3)$$

Clearly, $f, g \in C_c^\infty(\mathbf{R}^N)$, g is even and f is odd with respect to x_1 and $u = f + g$. Let

$$f_*(x) = \begin{cases} f(-x_1, x') = -f(x) & \text{if } x_1 < 0, \\ f(x_1, x') & \text{if } x_1 \geq 0. \end{cases} \quad (2.4)$$

Then f_* is even with respect to x_1 , $u_1 = g - f_*$ and $u_2 = g + f_*$.

We want to study the quantity $W(u_1) + W(u_2) - 2W(u)$, where W is given by (2.1). Later in Theorem 2.7 we specify the class of multipliers under consideration but, at this early stage, besides integrability conditions, we assume that

$$m(\xi) \text{ is real} \quad \text{and} \quad m(-\xi_1, \xi') = m(\xi_1, \xi'). \quad (2.5)$$

It is easy to see that

$$\widehat{g}(-\xi_1, \xi') = \widehat{g}(\xi_1, \xi') \quad \text{and} \quad \widehat{f}(-\xi_1, \xi') = -\widehat{f}(\xi_1, \xi'). \quad (2.6)$$

Therefore

$$\begin{aligned} &W(u_1) + W(u_2) - 2W(u) \\ &= \int_{\mathbf{R}^N} m(\xi_1, \xi') (|\widehat{g}(\xi) - \widehat{f}_*(\xi)|^2 + |\widehat{g}(\xi) + \widehat{f}_*(\xi)|^2 - 2|\widehat{g}(\xi) + \widehat{f}(\xi)|^2) d\xi \end{aligned}$$

$$\begin{aligned}
&= \int_{\mathbf{R}^N} m(\xi_1, \xi') (2|\widehat{f}_*(\xi)|^2 - 2|\widehat{f}(\xi)|^2 - 4\operatorname{Re}(\widehat{g}(\xi)\overline{\widehat{f}(\xi)})) d\xi \\
&= 2 \int_{\mathbf{R}^N} m(\xi_1, \xi') (|\widehat{f}_*(\xi)|^2 - |\widehat{f}(\xi)|^2) d\xi = 2W(f_*) - 2W(f)
\end{aligned} \tag{2.7}$$

because $\int_{\mathbf{R}^N} m(\xi_1, \xi') \operatorname{Re}(\widehat{g}(\xi)\overline{\widehat{f}(\xi)}) d\xi = 0$ in view of (2.5) and (2.6).

It is obvious that

$$\begin{aligned}
\widehat{f}(\xi_1, \xi') &= \int_{\mathbf{R}} \int_{\mathbf{R}^{N-1}} e^{-ix_1\xi_1 - ix'\cdot\xi'} f(x_1, x') dx' dx_1 \\
&= \int_0^\infty \int_{\mathbf{R}^{N-1}} (e^{-ix_1\xi_1} - e^{ix_1\xi_1}) e^{-ix'\cdot\xi'} f(x_1, x') dx' dx_1 \\
&= -2i \int_0^\infty \int_{\mathbf{R}^{N-1}} \sin(x_1\xi_1) e^{-ix'\cdot\xi'} f(x_1, x') dx' dx_1
\end{aligned}$$

and similarly

$$\widehat{f}_*(\xi_1, \xi') = 2 \int_0^\infty \int_{\mathbf{R}^{N-1}} \cos(x_1\xi_1) e^{-ix'\cdot\xi'} f(x_1, x') dx' dx_1.$$

We denote by $\mathcal{F}_{N-1}$ the partial Fourier transform in the last $N-1$ variables, that is

$$\mathcal{F}_{N-1} f(x_1, \xi') = \int_{\mathbf{R}^{N-1}} e^{-ix'\cdot\xi'} f(x_1, x') dx'.$$

Since $f \in C_c^\infty(\mathbf{R}^N)$ we may use Fubini's theorem to get

$$\begin{aligned}
|\widehat{f}(\xi_1, \xi')|^2 &= \widehat{f}(\xi_1, \xi') \overline{\widehat{f}(\xi_1, \xi')} \\
&= 4 \int_0^\infty \int_0^\infty \sin(x_1\xi_1) \sin(y_1\xi_1) (\mathcal{F}_{N-1} f)(x_1, \xi') \overline{(\mathcal{F}_{N-1} f)(y_1, \xi')} dx_1 dy_1.
\end{aligned}$$

In the same way,

$$|\widehat{f}_*(\xi_1, \xi')|^2 = 4 \int_0^\infty \int_0^\infty \cos(x_1\xi_1) \cos(y_1\xi_1) (\mathcal{F}_{N-1} f)(x_1, \xi') \overline{(\mathcal{F}_{N-1} f)(y_1, \xi')} dx_1 dy_1.$$

Consequently,

$$\begin{aligned}
& W(f_*) - W(f) \\
&= 4 \int_{\mathbf{R}^N} m(\xi_1, \xi') \int_0^\infty \int_0^\infty [\cos(x_1 \xi_1) \cos(y_1 \xi_1) - \sin(x_1 \xi_1) \sin(y_1 \xi_1)] \\
&\quad \times (\mathcal{F}_{N-1} f)(x_1, \xi') \overline{(\mathcal{F}_{N-1} f)(y_1, \xi')} dx_1 dy_1 d\xi \\
&= 4 \int_{\mathbf{R}^N} m(\xi_1, \xi') \int_0^\infty \int_0^\infty \cos((x_1 + y_1) \xi_1) (\mathcal{F}_{N-1} f)(x_1, \xi') \overline{(\mathcal{F}_{N-1} f)(y_1, \xi')} dx_1 dy_1 d\xi. \quad (2.8)
\end{aligned}$$

For an arbitrary (but fixed) $\xi' \in \mathbf{R}^{N-1}$, we define $\varphi_{\xi'}(t) = (\mathcal{F}_{N-1} f)(t, \xi')$. Since $f \in C_c^\infty(\mathbf{R}^N)$, it is clear that $\varphi_{\xi'} \in C_c^\infty(\mathbf{R})$. If $\text{supp}(f) \subset B_{\mathbf{R}^N}(0, R)$, then $\text{supp}(\varphi_{\xi'}) \subset [-R, R]$. For $z \in \mathbf{C}$, we define

$$h_{\xi'}(z) = \int_0^\infty \int_0^\infty e^{i(x_1 + y_1)z} \varphi_{\xi'}(x_1) \overline{\varphi_{\xi'}(y_1)} dx_1 dy_1. \quad (2.9)$$

Since $\varphi_{\xi'}$ is bounded and has compact support, $h_{\xi'}$ is well defined and is an holomorphic function on $\mathbf{C}$. For any $z \in \mathbf{R}$ we have

$$\begin{aligned}
\overline{h_{\xi'}(z)} &= \int_0^\infty \int_0^\infty e^{-i(x_1 + y_1)z} \overline{\varphi_{\xi'}(x_1)} \varphi_{\xi'}(y_1) dx_1 dy_1 = h_{\xi'}(-z) \quad \text{and} \\
\text{Re}(h_{\xi'}(z)) &= \frac{1}{2} (h_{\xi'}(z) + \overline{h_{\xi'}(z)}) = \int_0^\infty \int_0^\infty \cos((x_1 + y_1)z) \varphi_{\xi'}(x_1) \overline{\varphi_{\xi'}(y_1)} dx_1 dy_1.
\end{aligned}$$

From (2.7) and (2.8) we get

$$W(u_1) + W(u_2) - 2W(u) = 2W(f_*) - 2W(f) = 8 \int_{\mathbf{R}^{N-1}} \int_{-\infty}^\infty m(\xi_1, \xi') h_{\xi'}(\xi_1) d\xi_1 d\xi'. \quad (2.10)$$

Some properties of the function $h_{\xi'}$ are given in the next lemma. To simplify the notation, we shall write h instead of $h_{\xi'}$.

Lemma 2.1. *For any fixed ξ' , the function $h = h_{\xi'}$ given by (2.9) has the following properties:*

- (i) h is bounded in the upper half-plane $\{z \in \mathbf{C} \mid \text{Im}(z) \geq 0\}$.
- (ii) There exists a constant $C > 0$ (depending on f and ξ') such that for any $z \neq 0$ with $\text{Im}(z) \geq 0$ we have:

$$|h(z)| \leq \frac{C}{|z|^4} \quad \text{and} \quad |h'(z)| \leq \frac{C}{|z|^5}. \quad (2.11)$$

Proof. (i) If $b \geq 0$ and $x \geq 0$ then $|e^{iax-bx}| \leq 1$ and we have

$$\begin{aligned} |h(a+ib)| &= \left| \int_0^\infty \int_0^\infty e^{i(x_1+y_1)a-(x_1+y_1)b} \varphi_{\xi'}(x_1) \overline{\varphi_{\xi'}(y_1)} dx_1 dy_1 \right| \\ &\leq \left(\int_0^\infty |e^{iat-bt}| \cdot |\varphi_{\xi'}(t)| dt \right)^2 \leq \left(\int_0^\infty |\varphi_{\xi'}(t)| dt \right)^2. \end{aligned}$$

(ii) It is clear that

$$h(z) = \int_0^\infty e^{ix_1 z} \varphi_{\xi'}(x_1) dx_1 \cdot \int_0^\infty e^{iy_1 z} \overline{\varphi_{\xi'}(y_1)} dy_1 = \Psi_1(z) \Psi_2(z), \quad (2.12)$$

where $\Psi_1(z)$ and $\Psi_2(z)$ are defined in an obvious way. Notice that $\varphi_{\xi'}(0) = (\mathcal{F}_{N-1}f)(0, \xi') = 0$ because $f(0, x') = 0$ (recall that f is odd with respect to x_1). Moreover, for any $k \in \mathbf{N}$,

$$\frac{d^k}{dt^k} \varphi_{\xi'}(t) = \int_{\mathbf{R}^{N-1}} e^{-ix' \xi'} \frac{\partial^k f}{\partial x_1^k}(t, x') dx' = \left(\mathcal{F}_{N-1} \frac{\partial^k f}{\partial x_1^k} \right)(t, \xi')$$

is a C_c^∞ function of t , uniformly bounded for $(t, \xi') \in \mathbf{R} \times \mathbf{R}^{N-1}$. Integrating by parts, we get:

$$\begin{aligned} \Psi_1(z) &= \int_0^\infty e^{itz} \varphi_{\xi'}(t) dt = \frac{1}{iz} e^{itz} \varphi_{\xi'}(t) \Big|_{t=0}^\infty - \frac{1}{iz} \int_0^\infty e^{itz} \varphi_{\xi'}'(t) dt \\ &= -\frac{e^{itz}}{(iz)^2} \varphi_{\xi'}'(t) \Big|_{t=0}^\infty + \frac{1}{(iz)^2} \int_0^\infty e^{itz} \varphi_{\xi'}''(t) dt = -\frac{1}{z^2} \left[\varphi_{\xi'}'(0) + \int_0^\infty e^{itz} \varphi_{\xi'}''(t) dt \right]. \end{aligned}$$

It is clear that an analogous estimate is true for $\Psi_2(z)$ and the first inequality in (2.11) holds.

Similarly one can prove that $|\psi_j'(z)| \leq \frac{C_j}{|z|^3}$ for $j = 1, 2$ and $\operatorname{Re}(z) \geq 0$. Since $h'(z) = \Psi_1'(z) \Psi_2(z) + \Psi_1(z) \Psi_2'(z)$, the second estimate in (2.11) follows. $\square$

Remark 2.2. In general, $\frac{\partial f}{\partial x_1}(0, x')$ does not vanish identically; hence $\mathcal{F}_{N-1}f(0, \xi') \neq 0$ for some ξ' , i.e. there exists ξ' such that $\varphi_{\xi'}'(0) \neq 0$. For such ξ' , the functions Ψ_1 and Ψ_2 do not decay faster than $\frac{1}{|z|^2}$ and the estimate (2.11) is optimal.

Remark 2.3. Note that for any $t \in \mathbf{R}$ we have $h(it) = \left| \int_0^\infty e^{-x_1 t} \varphi_{\xi'}(x_1) dx_1 \right|^2 \in [0, \infty)$. Suppose that for any fixed $\xi' \in \mathbf{R}^{N-1}$, $m(\xi_1, \xi')$ admits an holomorphic extension $z \mapsto m(z, \xi')$ to the upper half-plane $\{z \in \mathbf{C} \mid \operatorname{Im}(z) > 0\}$, with possibly some singularities on the imaginary axis $\{it \mid t \in [0, \infty)\}$. If $|m(z, \xi')|$ increases more slowly than $|z|^3$ as $|z| \rightarrow \infty$, then $\int_{-\infty}^\infty m(\xi_1, \xi') h(\xi_1) d\xi_1$ should depend only on the values of h on the singular set of $m(\cdot, \xi')$. This simple idea will enable us to prove the identities that will be crucial in symmetry problems.

In order to clarify what kind of symbols may be considered, we start with some auxiliary technical results about holomorphic functions in a half-plane and their boundary values.

Given a function $\alpha \in L^p(\mathbf{R})$, $1 \leq p < \infty$, we recall that its Hilbert transform is defined by

$$(H\alpha)(x) = \lim_{\varepsilon \rightarrow 0} \frac{1}{\pi} \int_{\{|y| > \varepsilon\}} \frac{\alpha(x-y)}{y} dy \quad \text{or equivalently} \quad \widehat{H\alpha}(\xi) = -i \operatorname{sgn}(\xi) \widehat{\alpha}(\xi).$$

It is well known that H is a bounded linear mapping from $L^p(\mathbf{R})$ into $L^p(\mathbf{R})$ (see, e.g., [23, Chapter II], or [24, inequality (2.11), p. 188]).

In the next two lemmas we collect some classical facts that will be very useful in the sequel. Proofs can be found in [24, Chapters I, II, VI] or in [23].

Lemma 2.4. Consider $\alpha \in L^p(\mathbf{R})$, $1 < p < \infty$, and let $\beta = H\alpha$. For $x > 0$ and $y \in \mathbf{R}$ define

$$a(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{x}{x^2 + (y-t)^2} \alpha(t) dt = \int_{-\infty}^{\infty} P(y-t, x) \alpha(t) dt \quad \text{and} \\ b(x, y) = -\frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y-t}{x^2 + (y-t)^2} \alpha(t) dt = -\int_{-\infty}^{\infty} Q(y-t, x) \alpha(t) dt,$$

where $P(s, k) = \frac{1}{\pi} \frac{k}{s^2 + k^2}$ and $Q(s, k) = \frac{1}{\pi} \frac{s}{s^2 + k^2}$ are the Poisson kernel, respectively the conjugate Poisson kernel.

Then we have:

- (i) $b(x, y) = -\int_{-\infty}^{\infty} P(y-t, x) \beta(t) dt$ for any $x > 0$ and $t \in \mathbf{R}$.
- (ii) $\|a(x, \cdot)\|_{L^p(\mathbf{R})} \leq \|\alpha\|_{L^p(\mathbf{R})}$, $\|b(x, \cdot)\|_{L^p(\mathbf{R})} \leq \|\beta\|_{L^p(\mathbf{R})}$ and $\|a(x, \cdot) - \alpha\|_{L^p(\mathbf{R})} \rightarrow 0$, $\|b(x, \cdot) + \beta\|_{L^p(\mathbf{R})} \rightarrow 0$ as $x \rightarrow 0$. Moreover, $a(x, y) \rightarrow \alpha(y)$ for any y in the Lebesgue set of α (hence almost everywhere) and $b(x, y) \rightarrow -\beta(y)$ for any y in the Lebesgue set of β .
- (iii) The functions a and b are harmonic in $\{(x, y) \in \mathbf{R}^2 \mid x > 0\}$ and $r(z) = r(x + iy) := a(x, y) + ib(x, y)$ is holomorphic in $\{z \in \mathbf{C} \mid \operatorname{Re}(z) > 0\}$.
- (iv) There exists a constant $A > 0$ such that

$$|a(x, y)| \leq \frac{A \|\alpha\|_{L^p}}{x^{\frac{1}{p}}} \quad \text{and} \quad |b(x, y)| \leq \frac{A \|\alpha\|_{L^p}}{x^{\frac{1}{p}}} \quad \text{for any } x > 0 \text{ and } y \in \mathbf{R}, \quad (2.13)$$

and for any $\delta > 0$ we have

$$\lim_{|(x, y)| \rightarrow \infty, x \geq \delta} a(x, y) = 0 \quad \text{and} \quad \lim_{|(x, y)| \rightarrow \infty, x \geq \delta} b(x, y) = 0.$$

Lemma 2.5. Let μ be a finite Borel measure on $\mathbf{R}$. For $x > 0$ and $y \in \mathbf{R}$ define

$$a(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{x}{x^2 + (y-t)^2} d\mu(t) = \int_{-\infty}^{\infty} P(y-t, x) d\mu(t) \quad \text{and}$$

$$b(x, y) = -\frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y-t}{x^2 + (y-t)^2} d\mu(t) = -\int_{-\infty}^{\infty} Q(y-t, x) d\mu(t),$$

where $P(s, k)$ and $Q(s, k)$ are the Poisson kernel, respectively the conjugate Poisson kernel.

Then:

- (i) The functions a and b are harmonic in $\{(x, y) \in \mathbf{R}^2 \mid x > 0\}$ and $r(z) = r(x + iy) := a(x, y) + ib(x, y)$ is holomorphic in the right half-plane $\{z \in \mathbf{C} \mid \operatorname{Re}(z) > 0\}$.
- (ii) For any $x > 0$ and any p , $1 \leq p \leq \infty$, we have

$$\|a(x, \cdot)\|_{L^p(\mathbf{R})} \leq \frac{1}{\pi^{\frac{1}{q}} x^{\frac{1}{q}}} \|\mu\|, \quad (2.14)$$

where q is the conjugate exponent of p and $\|\mu\|$ is the total variation of μ . Furthermore,

$$\lim_{x \rightarrow 0} \int_{\mathbf{R}} a(x, y) \phi(y) dy = \int_{\mathbf{R}} \phi(y) d\mu(y) \quad (2.15)$$

for any function ϕ which is continuous on $\mathbf{R}$ and tends to zero at $\pm\infty$.

- (iii) For any $x > 0$ we have $b(x, \cdot) = -Ha(x, \cdot)$ and $|b(x, y)| \leq \frac{1}{2\pi x} \|\mu\|$. Moreover, for any $p \in (1, \infty)$ there exists $A_p > 0$ such that

$$\|b(x, \cdot)\|_{L^p(\mathbf{R})} \leq A_p x^{-\frac{p-1}{p}} \|\mu\|.$$

- (iv) For any $\delta > 0$ we have $\lim_{|(x,y)| \rightarrow \infty, x \geq \delta} a(x, y) = 0$ and $\lim_{|(x,y)| \rightarrow \infty, x \geq \delta} b(x, y) = 0$.
- (v) Suppose in addition that $\mu(S) = \mu(-S)$ and $\mu(S \cap [-\varepsilon, \varepsilon]) = 0$ for any Borel measurable set S . Then a and b are well defined, bounded and holomorphic in the strip $\{(x, y) \in \mathbf{R}^2 \mid -\frac{\varepsilon}{2} < y < \frac{\varepsilon}{2}\}$, the function $r(x + iy) = a(x, y) + ib(x, y)$ is holomorphic in that strip and $r(0) = 0$.

After this preparation, we come back to the study of the integral $\int_{\mathbf{R}} m(\xi_1, \xi') h_{\xi'}(\xi_1) d\xi_1$ which appears in the right-hand side of (2.10).

Lemma 2.6. Suppose that for a given $\xi' \in \mathbf{R}^{N-1}$ the symbol $m(\xi_1, \xi')$ can be written as

$$\begin{aligned} m(\xi_1, \xi') &= A_0(\xi') + A_1(\xi')|\xi_1| + A_2(\xi')\xi_1^2 + \frac{1}{\pi} \xi_1^4 \int_{\mathbf{R}} \frac{1}{\xi_1^2 + t^2} \alpha_{\xi'}(t) dt \\ &+ \frac{1}{\pi} \left[\int_{\mathbf{R}} \frac{1}{\xi_1^2 + t^2} d\mu_{\xi',0}(t) + \xi_1^2 \int_{\mathbf{R}} \frac{1}{\xi_1^2 + t^2} d\mu_{\xi',1}(t) + \xi_1^4 \int_{\mathbf{R}} \frac{1}{\xi_1^2 + t^2} d\mu_{\xi',2}(t) \right], \end{aligned} \quad (2.16)$$

where:

- (a) $A_0(\xi'), A_1(\xi'), A_2(\xi') \in \mathbf{R}$,

- (b) $\alpha_{\xi'} \in L^p(\mathbf{R})$ for some $p \in (1, \infty)$ and $\alpha_{\xi'}$ is an even function,
 (c) $\mu_{\xi',i}$ are finite Borel measures on $\mathbf{R}$ such that $\mu_{\xi',i}(S) = \mu_{\xi',i}(-S)$ for any Borel measurable set $S \subset \mathbf{R}$, $i = 0, 1, 2$. Moreover, there exists $\eta > 0$ such that $\mu_{\xi',0}(S) = 0$ for any Borel measurable set $S \subset [-\eta, \eta]$.

Let $h = h_{\xi'}$ be given by (2.9). Then we have the identity:

$$\begin{aligned} \frac{1}{2} \int_{-\infty}^{\infty} m(\xi_1, \xi') h(\xi_1) d\xi_1 &= -A_1(\xi') \int_0^{\infty} t h(it) dt + \int_0^{\infty} t^3 \alpha_{\xi'}(t) h(it) dt + \int_0^{\infty} \frac{h(it)}{t} d\mu_{\xi',0}(t) \\ &\quad - \int_0^{\infty} t h(it) d\mu_{\xi',1}(t) + \int_0^{\infty} t^3 h(it) d\mu_{\xi',2}(t). \end{aligned} \quad (2.17)$$

Proof. For $z = x + iy \in \mathbf{C}$ with $\operatorname{Re}(z) > 0$ we define

$$\begin{aligned} r(z) &= \frac{1}{\pi} \int_{\mathbf{R}} \frac{x}{x^2 + (y-t)^2} \alpha_{\xi'}(t) dt - \frac{i}{\pi} \int_{\mathbf{R}} \frac{y-t}{x^2 + (y-t)^2} \alpha_{\xi'}(t) dt \quad \text{and} \\ p_i(z) &= \frac{1}{\pi} \int_{\mathbf{R}} \frac{x}{x^2 + (y-t)^2} d\mu_{\xi',i}(t) - \frac{i}{\pi} \int_{\mathbf{R}} \frac{y-t}{x^2 + (y-t)^2} d\mu_{\xi',i}(t) \quad \text{for } i = 0, 1, 2. \end{aligned}$$

It follows from Lemmas 2.4 and 2.5 that r and p_i are well defined and holomorphic in the right half-plane $\{z \in \mathbf{C} \mid \operatorname{Re}(z) > 0\}$. Moreover, assumption (c) and Lemma 2.5(v) imply that p_0 admits an holomorphic extension to the domain $\{z \in \mathbf{C} \mid \operatorname{Re}(z) > 0 \text{ or } |\operatorname{Im}(z)| < \frac{\eta}{2}\}$, and $p_0(0) = 0$. Consequently, $\frac{p_0(z)}{z}$ is holomorphic in this domain and is bounded in a neighbourhood of zero.

Finally, we define

$$m_{\xi'}(z) = A_0(\xi') + A_1(\xi')z + A_2(\xi')z^2 + z^3 r(z) + \frac{p_0(z)}{z} + zp_1(z) + z^3 p_2(z). \quad (2.18)$$

It is obvious that $m_{\xi'}$ is well defined and holomorphic in the right half-plane. Since $\alpha_{\xi'}$ and $\mu_{\xi',i}$ are “even” and $t \mapsto \frac{t}{\xi_1^2 + t^2}$ is odd, for any $\xi_1 > 0$ we have $\operatorname{Im}(m_{\xi'}(\xi_1)) = 0$ and

$$m_{\xi'}(\xi_1) = \operatorname{Re}(m_{\xi'}(\xi_1)) = m(\xi_1, \xi').$$

For $\varepsilon, R > 0$, consider the closed continuous path $\gamma_{\varepsilon,R}$ composed by the following pieces:

$$\begin{aligned} \gamma_{1,\varepsilon,R}(t) &= t, & t &\in [\varepsilon, \varepsilon + R], \\ \gamma_{2,\varepsilon,R}(\theta) &= \varepsilon + Re^{i\theta}, & \theta &\in [0, \frac{\pi}{2}], \\ \gamma_{3,\varepsilon,R}(t) &= \varepsilon + i(R-t), & t &\in [0, R]. \end{aligned}$$

The function $z \mapsto m_{\xi'}(z)h(z)$ being holomorphic in the right half-plane we have $\int_{\gamma_{\varepsilon,R}} m_{\xi'}(z)h(z) dz = 0$, that is

$$\int_{\varepsilon}^R m(\xi_1, \xi') h(\xi_1) d\xi_1 + \int_{\gamma_{2,\varepsilon,R}} m_{\xi'}(z) h(z) dz + \int_{\gamma_{3,\varepsilon,R}} m_{\xi'}(z) h(z) dz = 0. \quad (2.19)$$

It follows from (2.18), Lemmas 2.4(iv) and 2.5(iv) that $\lim_{|z| \rightarrow \infty, \operatorname{Re}(z) \geq \varepsilon} \frac{m_{\xi'}(z)}{z^3} = 0$; hence, $\lim_{R \rightarrow \infty} \frac{m_{\xi'}(\varepsilon + Re^{i\theta})}{(\varepsilon + Re^{i\theta})^3} = 0$ uniformly with respect to $\theta \in [0, \frac{\pi}{2}]$. On the other hand, from Lemma 2.1(ii), we have $|h(\varepsilon + Re^{i\theta})| \leq \frac{C}{|\varepsilon + Re^{i\theta}|^4}$ and then $|(\varepsilon + Re^{i\theta})^3 h(\varepsilon + Re^{i\theta}) \cdot i Re^{i\theta}| \leq \frac{CR}{|\varepsilon + Re^{i\theta}|} \leq \frac{CR}{R-\varepsilon} \leq 2C$ for any $R \geq 2\varepsilon$. We infer that $\lim_{R \rightarrow \infty} \int_{\gamma_{2,\varepsilon,R}} m_{\xi'}(z) h(z) dz = 0$.

From (2.13), (2.14) and the boundedness of $\frac{p_0(z)}{z}$ near 0 it follows that $|m(\xi_1, \xi')| \leq C$ for $0 < \xi_1 < 1$ and $|m(\xi_1, \xi')| \leq C|\xi_1|^{3-\delta}$ for large ξ_1 and some $C, \delta > 0$. Since h is continuous and $|h(\xi_1)| \leq \frac{C}{|\xi_1|^4}$, the integral $\int_0^\infty m(\xi_1, \xi') h(\xi_1) d\xi_1$ converges absolutely.

Clearly we have $\int_{\gamma_{3,\varepsilon,R}} m_{\xi'}(z) h(z) dz = -i \int_0^R m_{\xi'}(\varepsilon + iy) h(\varepsilon + iy) dy$. Passing to the limit as $R \rightarrow \infty$ in (2.19) we infer that $\int_0^\infty m_{\xi'}(\varepsilon + iy) h(\varepsilon + iy) dy$ converges and

$$\int_{\varepsilon}^{\infty} m(\xi_1, \xi') h(\xi_1) d\xi_1 = i \int_0^{\infty} m_{\xi'}(\varepsilon + iy) h(\varepsilon + iy) dy. \quad (2.20)$$

Since $m(\xi_1, \xi')$ is real and symmetric with respect to ξ_1 and $h(-\xi_1) = \overline{h(\xi_1)}$, we have

$$\int_{-\infty}^{-\varepsilon} m(\xi_1, \xi') h(\xi_1) d\xi_1 = \int_{\varepsilon}^{\infty} m(\xi_1, \xi') \overline{h(\xi_1)} d\xi_1,$$

and then, taking (2.20) into account, we get

$$\int_{-\infty}^{-\varepsilon} m(\xi_1, \xi') h(\xi_1) d\xi_1 + \int_{\varepsilon}^{\infty} m(\xi_1, \xi') h(\xi_1) d\xi_1 = -2 \int_0^{\infty} \operatorname{Im}(m_{\xi'}(\varepsilon + iy) h(\varepsilon + iy)) dy; \quad (2.21)$$

hence

$$\int_{-\infty}^{\infty} m(\xi_1, \xi') h(\xi_1) d\xi_1 = -2 \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} \operatorname{Im}(m_{\xi'}(\varepsilon + iy) h(\varepsilon + iy)) dy. \quad (2.22)$$

Since $h(iy) \in \mathbf{R}$ for $y \in [0, \infty)$, using Lemma 2.1 and the Dominated Convergence Theorem we find

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} \operatorname{Im} \left[(A_0(\xi') + A_1(\xi')(\varepsilon + iy) + A_2(\xi')(\varepsilon + iy)^2) h(\varepsilon + iy) \right] dy \\
&= A_1(\xi') \int_0^{\infty} y h(iy) dy.
\end{aligned} \tag{2.23}$$

It is easy to see that $|(\varepsilon + iy)^\ell h(\varepsilon + iy) - (iy)^\ell h(iy)| \leq C_1 \varepsilon \min(1, \frac{1}{y^2})$ for $y \in (0, \infty)$, $\ell \in \{0, 1, 2, 3\}$ and $\varepsilon \in [0, 1]$. Hence there exists $C_2 > 0$ such that

$$\|(\varepsilon + iy)^\ell h(\varepsilon + iy) - (iy)^\ell h(iy)\|_{L^q(0, \infty)} \leq C_2 \varepsilon \tag{2.24}$$

for any $\varepsilon \in [0, 1]$, $\ell \in \{0, 1, 2, 3\}$ and $q \in [1, \infty]$. This implies that

$$\begin{aligned}
& \left| \int_0^{\infty} \operatorname{Im}((\varepsilon + iy)^3 h(\varepsilon + iy) r(\varepsilon + iy)) dy - \int_0^{\infty} \operatorname{Im}((iy)^3 h(iy) r(\varepsilon + iy)) dy \right| \\
& \leq (\|\operatorname{Re}(r(\varepsilon + i \cdot))\|_{L^p} + \|\operatorname{Im}(r(\varepsilon + i \cdot))\|_{L^p}) \|(\varepsilon + iy)^3 h(\varepsilon + iy) - (iy)^3 h(iy)\|_{L^{p'}(0, \infty)} \\
& \leq (\|\alpha_{\xi'}\|_{L^p} + \|H\alpha_{\xi'}\|_{L^p}) C_2 \varepsilon \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.
\end{aligned}$$

On the other hand, by Lemma 2.4(ii) we obtain

$$\begin{aligned}
\lim_{\varepsilon \rightarrow 0} \int_0^{\infty} \operatorname{Im}[(iy)^3 h(iy) r(\varepsilon + iy)] dy &= - \lim_{\varepsilon \rightarrow 0} \int_0^{\infty} y^3 h(iy) \operatorname{Re}[r(\varepsilon + iy)] dy \\
&= - \int_0^{\infty} y^3 h(iy) \alpha_{\xi'}(y) dy.
\end{aligned}$$

Therefore we have

$$\lim_{\varepsilon \rightarrow 0} \int_0^{\infty} \operatorname{Im}[(\varepsilon + iy)^3 h(\varepsilon + iy) r(\varepsilon + iy)] dy = - \int_0^{\infty} y^3 h(iy) \alpha_{\xi'}(y) dy. \tag{2.25}$$

Let $\chi \in C_c^\infty(\mathbf{R}, \mathbf{R}_+)$ be such that $\operatorname{supp}(\chi) \subset [-\frac{\eta}{4}, \frac{\eta}{4}]$ and $\chi \equiv 1$ on $[-\frac{\eta}{8}, \frac{\eta}{8}]$. Since the function $z \mapsto \frac{p_0(z)}{z} h(z)$ is uniformly continuous on $[-1, 1] \times [-\frac{\eta}{4}, \frac{\eta}{4}]$ we have

$$\begin{aligned}
\lim_{\varepsilon \rightarrow 0} \int_0^{\infty} \operatorname{Im} \left[\frac{p_0(\varepsilon + iy)}{\varepsilon + iy} h(\varepsilon + iy) \chi(y) \right] dy &= \int_0^{\infty} \operatorname{Im} \left(\frac{p_0(iy)}{iy} h(iy) \chi(y) \right) dy \\
&= - \int_0^{\infty} \frac{\operatorname{Re}(p_0(iy))}{y} h(iy) \chi(y) dy = 0.
\end{aligned} \tag{2.26}$$

By Lemma 2.1 we infer that there exists $C_3 > 0$ such that $|h(\varepsilon + iy) - h(iy)| \leq \varepsilon C_3 \min(1, \frac{1}{|y|^5})$ for any $y \in (0, \infty)$ and $\varepsilon \in [0, 1]$. It is easy to see that

$$\left| \left(\frac{h(\varepsilon + iy)}{\varepsilon + iy} - \frac{h(iy)}{iy} \right) (1 - \chi(y)) \right| \leq C_4 \varepsilon \min\left(\frac{1}{y^6}, 1\right)$$

for any $y \in (0, \infty)$ and some $C_4 > 0$. Consequently there exists $C_5 > 0$ such that

$$\left\| \left(\frac{h(\varepsilon + iy)}{\varepsilon + iy} - \frac{h(iy)}{iy} \right) (1 - \chi(y)) \right\|_{L^p(0, \infty)} \leq C_5 \varepsilon$$

for any $p \in [1, \infty]$. Using the Cauchy–Schwarz inequality and Lemma 2.5(ii) and (iii), we get

$$\begin{aligned} & \left| \int_0^\infty p_0(\varepsilon + iy) \left(\frac{h(\varepsilon + iy)}{\varepsilon + iy} - \frac{h(iy)}{iy} \right) (1 - \chi(y)) dy \right| \\ & \leq \|p_0(\varepsilon + i \cdot)\|_{L^2(\mathbf{R})} \left\| \left(\frac{h(\varepsilon + iy)}{\varepsilon + iy} - \frac{h(iy)}{iy} \right) (1 - \chi(y)) \right\|_{L^2(0, \infty)} \\ & \leq C_6 \varepsilon^{\frac{1}{2}} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \end{aligned} \quad (2.27)$$

We also have by (2.15) and assumption (c),

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \int_0^\infty \operatorname{Im} \left[p_0(\varepsilon + iy) \frac{h(iy)}{iy} (1 - \chi(y)) \right] dy \\ & = - \lim_{\varepsilon \rightarrow 0} \int_0^\infty \operatorname{Re}(p_0(\varepsilon + iy)) \frac{h(iy)}{y} (1 - \chi(y)) dy \\ & = - \int_0^\infty \frac{h(iy)}{y} (1 - \chi(y)) d\mu_{\xi', 0}(y) = - \int_0^\infty \frac{h(iy)}{y} d\mu_{\xi', 0}(y). \end{aligned} \quad (2.28)$$

From (2.26)–(2.28) we get

$$\lim_{\varepsilon \rightarrow 0} \int_0^\infty \operatorname{Im} \left[\frac{p_0(\varepsilon + iy)}{\varepsilon + iy} h(\varepsilon + iy) \right] dy = - \int_0^\infty \frac{h(iy)}{y} d\mu_{\xi', 0}(y). \quad (2.29)$$

Similarly we find

$$\lim_{\varepsilon \rightarrow 0} \int_0^\infty \operatorname{Im}((\varepsilon + iy) p_1(\varepsilon + iy) h(\varepsilon + iy)) dy = \int_0^\infty y h(iy) d\mu_{\xi', 1}(y) \quad \text{and} \quad (2.30)$$

$$\lim_{\varepsilon \rightarrow 0} \int_0^{\infty} \operatorname{Im}((\varepsilon + iy)^3 p_2(\varepsilon + iy) h(\varepsilon + iy)) dy = - \int_0^{\infty} y^3 h(iy) d\mu_{\xi', 2}(y). \quad (2.31)$$

Since $m_{\xi'}(z)$ is given by (2.18), replacing (2.23), (2.25), (2.29)–(2.31) into (2.22) we obtain the conclusion of Lemma 2.6. $\square$

Now we are ready to state and prove the main result of this section.

Theorem 2.7. Suppose that for any $\xi' \in \mathbf{R}^{N-1}$, $m(\xi_1, \xi')$ satisfies the assumptions of Lemma 2.6. For $u \in C_c^\infty(\mathbf{R}^N)$ define u_1, u_2, f, g and W as in (2.1)–(2.4). Then we have the identity:

$$\begin{aligned} & \frac{\pi^2}{16} (W(u_1) + W(u_2) - 2W(u)) \\ &= - \int_{\mathbf{R}^{N-1}} A_1(\xi') \int_0^\infty t \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt d\xi' \\ &+ \int_{\mathbf{R}^{N-1}} \int_0^\infty t^3 \alpha_{\xi'}(t) \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt d\xi' \\ &+ \int_{\mathbf{R}^{N-1}} \int_0^\infty \frac{1}{t} \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 d\mu_{\xi', 0}(t) d\xi' \\ &- \int_{\mathbf{R}^{N-1}} \int_0^\infty t \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 d\mu_{\xi', 1}(t) d\xi' \\ &+ \int_{\mathbf{R}^{N-1}} \int_0^\infty t^3 \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 d\mu_{\xi', 2}(t) d\xi'. \end{aligned} \quad (2.32)$$

Proof. Since $\mathcal{F}_{N-1} f \in \mathcal{S}(\mathbf{R}^N)$, the integral $\int_0^\infty e^{-x_1 t} (\mathcal{F}_{N-1} f)(x_1, \xi') dx_1$ is well defined for all $t > 0$ and $\xi' \in \mathbf{R}^{N-1}$. Using Plancherel's theorem we get

$$\begin{aligned} \int_0^\infty e^{-x_1 t} (\mathcal{F}_{N-1} f)(x_1, \xi') dx_1 &= \langle \mathcal{F}_{N-1} f(\cdot, \xi'), e^{-(\cdot)t} \chi_{[0, \infty)}(\cdot) \rangle_{L^2(\mathbf{R})} \\ &= (2\pi)^{-1} \langle \mathcal{F}_1(\mathcal{F}_{N-1} f(\cdot, \xi')), \mathcal{F}_1(e^{-(\cdot)t} \chi_{[0, \infty)}(\cdot)) \rangle_{L^2(\mathbf{R})}. \end{aligned} \quad (2.33)$$

Moreover, we have

$$\mathcal{F}_1(e^{-(\cdot)t} \chi_{[0, \infty)}(\cdot))(\xi_1) = \int_0^\infty e^{-ix_1 \xi_1} e^{-x_1 t} dx_1 = - \frac{1}{t + i\xi_1} e^{-(t+i\xi_1)x_1} \Big|_{x_1=0}^\infty = \frac{1}{t + i\xi_1}$$

and then, using (2.33) and the oddness of $\widehat{f}$ with respect to ξ_1 we get:

$$\begin{aligned} h_{\xi'}(it) &= \left| \int_0^\infty e^{-x_1 t} (\mathcal{F}_{N-1} f)(x_1, \xi') dx_1 \right|^2 = (2\pi)^{-2} \left| \int_{-\infty}^\infty \widehat{f}(\xi_1, \xi') \cdot \frac{1}{t - i\xi_1} d\xi_1 \right|^2 \\ &= \frac{1}{(2\pi)^2} \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \left(\frac{1}{t - i\xi_1} - \frac{1}{t + i\xi_1} \right) d\xi_1 \right|^2 \\ &= \frac{1}{\pi^2} \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2. \end{aligned} \quad (2.34)$$

Identity (2.32) is a simple consequence of (2.10), (2.17) and (2.34) and Theorem 2.7 is proved. $\square$

Remark 2.8. It is worth to note that we can prove an identity analogous to (2.32) whenever we work with a symbol $m(\xi) = m(\xi_1, \xi')$ symmetric with respect to ξ_1 and such that for any $\xi' \in \mathbf{R}^{N-1}$, $m(\cdot, \xi')$ admits an holomorphic extension $m_{\xi'}(z)$ to the domain $\{z \in \mathbf{C} \mid \operatorname{Re}(z) > 0, \operatorname{Im}(z) > 0\}$ having the following properties:

- P1.** $\lim_{z \rightarrow \xi_1, \operatorname{Im}(z) > 0} m_{\xi'}(z) = m(\xi_1, \xi')$.
- P2.** For any $\varepsilon > 0$, $\lim_{|z| \rightarrow \infty, \operatorname{Re}(z) \geq \varepsilon} \frac{m_{\xi'}(z)}{z^3} = 0$.
- P3.** $\lim_{\varepsilon \rightarrow 0} \int_0^\infty m_{\xi'}(\varepsilon + it) h_{\xi'}(\varepsilon + it) dt$ exists (and depends on ξ' and the values taken by $h_{\xi'}$ on the imaginary axis).

Note that assumption P1 implies that $m(\cdot, \xi')$ admits an holomorphic extension to the whole right half-plane. Indeed, it follows from Schwarz' reflection principle [8, p. 75] that the function

$$\widetilde{m}_{\xi'} = \begin{cases} m_{\xi'}(z) & \text{if } \operatorname{Im}(z) \geq 0, \\ \overline{m_{\xi'}(\bar{z})} & \text{if } \operatorname{Im}(z) < 0 \end{cases}$$

is holomorphic in $\{z \in \mathbf{C} \mid \operatorname{Re}(z) > 0\}$.

Assumption P2 is needed in the proof of Lemma 2.6 to show that

$$\lim_{R \rightarrow \infty} \int_{\gamma_{2,\varepsilon,R}} m_{\xi'}(z) h_{\xi'}(z) dz = 0$$

(where $\gamma_{2,\varepsilon,R}(\theta) = \varepsilon + Re^{i\theta}$, $\theta \in [0, \frac{\pi}{2}]$). We recall that $|h_{\xi'}(z)|$ behaves like $\frac{1}{|z|^4}$ as $|z| \rightarrow \infty$ (see Lemma 2.1 and Remark 2.2). This assumption could be replaced by a weaker one that guarantees at least that

$$\lim_{n \rightarrow \infty} \int_{\gamma_{2,\varepsilon,R_n}} m_{\xi'}(z) h_{\xi'}(z) dz = 0 \quad \text{for some sequence } R_n \rightarrow \infty.$$

In Theorem 2.7 assumption P3 is satisfied because of the special form of $m(\cdot, \xi')$ (see (2.16)).

Conversely, suppose that a function $m(z)$ has the properties P1–P3 above. Let $\tilde{m}$ be the holomorphic extension of m to the right half-plane and define $q(z) = \frac{\tilde{m}(z)}{z^3}$. Clearly, q is an holomorphic function in the right half-plane and $\lim_{|z| \rightarrow \infty, \operatorname{Re}(z) \geq \varepsilon} q(z) = 0$ for any $\varepsilon > 0$. Thus for any $x > \varepsilon$ we have the Poisson representation formulae

$$\begin{aligned} q(x+iy) &= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{x-\varepsilon}{(x-\varepsilon)^2 + (t-y)^2} \operatorname{Re}(q(\varepsilon+it)) dt \\ &\quad + \frac{i}{\pi} \int_{-\infty}^{\infty} \frac{t-y}{(x-\varepsilon)^2 + (t-y)^2} \operatorname{Re}(q(\varepsilon+it)) dt \end{aligned} \quad (2.35)$$

and

$$\begin{aligned} q(x+iy) &= \frac{-1}{\pi} \int_{-\infty}^{\infty} \frac{t-y}{(x-\varepsilon)^2 + (t-y)^2} \operatorname{Im}(q(\varepsilon+it)) dt \\ &\quad + \frac{i}{\pi} \int_{-\infty}^{\infty} \frac{x-\varepsilon}{(x-\varepsilon)^2 + (t-y)^2} \operatorname{Im}(q(\varepsilon+it)) dt. \end{aligned} \quad (2.36)$$

Multiplying (2.35) (respectively (2.36)) by $(x+iy)^3$, we find the expression of $m(x+iy)$ in terms of $\operatorname{Re}(q(\varepsilon+it))$ (respectively in terms of $\operatorname{Im}(q(\varepsilon+it))$). If $\operatorname{Re}(q(\varepsilon+it)) \rightarrow \alpha(t)$ as $\varepsilon \rightarrow 0$ and if it is possible to pass to the limit as $\varepsilon \rightarrow 0$ in (2.35) we obtain, at least formally,

$$m(\xi_1) = \xi_1^3 q(\xi_1) = \frac{\xi_1^4}{\pi} \int_{-\infty}^{\infty} \frac{\alpha(t)}{\xi_1^2 + t^2} dt.$$

However, as it will be seen later in applications, the function q may be singular at the origin. In this case it is not possible to pass to the limit as $\varepsilon \rightarrow 0$ in (2.35) or in (2.36) in order to express the function q (hence the function m) in terms of its “boundary values” on the imaginary axis. For this reason we have introduced “lower order terms” in the expression of $m_{\xi'}(z)$ in (2.16).

It is now clear that Theorem 2.7 can be generalized. For example, if the expression (2.16) of $m(\xi_1, \xi')$ contains other terms

$$\frac{1}{\pi} \sum_{k=0}^3 |\xi_1|^k \int_{\mathbf{R}} \frac{1}{\xi_1^2 + t^2} \alpha_{\xi',k}(t) dt,$$

where $\alpha_{\xi',k} \in L^{p_k}(\mathbf{R})$ for some $p_k \in (1, \infty)$, $\alpha_{\xi',k}$ are even functions and $\alpha_{\xi',0}$ vanishes in a neighborhood of zero, then we have to add terms

$$\int_{\mathbf{R}^{N-1}} \int_0^\infty \left[\frac{\alpha_{\xi',0}(t)}{t} + \beta_{\xi',1}(t) - t\alpha_{\xi',2}(t) - t^2\beta_{\xi',3}(t) \right] \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt d\xi'$$

in the right-hand side of (2.32), where $\beta_{\xi',1}$ and $\beta_{\xi',3}$ are Hilbert transforms of $\alpha_{\xi',1}$ and $\alpha_{\xi',3}$, respectively.

We give now some examples illustrating several situations that may be encountered in applications. Throughout $u \in C_c^\infty(\mathbf{R}^N)$ and we keep the notation introduced in (2.1)–(2.4).

Example 2.9. If the symbol m is of the form $m(\xi_1, \xi') = A_1(\xi')|\xi_1|$, then Theorem 2.7 gives

$$W(u_1) + W(u_2) - 2W(u) = -\frac{16}{\pi^2} \int_{\mathbf{R}^{N-1}} A_1(\xi') \int_0^\infty t \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt d\xi'. \quad (2.37)$$

This kind of symbol appears in problems involving operators of the type $H_1 \frac{\partial}{\partial x_1} P(\frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_N})$, where H_1 is the Hilbert transform with respect to the x_1 variable and P is a pseudo-differential operator in the last $N - 1$ variables.

Example 2.10. (i) Consider the symbol $m(\xi) = \frac{1}{|\xi|^2}$ appearing in Choquard's problem. It can be written as

$$m(\xi_1, \xi') = \frac{1}{\xi_1^2 + |\xi'|^2} = \frac{1}{\pi} \int_{\mathbf{R}} \frac{1}{\xi_1^2 + t^2} d\mu_{\xi',0}(t),$$

where $\mu_{\xi',0} = \frac{\pi}{2}(\delta_{-|\xi'|} + \delta_{|\xi'|})$ and δ_a is the Dirac measure with support $\{a\}$. From Theorem 2.7 we get the identity

$$W(u_1) + W(u_2) - 2W(u) = \frac{8}{\pi} \int_{\mathbf{R}^{N-1}} \frac{1}{|\xi'|} \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 \right|^2 d\xi'. \quad (2.38)$$

The same identity could be obtained by observing that the function $m_{\xi'}(z) = \frac{1}{z^2 + |\xi'|^2}$ is meromorphic in $\mathbf{C}$ and has exactly one pole in the upper half-plane, namely $i|\xi'|$. Using Residue's Theorem it is not hard to see that

$$\int_{-\infty}^{\infty} m_{\xi'}(z) h_{\xi'}(z) dz = 2\pi i \operatorname{Res}(m_{\xi'} h_{\xi'}, i|\xi'|),$$

and integrating this identity over $\mathbf{R}^{N-1}$ we get (2.38).

(ii) Consider the symbol $m(\xi) = \frac{1}{|\xi|^2 + a^2} = \frac{1}{\xi_1^2 + |\xi'|^2 + a^2}$ corresponding to the operator $(-\Delta + a^2)^{-1}$. It is obvious that

$$m(\xi_1, \xi') = \frac{1}{\pi} \int_{\mathbf{R}} \frac{1}{\xi_1^2 + t^2} d\mu_{\xi',0}(t),$$

where $\mu_{\xi',0} = \frac{\pi}{2}(\delta_{-\sqrt{|\xi'|^2 + a^2}} + \delta_{\sqrt{|\xi'|^2 + a^2}})$. From Theorem 2.7 we get the identity

$$W(u_1) + W(u_2) - 2W(u) = \frac{8}{\pi} \int_{\mathbf{R}^{N-1}} \frac{1}{\sqrt{|\xi'|^2 + a^2}} \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{a^2 + |\xi'|^2 + \xi_1^2} d\xi_1 \right|^2 d\xi'. \quad (2.39)$$

The same identity could be obtained by applying Residue's Theorem to the meromorphic function $z \mapsto \frac{1}{z^2 + |\xi'|^2 + a^2} h_{\xi'}(z)$.

(iii) More generally, consider a symbol of the form $m(\xi_1, \xi') = \frac{c(\xi')}{\xi_1^2 + r^2(\xi')}$. It can be written as

$$m(\xi_1, \xi') = \frac{1}{\pi} \int_{\mathbf{R}} \frac{1}{\xi_1^2 + t^2} d\mu_{\xi', 0}(t),$$

where $\mu_{\xi', 0} = \frac{\pi}{2} c(\xi') (\delta_{-r(\xi')} + \delta_{r(\xi')})$. Using Theorem 2.7 we obtain the identity

$$W(u_1) + W(u_2) - 2W(u) = \frac{8}{\pi} \int_{\mathbf{R}^{N-1}} \frac{c(\xi')}{r(\xi')} \cdot \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{r^2(\xi') + \xi_1^2} d\xi_1 \right|^2 d\xi'. \quad (2.40)$$

In particular, for the symbol $m(\xi_1, \xi') = \frac{\xi_j^{2k}}{\xi_1^2 + |\xi'|^2 + a^2}$, $j = 2, \dots, N$ (corresponding to the operator $(-1)^k \frac{\partial^{2k}}{\partial x_j^{2k}} (-\Delta + a^2)^{-1}$), we get

$$W(u_1) + W(u_2) - 2W(u) = \frac{8}{\pi} \int_{\mathbf{R}^{N-1}} \frac{\xi_j^{2k}}{\sqrt{|\xi'|^2 + a^2}} \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{a^2 + |\xi'|^2 + \xi_1^2} d\xi_1 \right|^2 d\xi'. \quad (2.41)$$

(iv) The symbol $m(\xi_1, \xi') = \frac{\xi_1^2}{\xi_1^2 + |\xi'|^2 + a^2}$ can be expressed as

$$m(\xi_1, \xi') = \frac{\xi_1^2}{\pi} \int_{\mathbf{R}} \frac{1}{\xi_1^2 + t^2} d\mu_{\xi', 1}(t),$$

where $\mu_{\xi', 1} = \frac{\pi}{2} (\delta_{-\sqrt{|\xi'|^2 + a^2}} + \delta_{\sqrt{|\xi'|^2 + a^2}})$. From Theorem 2.7 we find the identity

$$W(u_1) + W(u_2) - 2W(u) = -\frac{8}{\pi} \int_{\mathbf{R}^{N-1}} \sqrt{|\xi'|^2 + a^2} \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{a^2 + |\xi'|^2 + \xi_1^2} d\xi_1 \right|^2 d\xi'. \quad (2.42)$$

Notice that the right-hand side in (2.42) is nonpositive, while in (2.41) it is nonnegative.

(v) The symbol $m(\xi_1, \xi') = \frac{\xi_1^4}{\xi_1^2 + |\xi'|^2 + a^2}$ (corresponding to the operator $\frac{\partial^4}{\partial x_1^4}(-\Delta + a^2)^{-1}$) can be written as

$$m(\xi_1, \xi') = \frac{\xi_1^4}{\pi} \int_{\mathbf{R}} \frac{1}{\xi_1^2 + t^2} d\mu_{\xi', 2}(t),$$

where $\mu_{\xi', 2} = \frac{\pi}{2}(\delta_{-\sqrt{|\xi'|^2 + a^2}} + \delta_{\sqrt{|\xi'|^2 + a^2}})$. By Theorem 2.7 we have the identity

$$W(u_1) + W(u_2) - 2W(u) = \frac{8}{\pi} \int_{\mathbf{R}^{N-1}} (|\xi'|^2 + a^2)^{\frac{3}{2}} \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{a^2 + |\xi'|^2 + \xi_1^2} d\xi_1 \right|^2 d\xi'. \quad (2.43)$$

Obviously all the identities in (2.40)–(2.43) could be obtained by using the Residue Theorem.

Example 2.11. Consider the symbol $m(\xi) = |\xi|^{2s}$, corresponding to the operator $(-\Delta)^s$.

The complex logarithm $\log(z) = \ln|z| + i \arg(z)$ is well defined and holomorphic on $\mathbf{C} \setminus (-\infty, 0]$. For $z \in \Omega_{\xi'} := \mathbf{C} \setminus \{it \mid t \in (-\infty, -|\xi'|] \cup [|\xi'|, \infty)\}$, we have $z^2 + |\xi'|^2 \notin (-\infty, 0]$; hence we may define

$$m_{\xi'}(z) = e^{s \log(z^2 + |\xi'|^2)} = |z^2 + |\xi'|^2|^s e^{is \arg(z^2 + |\xi'|^2)}.$$

The function $m_{\xi'}$ is holomorphic in $\Omega_{\xi'}$ and $|m_{\xi'}(z)| = |z^2 + |\xi'|^2|^s$ for any $z \in \Omega_{\xi'}$. It is easy to see that, for $\xi' \neq 0$,

$$m_{\xi'}(z) = |\xi'|^{2s} \left(1 + s \frac{z^2}{|\xi'|^2} + \sum_{k=2}^\infty C_s^k \frac{z^{2k}}{|\xi'|^{2k}} \right), \quad (2.44)$$

where $C_s^k = \frac{s(s-1)\cdots(s-k+1)}{k!}$ and the series converges in the open ball $B_{\mathbf{C}}(0, |\xi'|)$.

For $s < \frac{3}{2}$ and $\xi' \neq 0$, the function $z \mapsto \frac{m_{\xi'}(z)}{z^3}$ is holomorphic in $\Omega_{\xi'} \setminus \{0\}$, tends to zero as $|z| \rightarrow \infty$ and has a third order pole at the origin. Consider the function $r_{\xi'}(z) = \frac{1}{z^3}(m_{\xi'}(z) - |\xi'|^{2s} - s|\xi'|^{2s-2}z^2)$. According to (2.44), $r_{\xi'}$ is a holomorphic function in $\Omega_{\xi'}$. If $s < \frac{3}{2}$, we have $r_{\xi'}(z) \rightarrow 0$ as $|z| \rightarrow \infty$. Consequently, the Poisson representation formula (2.35) holds for $r_{\xi'}$. Since $r_{\xi'}(\bar{z}) = \overline{r_{\xi'}(z)}$, the function $t \mapsto \operatorname{Re}(r_{\xi'}(\varepsilon + it))$ is even and we have, in particular,

$$\begin{aligned} m_{\xi'}(\xi_1) &= |\xi'|^{2s} + s|\xi'|^{2s-2}\xi_1^2 + \xi_1^3 r_{\xi'}(\xi_1) \\ &= |\xi'|^{2s} + s|\xi'|^{2s-2}\xi_1^2 + \frac{\xi_1^3}{\pi} \int_{-\infty}^\infty \frac{\xi_1 - \varepsilon}{(\xi_1 - \varepsilon)^2 + (t - y)^2} \operatorname{Re}(r_{\xi'}(\varepsilon + it)) dt. \end{aligned} \quad (2.45)$$

It is clear that for any $t \in (-|\xi'|, |\xi'|)$ we have $\lim_{\varepsilon \rightarrow 0} \operatorname{Re}(r_{\xi'}(\varepsilon + it)) = \operatorname{Re}(r_{\xi'}(it)) = 0$. For any $t > |\xi'|$ we have $\lim_{\varepsilon \downarrow 0} m_{\xi'}(\varepsilon + it) = (t^2 - |\xi'|^2)^s e^{is\pi}$ and $\lim_{\varepsilon \downarrow 0} \operatorname{Re}(r_{\xi'}(\varepsilon + it)) = -\sin(s\pi) \frac{(t^2 - |\xi'|^2)^s}{t^3}$.

On the other hand, it is straightforward to check that for $-1 < s < \frac{3}{2}$, there exists $p_s \in (1, \infty)$ and $C_{s,\xi'} > 0$ such that

$$\|r_{\xi'}(\varepsilon + i \cdot)\|_{L^{p_s}(\mathbf{R})} \leq C_{s,\xi'} \quad \text{for any } \varepsilon \in \left(0, \frac{|\xi'|}{2}\right). \quad (2.46)$$

It follows from (2.46) and [24, Theorem 2.5, p. 50] that there exists $k_{\xi'} \in L^{p_s}(\mathbf{R})$ such that $\operatorname{Re}(r_{\xi'}(x + iy)) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{x}{x^2 + (y-t)^2} k_{\xi'}(t) dt$. Moreover, from [24, Theorem 2.1, p. 47] we have $\lim_{\varepsilon \downarrow 0} \operatorname{Re}(r_{\xi'}(\varepsilon + it)) = k_{\xi'}(t)$ for almost every $t \in \mathbf{R}$ and $\|\operatorname{Re}(r_{\xi'}(\varepsilon + i \cdot)) - k_{\xi'}\|_{L^{p_s}} \rightarrow 0$ as $\varepsilon \rightarrow 0$. In view of the pointwise convergence, we infer that $k_{\xi'}(-t) = k_{\xi'}(t)$ a.e. and

$$k_{\xi'}(t) = \begin{cases} 0 & \text{if } t \in (-|\xi'|, |\xi'|), \\ -\sin(s\pi) \frac{(t^2 - |\xi'|^2)^s}{|t|^3} & \text{if } |t| > |\xi'|, \end{cases} \quad \text{a.e. on } \mathbf{R}.$$

Now it is clear that the symbol $m(\xi_1, \xi') = (\xi_1^2 + |\xi'|^2)^s$ can be written as

$$\begin{aligned} m(\xi_1, \xi') &= |\xi'|^{2s} + s|\xi'|^{2s-2}\xi_1^2 + \xi_1^3 r_{\xi'}(\xi_1) \\ &= |\xi'|^{2s} + s|\xi'|^{2s-2}\xi_1^2 + \frac{\xi_1^4}{\pi} \int_{-\infty}^{\infty} \frac{1}{\xi_1^2 + t^2} k_{\xi'}(t) dt. \end{aligned} \quad (2.47)$$

Thus we may apply Theorem 2.7 to get, for any $u \in C_c^\infty(\mathbf{R}^N)$ and $s \in (-1, \frac{3}{2})$,

$$\begin{aligned} W(u_1) + W(u_2) - 2W(u) &= \frac{16}{\pi^2} \int_{\mathbf{R}^{N-1}} \int_0^\infty t^3 k_{\xi'}(t) \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt d\xi' \\ &= -\frac{16 \sin(s\pi)}{\pi^2} \int_{\mathbf{R}^{N-1}} \int_{|\xi'|}^\infty (t^2 - |\xi'|^2)^s \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt d\xi'. \end{aligned} \quad (2.48)$$

Similarly, if we consider the symbol $m(\xi) = (|\xi|^2 + a^2)^s$ we get the identity

$$\begin{aligned} W(u_1) + W(u_2) - 2W(u) &= -\frac{16 \sin(s\pi)}{\pi^2} \int_{\mathbf{R}^{N-1}} \int_{\sqrt{|\xi'|^2 + a^2}}^\infty (t^2 - |\xi'|^2 - a^2)^s \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt d\xi'. \end{aligned} \quad (2.49)$$

3. Symmetry and function spaces

For any $u \in C_c^\infty(\mathbf{R}^N)$ we define u_1 and u_2 as in (2.2) and we put $T_1 u = u_1$, $T_2 u = u_2$. Clearly, T_1 and T_2 are linear continuous mappings from $C_c^\infty(\mathbf{R}^N)$ to $C_c^0(\mathbf{R}^N)$. In this section we consider the following intimately related problems.

Problem 1. Determine significant subspaces $\mathcal{X} \subset \mathcal{D}'(\mathbf{R}^N)$ such that T_1 and T_2 can be extended to linear continuous mappings from $\mathcal{X}$ to $\mathcal{X}$. (Or, equivalently, find subspaces $\mathcal{X}$ such that $u \in \mathcal{X}$ implies $T_1 u, T_2 u \in \mathcal{X}$ and $u \mapsto T_1 u, u \mapsto T_2 u$ are continuous for the $\mathcal{X}$ topology.)

Problem 2. If $\mathcal{X}$ is a subspace as above, how the identities proved in the previous section can be extended to $\mathcal{X}$?

The answer to these questions is of great importance in symmetry problems. For instance, suppose that a function space $\mathcal{X}$ has the two properties described above and that the solutions of the variational problem

$$\begin{aligned} &\text{minimize} \quad E(u) := \int_{\mathbf{R}^N} m(\xi) |\widehat{u}(\xi)|^2 d\xi + \int_{\mathbf{R}^N} F(u) dx \\ &\text{under the constraint} \quad \int_{\mathbf{R}^N} G(u) dx = \lambda \neq 0 \end{aligned} \quad (3.1)$$

belong to $\mathcal{X}$. As before, the symbol $m(\xi) = m(\xi_1, \xi')$ is assumed to be symmetric with respect to ξ_1 . Defining $W(u) := \int_{\mathbf{R}^N} m(\xi) |\widehat{u}(\xi)|^2 d\xi$, we suppose also that an identity of type (2.32) holds for $W(u)$ and it can be extended to $\mathcal{X}$ in such a way that

$$W(T_1 u) + W(T_2 u) - 2W(u) < 0 \quad \text{whenever } T_1 u \neq u, T_2 u \neq u.$$

(We will see later that most of the symbols in Examples 2.9–2.11 have this property.) Then, we claim that after a translation in the x_1 direction, any solution of (3.1) is symmetric with respect to x_1 . Indeed, let u be a minimizer. After a translation in the x_1 direction, we may assume that

$$\int_{\{x_1 < 0\}} G(u(x)) dx = \int_{\{x_1 > 0\}} G(u(x)) dx = \frac{\lambda}{2}.$$

Denoting $u_1 = T_1 u, u_2 = T_2 u$, this implies

$$\int_{\mathbf{R}^N} G(u_1(x)) dx = 2 \int_{\{x_1 < 0\}} G(u(x)) dx = \lambda \quad \text{and} \quad \int_{\mathbf{R}^N} G(u_2(x)) dx = 2 \int_{\{x_1 > 0\}} G(u(x)) dx = \lambda;$$

consequently u_1 and u_2 (which belong to $\mathcal{X}$) also satisfy the constraint. It is obvious that

$$\int_{\mathbf{R}^N} F(u_1(x)) dx + \int_{\mathbf{R}^N} F(u_2(x)) dx = 2 \int_{\mathbf{R}^N} F(u(x)) dx.$$

Suppose by contradiction that u is not symmetric with respect to x_1 . Then we get

$$E(u_1) + E(u_2) - 2E(u) = W(u_1) + W(u_2) - 2W(u) < 0,$$

and this implies that either $E(u_1) < E(u)$ or $E(u_2) < E(u)$. Therefore u cannot be a minimizer and this proves the claim.

Given the motivation above, we will study the behavior of T_1 and T_2 from $H^s(\mathbf{R}^N)$ to $H^s(\mathbf{R}^N)$, respectively from $\dot{H}^s(\mathbf{R}^N)$ to $\dot{H}^s(\mathbf{R}^N)$, where

$$H^s(\mathbf{R}^N) = \left\{ u \in \mathcal{S}'(\mathbf{R}^N) \mid \widehat{u} \in L^1_{\text{loc}}(\mathbf{R}^N) \text{ and } \int_{\mathbf{R}^N} (1 + |\xi|^2)^s |\widehat{u}(\xi)|^2 d\xi < \infty \right\},$$

$$\dot{H}^s(\mathbf{R}^N) = \left\{ u \in \mathcal{S}'(\mathbf{R}^N) \mid \widehat{u} \in L^1_{\text{loc}}(\mathbf{R}^N) \text{ and } \int_{\mathbf{R}^N} |\xi|^{2s} |\widehat{u}(\xi)|^2 d\xi < \infty \right\}.$$

It happens that T_1 and T_2 are not well defined from $H^s(\mathbf{R}^N)$ to $H^s(\mathbf{R}^N)$ (respectively from $\dot{H}^s(\mathbf{R}^N)$ to $\dot{H}^s(\mathbf{R}^N)$) if $s \geq \frac{3}{2}$ or if $s \leq -\frac{1}{2}$, as it can be seen in the following example.

Example 3.1. (i) Define $\varphi: \mathbf{R} \rightarrow \mathbf{R}$, $\varphi(x) = xe^{-|x|}$. An easy computation shows that $\widehat{\varphi}(\xi) = \frac{-4i\xi}{(1+\xi^2)^2}$, hence $\varphi \in H^s(\mathbf{R})$ for any $s < \frac{5}{2}$ and $\varphi \in \dot{H}^s(\mathbf{R})$ for any $s \in (-\frac{3}{2}, \frac{5}{2})$. It is clear that $(T_1\varphi)(x) = -|x|e^{-|x|}$ and $\widehat{T_1\varphi}(\xi) = \frac{2(\xi^2-1)}{(1+\xi^2)^2}$. Consequently, $T_1\varphi \in H^s(\mathbf{R})$ for $s < \frac{3}{2}$ (respectively $T_1\varphi \in \dot{H}^s(\mathbf{R})$ for $-\frac{1}{2} < s < \frac{3}{2}$), but $T_1\varphi \notin H^s(\mathbf{R})$ and $T_1\varphi \notin \dot{H}^s(\mathbf{R})$ for $s \geq \frac{3}{2}$.

In dimension $N \geq 2$ it suffices to take $\psi(x) = \varphi(x_1)\varphi_1(x_2, \dots, x_N)$, where $\varphi_1 \in C_c^\infty(\mathbf{R}^{N-1})$, to see that T_1 and T_2 are not well defined from $H^s(\mathbf{R}^N)$ to $H^s(\mathbf{R}^N)$ (respectively from $\dot{H}^s(\mathbf{R}^N)$ to $\dot{H}^s(\mathbf{R}^N)$) if $\frac{3}{2} \leq s < \frac{5}{2}$.

(ii) If $s < 0$, the elements of $H^s(\mathbf{R}^N)$ or $\dot{H}^s(\mathbf{R}^N)$ are not necessarily measurable functions. In this case we extend T_1 and T_2 to $H^s(\mathbf{R}^N)$ or $\dot{H}^s(\mathbf{R}^N)$ by duality. For $u, \varphi \in C_c^\infty(\mathbf{R}^N)$ we have

$$\begin{aligned} \langle T_1 u, \varphi \rangle_{\mathcal{S}', \mathcal{S}} &= \int_{\mathbf{R}^N} (T_1 u)(x) \varphi(x) dx = \int_{\{x_1 < 0\}} u(x) \varphi(x) dx + \int_{\{x_1 > 0\}} u(-x_1, x') \varphi(x) dx \\ &= \int_{\{x_1 < 0\}} u(x) \varphi(x) dx + \int_{\{x_1 < 0\}} u(x_1, x') \varphi(-x_1, x') dx = \langle u, T_1^* \varphi \rangle_{L^2, L^2}, \end{aligned}$$

where $(T_1^* \varphi)(x) = \chi_{\{x_1 < 0\}}(\varphi(x_1, x') + \varphi(-x_1, x'))$. Hence, for $u \in H^s(\mathbf{R}^N)$ with $s < 0$ we should define $T_1 u$ by

$$\langle T_1 u, \varphi \rangle_{H^s, H^{-s}} = \langle u, T_1^* \varphi \rangle_{H^s, H^{-s}}$$

for any test function $\varphi \in C_c^\infty(\mathbf{R}^N)$. However, the operator T_1^* does not map $H^k(\mathbf{R}^N)$ into $H^k(\mathbf{R}^N)$ if $k \geq \frac{1}{2}$ (as it can be easily seen by taking the function $\eta(x) = e^{-|x|}$ in one dimension, respectively $\eta(x_1)\eta_1(x_2, \dots, x_N)$, where $\eta_1 \in C_c^\infty(\mathbf{R}^{N-1})$ in dimension $N \geq 2$). This shows that we cannot define T_1 and T_2 on $H^s(\mathbf{R}^N)$ and on $\dot{H}^s(\mathbf{R}^N)$ if $s \leq -\frac{1}{2}$.

Our next goal is to prove that the operators T_1 and T_2 are well defined and continuous from $H^s(\mathbf{R}^N)$ to $H^s(\mathbf{R}^N)$ (respectively from $\dot{H}^s(\mathbf{R}^N)$ to $\dot{H}^s(\mathbf{R}^N)$) if $-\frac{1}{2} < s < \frac{3}{2}$. It is obvious that T_1 and T_2 are well defined and continuous from $L^2(\mathbf{R}^N)$ to $L^2(\mathbf{R}^N)$. It is well known that $H^1(\mathbf{R}^N) = W^{1,2}(\mathbf{R}^N) = \{\varphi \in L^2(\mathbf{R}^N) \mid \frac{\partial \varphi}{\partial x_i} \in L^2(\mathbf{R}^N), i = 1, \dots, N\}$ and that $T_1, T_2: W^{1,2}(\mathbf{R}^N) \rightarrow W^{1,2}(\mathbf{R}^N)$ are well defined and continuous. Using interpolation theory we

conclude that T_1 and T_2 are well defined and continuous from $H^s(\mathbf{R}^N)$ to $H^s(\mathbf{R}^N)$ if $0 \leq s \leq 1$. However, interpolation gives no information if either $s < 0$ or $s > 1$. Our next result deals with any value of s in $(-\frac{1}{2}, \frac{3}{2})$.

Theorem 3.2. *The operators T_1 and T_2 are well defined and continuous from $H^s(\mathbf{R}^N)$ to $H^s(\mathbf{R}^N)$ and from $\dot{H}^s(\mathbf{R}^N)$ to $\dot{H}^s(\mathbf{R}^N)$ for any $s \in (-\frac{1}{2}, \frac{3}{2})$.*

Proof. We will prove that there exists $C_s > 0$ such that for any $u \in C_c^\infty(\mathbf{R}^N)$ we have

$$\|T_i u\|_{H^s} \leq C_s \|u\|_{H^s}, \quad \text{respectively } \|T_i u\|_{\dot{H}^s} \leq C_s \|u\|_{\dot{H}^s}, \quad i = 1, 2, \quad (3.2)$$

and then the theorem will follow by density.

Therefore, suppose $u \in C_c^\infty(\mathbf{R}^N)$. By (2.48) and (2.49) we have

$$\begin{aligned} & \|T_1 u\|_{\dot{H}^s}^2 + \|T_2 u\|_{\dot{H}^s}^2 - 2\|u\|_{\dot{H}^s}^2 \\ &= -\frac{16 \sin(s\pi)}{\pi^2} \int_{\mathbf{R}^{N-1}} \int_{|\xi'|}^\infty (t^2 - |\xi'|^2)^s \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt d\xi', \end{aligned} \quad (3.3)$$

respectively

$$\begin{aligned} & \|T_1 u\|_{H^s}^2 + \|T_2 u\|_{H^s}^2 - 2\|u\|_{H^s}^2 \\ &= -\frac{16 \sin(s\pi)}{\pi^2} \int_{\mathbf{R}^{N-1}} \int_{\sqrt{|\xi'|^2+1}}^\infty (t^2 - |\xi'|^2 - 1)^s \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt d\xi'. \end{aligned} \quad (3.4)$$

If $N = 1$ we use the convention $\mathbf{R}^0 = \{0\}$ and the measure of $\{0\}$ is 1.

We begin by proving that T_1 and T_2 are bounded from $\dot{H}^s(\mathbf{R})$ to $\dot{H}^s(\mathbf{R})$, $-\frac{1}{2} < s < \frac{3}{2}$. For $N = 1$, the integral in the right-hand side of (3.3) can be formally written as

$$\int_0^\infty \int_0^\infty \int_0^\infty t^{2s} \frac{\xi}{t^2 + \xi^2} \cdot \frac{\eta}{t^2 + \eta^2} \widehat{f}(\xi) \overline{\widehat{f}(\eta)} d\xi d\eta dt. \quad (3.5)$$

Our strategy is as follows: first we compute explicitly the integral

$$I_s(\xi, \eta) = \int_0^\infty t^{2s} \frac{\xi}{t^2 + \xi^2} \cdot \frac{\eta}{t^2 + \eta^2} dt = \xi \eta \int_0^\infty t^{2s} \frac{1}{t^2 + \xi^2} \cdot \frac{1}{t^2 + \eta^2} dt. \quad (3.6)$$

Observe that $I_s(\xi, \eta) > 0$ if $\xi > 0$, $\eta > 0$. Then we will prove that for any $s \in (-\frac{1}{2}, \frac{3}{2})$ and any $\varphi, \psi \in L^2(0, \infty)$ we have

$$\left| \int_0^\infty \int_0^\infty \xi^{-s} \eta^{-s} I_s(\xi, \eta) \varphi(\xi) \psi(\eta) d\xi d\eta \right| \leq C(s) \|\varphi\|_{L^2(0, \infty)} \cdot \|\psi\|_{L^2(0, \infty)}.$$

This will be done in Lemma 3.3. Thereafter it will be clear that for any $f \in \dot{H}^s(\mathbf{R})$ we have

$$\begin{aligned} & \int_0^\infty \int_0^\infty I_s(\xi, \eta) |\widehat{f}(\xi)| \cdot |\overline{\widehat{f}(\eta)}| d\xi d\eta \\ &= \int_0^\infty \int_0^\infty \xi^{-s} \eta^{-s} I_s(\xi, \eta) |\xi^s \widehat{f}(\xi)| \cdot |\eta^s \overline{\widehat{f}(\eta)}| d\xi d\eta \\ &\leq C(s) \|\cdot\|^2_{L^2(0, \infty)} \leq C(s) \|f\|^2_{\dot{H}^s(\mathbf{R})}. \end{aligned} \quad (3.7)$$

This justifies the use of Fubini's Theorem in evaluating (3.5) and proves that the right-hand side of (3.3) is less than $C_1(s) \|f\|^2_{\dot{H}^s(\mathbf{R})}$, where $C_1(s)$ is a constant depending only on s . Thus we infer that there exists $C_s > 0$ such that $\|T_1 u\|_{\dot{H}^s(\mathbf{R})} \leq C_s \|u\|_{\dot{H}^s(\mathbf{R})}$ and $\|T_2 u\|_{\dot{H}^s(\mathbf{R})} \leq C_s \|u\|_{\dot{H}^s(\mathbf{R})}$ for any $u \in C_c^\infty(\mathbf{R})$. Consequently, T_1 and T_2 can be extended as continuous linear mappings from $\dot{H}^s(\mathbf{R})$ to $\dot{H}^s(\mathbf{R})$, $-\frac{1}{2} < s < \frac{3}{2}$, as claimed.

To carry out the first step of this strategy, we come back to $I_s(\xi, \eta)$ given by (3.6). Since the complex logarithm can be defined analytically on $\mathbf{C} \setminus \{it \mid t \in (-\infty, 0]\}$, we may define the holomorphic function $z \mapsto z^{2s} := e^{2s \log(z)} = |z|^{2s} e^{2is \arg(z)}$ on $\mathbf{C} \setminus \{it \mid t \in (-\infty, 0]\}$. With this definition the function $k(z) = \frac{z^{2s}}{(z^2 + \xi^2)(z^2 + \eta^2)}$ is meromorphic on $\mathbf{C} \setminus \{it \mid t \in (-\infty, 0]\}$. If $\xi \neq \eta$, k has two simple poles, namely $i\xi$ and $i\eta$; if $\xi = \eta$ it has a double pole at $i\xi$. For $0 < \varepsilon < \min(\xi, \eta)$, and $R > \max(\xi, \eta)$, consider the closed path $\beta_{\varepsilon, R}$ composed by the following pieces:

$$\begin{aligned} \beta_{1, \varepsilon, R}(t) &= t, & t &\in [-R, -\varepsilon], \\ \beta_{2, \varepsilon}(\theta) &= \varepsilon e^{i(\pi - \theta)}, & \theta &\in [0, \pi], \\ \beta_{3, \varepsilon, R}(t) &= t, & t &\in [\varepsilon, R], \\ \beta_{4, R}(\theta) &= R e^{i\theta}, & \theta &\in [0, \pi]. \end{aligned}$$

If $\xi \neq \eta$, using the Residue Theorem we get

$$\int_{\beta_{\varepsilon, R}} k(z) dz = 2\pi i [\text{Res}(k, i\xi) + \text{Res}(k, i\eta)] = \pi e^{is\pi} \left[\frac{\xi^{2s}}{\xi(\eta^2 - \xi^2)} + \frac{\eta^{2s}}{\eta(\xi^2 - \eta^2)} \right]. \quad (3.8)$$

Since $s > -\frac{1}{2}$ we have $\lim_{\varepsilon \rightarrow 0} \int_{\beta_{2, \varepsilon}} k(z) dz = 0$. We have also $\lim_{R \rightarrow \infty} \int_{\beta_{4, R}} k(z) dz = 0$ because $s < \frac{3}{2}$. Passing to the limit as $\varepsilon \rightarrow 0$ in (3.8) and then passing to the limit as $R \rightarrow \infty$ in the resulting equation, we get

$$\int_{-\infty}^0 k(z) dz + \int_0^\infty k(z) dz = \pi e^{is\pi} \frac{\xi^{2s-1} - \eta^{2s-1}}{\eta^2 - \xi^2},$$

that is

$$(e^{2is\pi} + 1) \int_0^\infty \frac{t^{2s}}{(t^2 + \xi^2)(t^2 + \eta^2)} dt = \pi e^{is\pi} \frac{\xi^{2s-1} - \eta^{2s-1}}{\eta^2 - \xi^2}.$$

For $s \neq \frac{1}{2}$ we obtain

$$\int_0^\infty \frac{t^{2s}}{(t^2 + \xi^2)(t^2 + \eta^2)} dt = \frac{\pi}{2 \cos(s\pi)} \frac{\xi^{2s-1} - \eta^{2s-1}}{\eta^2 - \xi^2}. \quad (3.9)$$

For $s = \frac{1}{2}$ we compute directly

$$\int_0^\infty \frac{t}{(t^2 + \xi^2)(t^2 + \eta^2)} dt = \frac{1}{\eta^2 - \xi^2} \int_0^\infty \frac{t}{t^2 + \xi^2} - \frac{t}{t^2 + \eta^2} dt = \frac{\ln \eta - \ln \xi}{\eta^2 - \xi^2}. \quad (3.10)$$

Hence

$$\begin{aligned} I_s(\xi, \eta) &= \frac{\pi}{2 \cos(s\pi)} \frac{\xi \eta (\xi^{2s-1} - \eta^{2s-1})}{\eta^2 - \xi^2} \quad \text{if } s \neq \frac{1}{2}, \quad \text{and} \\ I_{\frac{1}{2}}(\xi, \eta) &= \frac{\xi \eta (\ln \eta - \ln \xi)}{\eta^2 - \xi^2}. \end{aligned} \quad (3.11)$$

This gives

$$\xi^{-s} \eta^{-s} I_s(\xi, \eta) = \frac{\pi}{2 \cos(s\pi)} \frac{\xi^s \eta^{1-s} - \xi^{1-s} \eta^s}{\eta^2 - \xi^2} \quad \text{if } s \neq \frac{1}{2},$$

and

$$\xi^{-\frac{1}{2}} \eta^{-\frac{1}{2}} I_{\frac{1}{2}}(\xi, \eta) = \xi^{\frac{1}{2}} \eta^{\frac{1}{2}} \frac{\ln \eta - \ln \xi}{\eta^2 - \xi^2}.$$

An interesting property of these functions is given by the next lemma.

Lemma 3.3. Let $K_s(\xi, \eta) = \frac{\xi^s \eta^{1-s} - \xi^{1-s} \eta^s}{\eta^2 - \xi^2}$ if $s \neq \frac{1}{2}$, respectively $K_{\frac{1}{2}}(\xi, \eta) = \xi^{\frac{1}{2}} \eta^{\frac{1}{2}} \frac{\ln \eta - \ln \xi}{\eta^2 - \xi^2}$. For any $s \in (-\frac{1}{2}, \frac{3}{2})$ there exists a constant $C(s)$ (depending only on s) such that for any $\varphi, \psi \in L^2(0, \infty)$ we have

$$\left| \int_0^\infty \int_0^\infty \varphi(\xi) K_s(\xi, \eta) \psi(\eta) d\xi d\eta \right| \leq C(s) \|\varphi\|_{L^2(0, \infty)} \|\psi\|_{L^2(0, \infty)}.$$

Proof. Using polar coordinates we write $\xi = r \cos(\theta)$, $\eta = r \sin(\theta)$, where $r = \sqrt{\xi^2 + \eta^2}$ and $\theta = \arctan \frac{\eta}{\xi}$. It is easy to see that $K_s(\xi, \eta) = \frac{1}{r} L_s(\theta)$, where

$$L_s(\theta) = \frac{(\sin \theta)^s (\cos \theta)^{1-s} - (\cos \theta)^s (\sin \theta)^{1-s}}{\cos^2 \theta - \sin^2 \theta} \quad \text{if } s \neq \frac{1}{2},$$

and

$$L_{\frac{1}{2}}(\theta) = \frac{-\ln \tan \theta}{(1 - \tan^2 \theta) \cos^2 \theta} (\sin \theta)^{\frac{1}{2}} (\cos \theta)^{\frac{1}{2}}.$$

By a change of variables we get

$$\int_0^\infty \int_0^\infty |\varphi(\xi) K_s(\xi, \eta) \psi(\eta)| d\xi d\eta = \int_0^{\frac{\pi}{2}} \int_0^\infty |\varphi(r \cos \theta) \psi(r \sin \theta)| dr |L_s(\theta)| d\theta.$$

Using the Cauchy–Schwarz inequality we have

$$\int_0^\infty |\varphi(r \cos \theta) \psi(r \sin \theta)| dr \leq \|\varphi(\cdot \cos \theta)\|_{L^2(0, \infty)} \|\psi(\cdot \sin \theta)\|_{L^2(0, \infty)} = \frac{\|\varphi\|_{L^2(0, \infty)} \|\psi\|_{L^2(0, \infty)}}{\sqrt{\cos \theta \cdot \sin \theta}}.$$

Consequently,

$$\int_0^\infty \int_0^\infty |\varphi(\xi) K_s(\xi, \eta) \psi(\eta)| d\xi d\eta \leq \|\varphi\|_{L^2(0, \infty)} \|\psi\|_{L^2(0, \infty)} \int_0^{\frac{\pi}{2}} \frac{|L_s(\theta)|}{\sqrt{\cos \theta \cdot \sin \theta}} d\theta. \quad (3.12)$$

The lemma will be proved if we show that the last integral in (3.12) is finite. If $s \neq \frac{1}{2}$ we have

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \frac{|L_s(\theta)|}{\sqrt{\cos \theta \cdot \sin \theta}} d\theta &= \int_0^{\frac{\pi}{2}} \left| \frac{(\sin \theta)^{s-\frac{1}{2}} (\cos \theta)^{\frac{1}{2}-s} - (\cos \theta)^{s-\frac{1}{2}} (\sin \theta)^{\frac{1}{2}-s}}{\cos^2 \theta - \sin^2 \theta} \right| d\theta \\ &= \int_0^{\frac{\pi}{2}} \left| \frac{(\tan \theta)^{s-\frac{1}{2}} - (\tan \theta)^{\frac{1}{2}-s}}{1 - \tan^2 \theta} \right| \cdot \frac{1}{\cos^2 \theta} d\theta \\ &= \int_0^\infty \left| \frac{t^{s-\frac{1}{2}} - t^{\frac{1}{2}-s}}{1 - t^2} \right| dt. \end{aligned} \quad (3.13)$$

Using l'Hôspital's rule it is easy to see that $\lim_{t \rightarrow 1} \frac{t^{s-\frac{1}{2}} - t^{\frac{1}{2}-s}}{1 - t^2} = \frac{1}{2} - s$; hence the function $t \mapsto \frac{t^{s-\frac{1}{2}} - t^{\frac{1}{2}-s}}{1 - t^2}$ is bounded near 1. Since $s - \frac{1}{2} \in (-1, 1)$, the last integral in (3.13) converges.

If $s = \frac{1}{2}$ we have

$$\int_0^{\frac{\pi}{2}} \frac{|L_{\frac{1}{2}}(\theta)|}{\sqrt{\cos \theta \cdot \sin \theta}} d\theta = \int_0^{\frac{\pi}{2}} \left| \frac{-\ln \tan \theta}{1 - \tan^2 \theta} \right| \cdot \frac{1}{\cos^2 \theta} d\theta = \int_0^{\infty} \left| \frac{\ln y}{y^2 - 1} \right| dy. \quad (3.14)$$

Note that $\lim_{y \rightarrow 1} \frac{\ln y}{y^2 - 1} = \frac{1}{2}$ and this implies that the last integral in (3.14) converges. This completes the proof of Lemma 3.3. $\square$

In view of (3.3), (3.5), (3.7), (3.11) and Lemma 3.3, it follows that T_1 and T_2 are well defined and continuous from $\dot{H}^s(\mathbf{R})$ to $\dot{H}^s(\mathbf{R})$ for $-\frac{1}{2} < s < \frac{3}{2}$.

Next we prove that T_1 and T_2 are continuous from $H^s(\mathbf{R})$ to $H^s(\mathbf{R})$. We estimate the integral in the right-hand side of (3.4) for $N = 1$. If $s \in [0, \frac{3}{2})$ we have by (3.5)–(3.7)

$$\begin{aligned} \int_1^{\infty} (t^2 - 1)^s \left| \int_0^{\infty} \widehat{f}(\xi) \frac{\xi}{t^2 + \xi^2} d\xi \right|^2 dt &\leq \int_0^{\infty} t^{2s} \left| \int_0^{\infty} \widehat{f}(\xi) \frac{\xi}{t^2 + \xi^2} d\xi \right|^2 dt \\ &\leq C(s) \|f\|_{\dot{H}^s}^2 \leq C(s) \|f\|_{H^s}^2. \end{aligned} \quad (3.15)$$

If $s \in (-\frac{1}{2}, 0)$, using the change of variable $\tau = \sqrt{t^2 - 1}$ and (3.9) we get

$$\begin{aligned} \int_1^{\infty} \frac{(t^2 - 1)^s}{(t^2 + \xi^2)(t^2 + \eta^2)} dt &= \int_0^{\infty} \frac{\tau^{2s}}{(\tau^2 + 1 + \xi^2)(\tau^2 + 1 + \eta^2)} \cdot \frac{\tau}{\sqrt{\tau^2 + 1}} d\tau \\ &\leq \int_0^{\infty} \frac{\tau^{2s}}{(\tau^2 + 1 + \xi^2)(\tau^2 + 1 + \eta^2)} d\tau \\ &= \frac{\pi}{2 \cos(s\pi)} \cdot \frac{(1 + \xi^2)^{\frac{2s-1}{2}} - (1 + \eta^2)^{\frac{2s-1}{2}}}{\eta^2 - \xi^2}. \end{aligned} \quad (3.16)$$

Consequently,

$$\begin{aligned} \int_1^{\infty} (t^2 - 1)^s \left| \int_0^{\infty} \widehat{f}(\xi) \frac{\xi}{t^2 + \xi^2} d\xi \right|^2 dt &\leq \int_0^{\infty} \int_0^{\infty} |\widehat{f}(\xi)| \cdot |\overline{\widehat{f}(\eta)}| \int_1^{\infty} (t^2 - 1)^s \frac{\xi \eta}{(t^2 + \xi^2)(t^2 + \eta^2)} dt d\xi d\eta \\ &\leq \frac{\pi}{2 \cos(s\pi)} \int_0^{\infty} \int_0^{\infty} |\widehat{f}(\xi)| \cdot |\overline{\widehat{f}(\eta)}| \cdot \xi \eta \frac{(1 + \xi^2)^{\frac{2s-1}{2}} - (1 + \eta^2)^{\frac{2s-1}{2}}}{\eta^2 - \xi^2} d\xi d\eta \end{aligned}$$

$$\begin{aligned}
 &= \frac{\pi}{2\cos(s\pi)} \int_0^\infty \int_0^\infty (1+\xi^2)^{\frac{s}{2}} |\widehat{f}(\xi)| \cdot (1+\eta^2)^{\frac{s}{2}} |\overline{\widehat{f}(\eta)}| \\
 &\quad \times \frac{\xi\eta}{\eta^2 - \xi^2} \cdot \frac{(1+\xi^2)^{\frac{2s-1}{2}} - (1+\eta^2)^{\frac{2s-1}{2}}}{(1+\xi^2)^{\frac{s}{2}}(1+\eta^2)^{\frac{s}{2}}} d\xi d\eta.
 \end{aligned} \tag{3.17}$$

It is elementary to prove that for any $\xi, \eta > 0, \xi \neq \eta$ we have

$$\frac{\xi\eta}{\eta^2 - \xi^2} \cdot \frac{(1+\xi^2)^{\frac{2s-1}{2}} - (1+\eta^2)^{\frac{2s-1}{2}}}{(1+\xi^2)^{\frac{s}{2}}(1+\eta^2)^{\frac{s}{2}}} \leq \frac{\xi^s \eta^{1-s} - \xi^{1-s} \eta^s}{\eta^2 - \xi^2} = K_s(\xi, \eta). \tag{3.18}$$

Coming back to (3.17) and using Lemma 3.3 we obtain

$$\begin{aligned}
 \int_1^\infty (t^2 - 1)^s \left| \int_0^\infty \widehat{f}(\xi) \frac{\xi}{t^2 + \xi^2} d\xi \right|^2 dt &\leq \frac{\pi C(s)}{2\cos(s\pi)} \|(1 + |\cdot|^2)^{\frac{s}{2}} \widehat{f}\|_{L^2(0,\infty)}^2 \\
 &\leq C'(s) \|f\|_{H^s}^2.
 \end{aligned} \tag{3.19}$$

From (3.4) and (3.15) if $s \in [0, \frac{3}{2})$, respectively from (3.4) and (3.19) if $s \in (-\frac{1}{2}, 0)$, we infer that T_1 and T_2 can be extended as linear continuous operators from $H^s(\mathbf{R})$ to $H^s(\mathbf{R})$.

Now we prove Theorem 3.2 in the case $N \geq 2$.

If $s \in [0, \frac{3}{2})$, arguing as in (3.5)–(3.7) and using Lemma 3.3 we have

$$\begin{aligned}
 &\int_{|\xi'|}^\infty (t^2 - |\xi'|^2)^s \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt \\
 &\leq \int_0^\infty t^{2s} \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt \\
 &\leq \int_0^\infty \int_0^\infty |\widehat{f}(\xi_1, \xi')| \xi_1^s \cdot |\overline{\widehat{f}(\eta_1, \xi')}| \eta_1^s \cdot (\xi_1^{-s} \eta_1^{-s} I_s(\xi_1, \eta_1)) d\xi_1 d\eta_1 \\
 &\leq C(s) \| |\cdot|^s \widehat{f}(\cdot, \xi') \|_{L^2(0,\infty)}^2 \leq C(s) \int_{-\infty}^\infty (\xi_1^2 + |\xi'|^2)^s |\widehat{f}(\xi_1, \xi')|^2 d\xi_1.
 \end{aligned} \tag{3.20}$$

If $s \in (-\frac{1}{2}, 0)$, using the change of variable $\tau = \sqrt{t^2 - |\xi'|^2}$, arguing as in the proof of (3.16), then taking (3.9) into account we obtain

$$\int_{|\xi'|}^\infty \frac{(t^2 - |\xi'|^2)^s}{(t^2 + \xi^2)(t^2 + \eta^2)} dt = \int_0^\infty \frac{\tau^{2s}}{(\tau^2 + |\xi'|^2 + \xi_1^2)(\tau^2 + |\xi'|^2 + \eta_1^2)} \cdot \frac{\tau}{\sqrt{\tau^2 + |\xi'|^2}} d\tau$$

$$\begin{aligned}
&\leq \int_0^\infty \frac{\tau^{2s}}{(\tau^2 + |\xi'|^2 + \xi_1^2)(\tau^2 + |\xi'|^2 + \eta_1^2)} d\tau \\
&= \frac{\pi}{2\cos(s\pi)} \cdot \frac{(|\xi'|^2 + \xi_1^2)^{\frac{2s-1}{2}} - (|\xi'|^2 + \eta_1^2)^{\frac{2s-1}{2}}}{\eta_1^2 - \xi_1^2}.
\end{aligned}$$

We also have

$$\frac{\xi_1 \eta_1}{\eta_1^2 - \xi_1^2} \cdot \frac{(\xi_1^2 + |\xi'|^2)^{\frac{2s-1}{2}} - (\eta_1^2 + |\xi'|^2)^{\frac{2s-1}{2}}}{(\xi_1^2 + |\xi'|^2)^{\frac{s}{2}} (\eta_1^2 + |\xi'|^2)^{\frac{s}{2}}} \leq K_s(\xi_1, \eta_1)$$

(this inequality is analogous to (3.18)). Arguing as in (3.17), using the two previous inequalities and Lemma 3.3 we get

$$\begin{aligned}
&\int_{|\xi'|}^\infty (t^2 - |\xi'|^2)^s \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right|^2 dt \\
&\leq \frac{\pi C(s)}{2\cos(s\pi)} \left\| (|\xi'|^2 + |\cdot|^2)^{\frac{s}{2}} \widehat{f}(\cdot, \xi') \right\|_{L^2(0, \infty)}^2 \\
&\leq C'(s) \int_{-\infty}^\infty (\xi_1^2 + |\xi'|^2)^s |\widehat{f}(\xi_1, \xi')|^2 d\xi_1. \tag{3.21}
\end{aligned}$$

Integrating (3.20), respectively (3.21), over $\mathbf{R}^{N-1}$ we infer that the integral in the right-hand side of (3.3) is less than $C''(s) \|f\|_{\dot{H}^s}^2$. This proves that T_1 and T_2 can be extended by continuity from $\dot{H}^s(\mathbf{R}^N)$ to $\dot{H}^s(\mathbf{R}^N)$ for $s \in (-\frac{1}{2}, \frac{3}{2})$.

In a similar way we show that T_1 and T_2 can be extended by continuity from $H^s(\mathbf{R}^N)$ to $H^s(\mathbf{R}^N)$ for $s \in (-\frac{1}{2}, \frac{3}{2})$. Theorem 3.2 is now proved. $\square$

For a measurable function u defined on $\mathbf{R}^N$, we define its antisymmetric part in the x_1 direction by $Au(x_1, x') = \frac{1}{2}(u(x_1, x') - u(-x_1, x'))$. If u is a tempered distribution, we define Au by $\langle Au, \phi \rangle_{\mathcal{S}', \mathcal{S}} = \langle u, A\phi \rangle_{\mathcal{S}', \mathcal{S}}$ for any $\phi \in \mathcal{S}$. Obviously, Au is odd with respect to x_1 (for distributions, this means that $\langle Au, \phi(-x_1, x') \rangle_{\mathcal{S}', \mathcal{S}} = -\langle Au, \phi \rangle_{\mathcal{S}', \mathcal{S}}$). It is clear from the definition that A defines a linear continuous map from $H^s(\mathbf{R}^N)$ to $H^s(\mathbf{R}^N)$ (respectively from $\dot{H}^s(\mathbf{R}^N)$ to $\dot{H}^s(\mathbf{R}^N)$) for any s . Moreover, for any tempered distribution u , the distribution $\mathcal{F}(Au)$ is odd with respect to x_1 .

It follows from the proof of Theorem 3.2 that for any $s \in (-\frac{1}{2}, \frac{3}{2})$, the following complex bilinear forms are continuous:

$$B_{N,s} : \dot{H}^s(\mathbf{R}^N) \times \dot{H}^s(\mathbf{R}^N) \rightarrow \mathbf{C},$$

$$B_{N,s}(u, v) = \int_{\mathbf{R}^{N-1}} \int_{|\xi'|}^{\infty} (t^2 - |\xi'|^2)^s \int_0^{\infty} \widehat{Au}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \\ \times \int_0^{\infty} \overline{\widehat{Av}(\eta_1, \xi')} \frac{\eta_1}{t^2 + \eta_1^2} d\eta_1 dt d\xi',$$

$$\widetilde{B}_{N,s} : H^s(\mathbf{R}^N) \times H^s(\mathbf{R}^N) \rightarrow \mathbf{C},$$

$$\widetilde{B}_{N,s}(u, v) = \int_{\mathbf{R}^{N-1}} \int_{\sqrt{|\xi'|^2+1}}^{\infty} (t^2 - |\xi'|^2 - 1)^s \int_0^{\infty} \widehat{Au}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \\ \times \int_0^{\infty} \overline{\widehat{Av}(\eta_1, \xi')} \frac{\eta_1}{t^2 + \eta_1^2} d\eta_1 dt d\xi'.$$

Moreover, from (3.3) and (3.4) we have the identities

$$\|T_1 u\|_{\dot{H}^s(\mathbf{R}^N)}^2 + \|T_2 u\|_{\dot{H}^s(\mathbf{R}^N)}^2 - 2\|u\|_{\dot{H}^s(\mathbf{R}^N)}^2 = -\frac{16 \sin(s\pi)}{\pi^2} B_{N,s}(Au, Au), \quad (3.22)$$

$$\|T_1 u\|_{H^s(\mathbf{R}^N)}^2 + \|T_2 u\|_{H^s(\mathbf{R}^N)}^2 - 2\|u\|_{H^s(\mathbf{R}^N)}^2 = -\frac{16 \sin(s\pi)}{\pi^2} \widetilde{B}_{N,s}(Au, Au) \quad (3.23)$$

for any $u \in C_c^\infty(\mathbf{R}^N)$. From Theorem 3.2, the continuity of $B_{N,s}$ and of $\widetilde{B}_{N,s}$ and the density of $C_c^\infty(\mathbf{R}^N)$ in $\dot{H}^s(\mathbf{R}^N)$ and in $H^s(\mathbf{R}^N)$ we infer that we have the following.

Corollary 3.4. *Let $s \in (-\frac{1}{2}, \frac{3}{2})$. The identity (3.22) holds for any $u \in \dot{H}^s(\mathbf{R}^N)$ and (3.23) holds for any $u \in H^s(\mathbf{R}^N)$.*

Our next aim is to show that the quadratic forms $B_{N,s}$ and $\widetilde{B}_{N,s}$ define norms in some spaces of odd functions. We start with the following proposition.

Lemma 3.5. *Assume that $g : \mathbf{R} \rightarrow \mathbf{C}$ is a measurable function, $g(-t) = -g(t)$ a.e. and*

- *either $g \in L^p(\mathbf{R})$ for some $p \in (1, \infty)$,*
- *or $(k^2 + \xi^2)^{\frac{s}{2}} g(\xi) \in L^2(\mathbf{R})$ for some $k \in \mathbf{R}$ and $s \in (-\frac{1}{2}, \frac{3}{2})$.*

Suppose that the set

$$A = \left\{ x > 0 \mid \int_0^{\infty} \frac{\xi}{x^2 + \xi^2} g(\xi) d\xi = 0 \right\}$$

has a limit point $x_0 > 0$.

Then $g = 0$ almost everywhere on $\mathbf{R}$.

Proof. We may suppose without loss of generality that g is real (otherwise we carry out the proof for its real and imaginary parts).

First we deal with the simpler case $g \in L^p(\mathbf{R})$ for some p , $1 < p < \infty$. We define the Poisson integrals for g ,

$$\begin{aligned} a(x, y) &= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{x}{x^2 + (y-t)^2} g(t) dt \quad \text{and} \\ b(x, y) &= -\frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y-t}{x^2 + (y-t)^2} g(t) dt. \end{aligned} \quad (3.24)$$

It follows from Lemma 2.4(iii) that the functions a and b are well defined and harmonic in the right half-plane and $r(x+iy) := a(x, y) + ib(x, y)$ is holomorphic in $\{z \in \mathbf{C} \mid \operatorname{Re}(z) > 0\}$. Since g is odd, we have $a(x, 0) = 0$ for any $x > 0$. If $x \in A$, we have also $b(x, 0) = 0$. Consequently, $r(x) = 0$ for any $x \in A$. But r is holomorphic and A has a limit point $x_0 > 0$, thus necessarily $r \equiv 0$. By Lemma 2.4(ii) we know that $a(x, y) \rightarrow g(y)$ as $x \rightarrow 0$ for almost every y , hence $g = 0$ a.e. on $\mathbf{R}$.

Suppose that $(k^2 + |\cdot|^2)^{\frac{s}{2}} g \in L^2(\mathbf{R})$ for some $k \neq 0$ and $s \in (-\frac{1}{2}, \frac{3}{2})$. We may assume that $k = 1$. If $s \in [0, \frac{3}{2})$, then obviously $g \in L^2(\mathbf{R})$ and the conclusion of the lemma follows from the above considerations. If $s \in (-\frac{1}{2}, 0)$, then for any $x > 0$ and $y \in \mathbf{R}$ the functions $\varphi_{x,y}(t) = (1+t^2)^{-\frac{s}{2}} \frac{x}{x^2+(y-t)^2}$ and $\psi_{x,y}(t) = (1+t^2)^{-\frac{s}{2}} \frac{y-t}{x^2+(y-t)^2}$ belong to $L^2(\mathbf{R})$. We may write

$$\int_{-\infty}^{\infty} \frac{x}{x^2 + (y-t)^2} g(t) dt = \int_{-\infty}^{\infty} \varphi_{x,y}(t) (1+t^2)^{\frac{s}{2}} g(t) dt$$

and

$$\int_{-\infty}^{\infty} \frac{y-t}{x^2 + (y-t)^2} g(t) dt = \int_{-\infty}^{\infty} \psi_{x,y}(t) (1+t^2)^{\frac{s}{2}} g(t) dt.$$

Using the Cauchy–Schwarz inequality, we see that the functions a and b in (3.24) are well defined in the right half-plane (in particular, $\int_0^\infty \frac{\xi}{x^2+\xi^2} g(\xi) d\xi$ exists for any $x > 0$). Clearly the function $r(x+iy) := a(x, y) + ib(x, y)$ is holomorphic and, as above we have $r(x) = 0$ for $x \in A$. Since A has a limit point $x_0 > 0$, we infer that $r \equiv 0$. Next, we have $\lim_{x \downarrow 0} a(x, y) = g(y)$ whenever y is a Lebesgue point of g (the proof of this fact follows from standard arguments and it is quite similar to the proof of [24, Theorem 1.25, p. 15]; for brevity, we omit it). This obviously implies $g = 0$ a.e., as desired.

Now let us consider the case $k = 0$. If $|\cdot|^s g \in L^2(\mathbf{R})$ and $s \in (-\frac{1}{2}, \frac{1}{2})$, we may repeat almost word by word the proof above (we have only to replace the functions $\varphi_{x,y}$ and $\psi_{x,y}$ by $t \mapsto t^{-s} \frac{x}{x^2+(y-t)^2}$, respectively by $t \mapsto t^{-s} \frac{y-t}{x^2+(y-t)^2}$).

If $|\cdot|^s g \in L^2(\mathbf{R})$ and $s \in [\frac{1}{2}, \frac{3}{2})$, the integrals defining a and b in (3.24) do not necessarily converge. In this case we define

$$\begin{aligned}
 a_1(x, y) &= \frac{1}{\pi} \int_0^\infty \frac{4xyt}{[x^2 + (y-t)^2][x^2 + (y+t)^2]} g(t) dt \quad \text{and} \\
 b_1(x, y) &= \frac{1}{\pi} \int_0^\infty \frac{2t(t^2 + x^2 - y^2)}{[x^2 + (y-t)^2][x^2 + (y+t)^2]} g(t) dt.
 \end{aligned} \tag{3.25}$$

Notice that if $g \in L^1_{\text{loc}}(\mathbf{R})$ is odd and $\frac{g(t)}{t} \in L^1([1, \infty))$, then $a = a_1$ and $b = b_1$. It is obvious that for fixed $x > 0$, $y \in \mathbf{R}$ and $s \in (-\frac{1}{2}, \frac{3}{2})$, the functions $\varphi_1(t) = t^{-s} \frac{4xyt}{[x^2 + (y-t)^2][x^2 + (y+t)^2]}$ and $\psi_1(t) = t^{-s} \frac{2t(t^2 + x^2 - y^2)}{[x^2 + (y-t)^2][x^2 + (y+t)^2]}$ belong to $L^2((0, \infty))$ and this implies that a_1 and b_1 are well defined. It is straightforward that $r_1(x + iy) := a_1(x, y) + ib_1(x, y)$ is holomorphic in the right half-plane. Obviously $a_1(x, 0) = 0$ for any $x > 0$ and $b_1(x, 0) = \frac{2}{\pi} \int_0^\infty \frac{t}{x^2 + t^2} g(t) dt = 0$ for $x \in A$. Consequently $r = 0$ on A . Since A has a limit point $x_0 > 0$, we infer that $r \equiv 0$ in the right half-plane. Let $y > 0$ be a Lebesgue point of g . Since

$$\int_0^\infty \frac{4xyt}{[x^2 + (y-t)^2][x^2 + (y+t)^2]} dt = 2 \arctan \frac{y}{x},$$

proceeding as in the previous cases, one can show that $|a_1(x, y) - \frac{2}{\pi}(\arctan \frac{y}{x})g(y)| \rightarrow 0$ as $x \rightarrow 0$, hence $\lim_{x \downarrow 0} a_1(x, y) = g(y)$. Consequently we have $\lim_{x \downarrow 0} a_1(x, y) = g(y)$ for almost every y and the lemma is proved. $\square$

We set

$$\begin{aligned}
 H^s_{1,\text{odd}}(\mathbf{R}^N) &= \{f \in H^s(\mathbf{R}^N) \mid f \text{ is odd with respect to } x_1\} = \{f \in H^s(\mathbf{R}^N) \mid f = Af\}, \\
 \dot{H}^s_{1,\text{odd}}(\mathbf{R}^N) &= \{f \in \dot{H}^s(\mathbf{R}^N) \mid f \text{ is odd with respect to } x_1\} = \{f \in \dot{H}^s(\mathbf{R}^N) \mid f = Af\},
 \end{aligned}$$

where, as before, Af is the antisymmetric part of f in the x_1 direction. For $f \in \dot{H}^s_{1,\text{odd}}(\mathbf{R}^N)$ we define $N_s(f) = (B_{N,s}(f, f))^{\frac{1}{2}}$ and for $f \in H^s_{1,\text{odd}}(\mathbf{R}^N)$ we define $\tilde{N}_s(f) = (\tilde{B}_{N,s}(f, f))^{\frac{1}{2}}$.

Theorem 3.6. $\tilde{N}_s$ is a norm on $H^s_{1,\text{odd}}(\mathbf{R}^N)$, continuous with respect to the usual H^s norm, and N_s is a norm on $\dot{H}^s_{1,\text{odd}}(\mathbf{R}^N)$, continuous with respect to the $\dot{H}^s$ norm.

Endowed with these norms, $H^s_{1,\text{odd}}(\mathbf{R}^N)$ and $\dot{H}^s_{1,\text{odd}}(\mathbf{R}^N)$ are pre-Hilbert spaces.

Proof. It is clear that $\tilde{B}_{N,s}$ and $B_{N,s}$ are complex-symmetric bilinear forms on $H^s(\mathbf{R}^N)$ (respectively on $\dot{H}^s(\mathbf{R}^N)$) and that $\tilde{B}_{N,s}(f, f) \geq 0$ and $B_{N,s}(f, f) \geq 0$ for any f (thus, in particular, $\tilde{N}_s$ and N_s are well defined). Suppose, for instance, that $\hat{f} \in H^s_{1,\text{odd}}(\mathbf{R}^N)$ and $\tilde{B}_{N,s}(f, f) = 0$. This implies that for almost every $\xi' \in \mathbf{R}^{N-1}$ we have $\hat{f}(\cdot, \xi') = -\hat{f}(\cdot, \xi')$ a.e., $(|\cdot|^2 + |\xi'|^2)^{\frac{s}{2}} \hat{f}(\cdot, \xi') \in L^2(\mathbf{R})$ and

$$\int_{\sqrt{|\xi'|^2+1}}^{\infty} (t^2 - |\xi'|^2 - 1)^s \left| \int_0^{\infty} \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 \right| dt = 0.$$

For such ξ' we must have

$$\int_0^{\infty} \widehat{f}(\xi_1, \xi') \frac{\xi_1}{t^2 + \xi_1^2} d\xi_1 = 0 \quad \text{for almost every } t \in (\sqrt{|\xi'|^2 + 1}, \infty)$$

and using Lemma 3.5 we infer that $\widehat{f}(\cdot, \xi') = 0$ a.e. on $\mathbf{R}$, so $\int_{\mathbf{R}} (\xi_1^2 + |\xi'|^2)^s |\widehat{f}(\xi_1, \xi')|^2 d\xi_1 = 0$. Consequently

$$\|f\|_{H^s}^2 = \int_{\mathbf{R}^{N-1}} \int_{\mathbf{R}} (\xi_1^2 + |\xi'|^2)^s |\widehat{f}(\xi_1, \xi')|^2 d\xi_1 d\xi' = 0,$$

i.e. $f = 0$ a.e. The proof is the same for $f \in \dot{H}^s(\mathbf{R}^N)$. Finally, the continuity of $\widetilde{N}_s$ and N_s with respect to the usual norms follows from Theorem 3.2 and Corollary 3.4. $\square$

4. Applications

In this section we illustrate how the results in Sections 2 and 3 can be used to prove the symmetry of minimizers in some concrete examples.

4.1. Problems involving fractional powers of the Laplace operator

Theorem 4.1. *Let $s \in (0, 1)$ and assume that $F, G : \mathbf{R} \rightarrow \mathbf{R}$ are such that $u \mapsto F(u)$ and $u \mapsto G(u)$ map $\dot{H}^s(\mathbf{R}^N)$ (or $H^s(\mathbf{R}^N)$) into $L^1(\mathbf{R}^N)$. Suppose that either:*

Case A. $u \in \dot{H}^s(\mathbf{R}^N)$ and u is a solution of the minimization problem

$$\text{minimize } E(u) := \int_{\mathbf{R}^N} |\xi|^{2s} |\widehat{u}(\xi)|^2 d\xi + \int_{\mathbf{R}^N} F(u(x)) dx$$

$$\text{under the constraint } I(u) = \int_{\mathbf{R}^N} G(u(x)) dx = \lambda \neq 0, \quad \text{or}$$

Case B. $u \in H^s(\mathbf{R}^N)$ and u is a solution of the minimization problem

$$\text{minimize } E(u) := \int_{\mathbf{R}^N} (1 + |\xi|^2)^s |\widehat{u}(\xi)|^2 d\xi + \int_{\mathbf{R}^N} F(u(x)) dx$$

$$\text{under the constraint } I(u) = \int_{\mathbf{R}^N} G(u(x)) dx = \lambda \neq 0.$$

Then, after a translation in $\mathbf{R}^N$, u is radially symmetric.

Proof. Let us prove first that u is symmetric with respect to x_1 . Making a translation in the x_1 direction if necessary, we may assume that $\int_{\{x_1 < 0\}} G(u(x)) dx = \int_{\{x_1 > 0\}} G(u(x)) dx = \frac{\lambda}{2}$. Let $u_1 = T_1 u$ and $u_2 = T_2 u$. It follows from Theorem 3.2 that $u_1, u_2 \in \dot{H}^s(\mathbf{R}^N)$ in case A, respectively $u_1, u_2 \in H^s(\mathbf{R}^N)$ in case B. It is obvious that we have $\int_{\mathbf{R}^N} G(u_1(x)) dx = 2 \int_{\{x_1 < 0\}} G(u(x)) dx = \lambda$ and $\int_{\mathbf{R}^N} G(u_2(x)) dx = 2 \int_{\{x_1 > 0\}} G(u(x)) dx = \lambda$; hence u_1 and u_2 also satisfy the constraint. From (3.22) and (3.23) we have

$$E(u_1) + E(u_2) - 2E(u) = -\frac{16 \sin(s\pi)}{\pi^2} N_s^2(Au) \quad \text{in case A, respectively}$$

$$E(u_1) + E(u_2) - 2E(u) = -\frac{16 \sin(s\pi)}{\pi^2} \tilde{N}_s^2(Au) \quad \text{in case B,}$$

where, as before, $Au(x_1, x') = \frac{1}{2}(u(x_1, x') - u(-x_1, x'))$ is the antisymmetric part of u in the x_1 direction. If $Au \neq 0$, then Theorem 3.6 implies $N_s^2(Au) > 0$ (respectively $\tilde{N}_s^2(Au) > 0$) and we infer that $E(u_1) + E(u_2) - 2E(u) < 0$, contradicting the fact that u is a minimizer. Thus necessarily $Au \equiv 0$ and this means that u is symmetric with respect to x_1 .

Arguing similarly with the remaining variables $x_2, \dots, x_N$, we find a new origin O' such that u is symmetric with respect to any of the variables $x_1, \dots, x_N$; in particular, $u(-x) = u(x)$ a.e. on $\mathbf{R}^N$. Now let Π be any hyperplane containing the new origin O' and let Π_+ and Π_- be the halfspaces determined by Π . Since the transformation $x \mapsto -x$ maps Π_- into Π_+ , we see that $\int_{\Pi_-} G(u(x)) dx = \int_{\Pi_+} G(u(x)) dx = \frac{\lambda}{2}$. Arguing as above we conclude that u must be symmetric with respect to Π . This implies that u is radially symmetric with respect to the new origin O' . $\square$

An application of Theorem 4.1 concerns the solitary waves to the generalized Benjamin–Ono equation

$$A_t + \alpha A A_x - \beta (-\Delta)^{\frac{1}{2}} A_x = 0, \quad (x, y) \in \mathbf{R}^2, \quad t \in \mathbf{R},$$

where $\alpha, \beta > 0$. Solitary waves are solutions of the form $A(t, x, y) = u(x - ct, y)$. After a scale change, a solitary wave $u(x, y)$ satisfies the equation

$$u + (-\Delta)^{\frac{1}{2}} u = u^2 \quad \text{in } \mathbf{R}^2.$$

The existence of solitary waves was proved in [21] by minimizing the functional

$$V(u) = \frac{1}{2} \int_{\mathbf{R}^2} |(-\Delta)^{\frac{1}{4}} u|^2 dx + \int_{\mathbf{R}^2} u^2 dx = \frac{1}{2(2\pi)^2} \int_{\mathbf{R}^2} |\xi| |\widehat{u}(\xi)|^2 d\xi + \int_{\mathbf{R}^2} u^2 dx$$

under the constraint $I(u) = \frac{1}{3} \int_{\mathbf{R}^2} u^3 dx = \text{constant}$. It has been shown in [21] that any solution u_* of the above problem also minimizes

$$E(v) := \frac{1}{2} \int_{\mathbf{R}^2} |(-\Delta)^{\frac{1}{4}} v|^2 dx - \frac{1}{3} \int_{\mathbf{R}^2} v^3 dx$$

under the constraint $Q(v) = Q(u_*)$, where $Q(v) = \frac{1}{2} \int_{\mathbf{R}^2} |u|^2 dx$.

It follows directly from Theorem 4.1 that, except for translation, any minimizer of these problems is radially symmetric.

Next we apply our method to a variational problem involving two unknown functions (the vector case). Consider the functionals

$$E(u, v) = \frac{1}{2} \int_{\mathbf{R}^N} (|(-\Delta)^{\frac{s}{2}} u|^2 + |\nabla v|^2) dx + \int_{\mathbf{R}^N} F(u, v) dx$$

where $0 < s < 1$, and

$$Q(u, v) = \int_{\mathbf{R}^N} G(u, v) dx.$$

We make the following assumptions:

A1. $F, G: \mathbf{R}^2 \rightarrow \mathbf{R}$ are C^2 functions satisfying $F(0, 0) = \partial_1 F(0, 0) = \partial_2 F(0, 0) = 0$, $G(0, 0) = \partial_1 G(0, 0) = \partial_2 G(0, 0) = 0$ and the growth conditions

$$|\partial_i F(u, v)| \leq C(|u|^{p-1} + |v|^{q-1}) \quad \text{and} \quad |\partial_i G(u, v)| \leq C(|u|^{p-1} + |v|^{q-1}) \quad \text{if } |(u, v)| \geq 1,$$

where $i \in \{1, 2\}$, C is a positive constant, $2 < p < \frac{2N}{N-2s}$ and $2 < q < \frac{2N}{N-2}$.

A2. If $(u, v) \in H^s(\mathbf{R}^N) \times H^1(\mathbf{R}^N)$ and $(u, v) \neq (0, 0)$, then either $\partial_1 G(u, v) \neq 0$ or $\partial_2 G(u, v) \neq 0$ (a manifold condition).

Theorem 4.2. Under assumptions A1 and A2, any minimizer $(u, v) \in H^s(\mathbf{R}^N) \times H^1(\mathbf{R}^N)$ of $E(u, v)$ subject to the constraint $Q(u, v) = \lambda \neq 0$ is radially symmetric (except for translation).

Proof. First we prove that after a translation, (u, v) is symmetric with respect to x_1 . In fact, after possibly a translation in the x_1 direction we may assume that

$$\int_{\{x_1 < 0\}} G(u, v) dx = \int_{\{x_1 > 0\}} G(u, v) dx = \frac{\lambda}{2}. \quad (4.1)$$

We put $u_1 = T_1 u$, $u_2 = T_2 u$, $v_1 = T_1 v$ and $v_2 = T_2 v$. By Theorem 3.2, the pairs (u_1, v_1) and (u_2, v_2) belong to $H^s(\mathbf{R}^N) \times H^1(\mathbf{R}^N)$ and in view of (4.1) they also satisfy the constraint $Q(u_1, v_1) = Q(u_2, v_2) = \lambda$. Moreover, defining $W(\varphi) = \int_{\mathbf{R}^N} |\xi|^{2s} |\widehat{\varphi}(\xi)|^2 d\xi$ and using (3.22) we see that

$$\begin{aligned} E(u_1, v_1) + E(u_2, v_2) - 2E(u, v) &= \frac{1}{2} \frac{1}{(2\pi)^N} (W(u_1) + W(u_2) - 2W(u)) \\ &= -\frac{1}{(2\pi)^N} \frac{8 \sin(s\pi)}{\pi^2} B_{N,s}(Au, Au) \leq 0. \end{aligned}$$

We conclude that (u_1, v_1) and (u_2, v_2) are also minimizers and we must have $B_{N,s}(Au, Au) = 0$. Theorem 3.6 implies that $Au = 0$, that is u is symmetric with respect to x_1 , i.e. $u = u_1 = u_2$.

Since (u, v) and $(u_1, v_1) = (u, v_1)$ are minimizers, they satisfy the Euler–Lagrange equations

$$\begin{cases} (-\Delta)^s u + \partial_1 F(u, v) + \alpha \partial_1 G(u, v) = 0, \\ -\Delta v + \partial_2 F(u, v) + \alpha \partial_2 G(u, v) = 0, \end{cases} \quad (4.2)$$

respectively

$$\begin{cases} (-\Delta)^s u + \partial_1 F(u, v_1) + \beta \partial_1 G(u, v_1) = 0, \\ -\Delta v_1 + \partial_2 F(u, v_1) + \beta \partial_2 G(u, v_1) = 0. \end{cases} \quad (4.3)$$

From (4.2), A1, the elliptic regularity for the Laplacian and its fractional powers and the usual boot-strap argument we get $u \in H^{2s}(\mathbf{R}^N) \cap L^\infty(\mathbf{R}^N)$ and $v \in H^2(\mathbf{R}^N) \cap L^\infty(\mathbf{R}^N)$. Of course that the same conclusion holds for (u, v_1) . Notice that the L^p elliptic regularity for fractional powers of the Laplacian and for $1 < p < \infty$ follows from the fact that the multiplier $m(\xi) = \frac{(1+|\xi|^2)^s}{1+|\xi|^{2s}}$ satisfies the estimate $|D^\alpha m(\xi)| \leq \frac{B(\alpha)}{|\xi|^\alpha}$ and from the theorem of Mihlin–Hörmander.

We recall the following well-known result.

Unique Continuation Principle. Assume that $\Phi \in H^2(\mathbf{R}^N, \mathbf{R}^m)$ solves the linear system

$$-\Delta \Phi + A(x)\Phi(x) = 0 \quad \text{in } \mathbf{R}^N, \quad (4.4)$$

where $A(x)$ is an $m \times m$ matrix whose elements belong to $L^\infty(\mathbf{R}^N)$. If $\Phi \equiv 0$ in some open set $\omega \subset \mathbf{R}^N$, then $\Phi \equiv 0$ in $\mathbf{R}^N$.

A proof for the Unique Continuation Principle is given in [13, Chapter VIII] in the scalar case and in the appendix of [18] in the vector case. Notice that the Unique Continuation Principle is essentially a local result. Although it is stated for functions $\Phi \in H^2(\mathbf{R}^N)$, it is also valid for functions $\Phi \in W^{2,p}(\mathbf{R}^N)$ with $p > 2$ because $W_{\text{loc}}^{2,p}(\mathbf{R}^N) \subset H_{\text{loc}}^2(\mathbf{R}^N)$. This observation will be useful later.

Now let us come back to the proof of Theorem 4.2.

If $(u_1, v_1) = (0, 0)$, since $G(0, 0) = 0$ we have $\lambda = Q(u_1, v_1) = 0$, a contradiction. Thus $(u_1, v_1) \neq (0, 0)$ and it follows from A2 that there exists $(x_1, x') \in (-\infty, 0) \times \mathbf{R}^{N-1}$ such that $\partial_1 G(u_1, v_1)(x_1, x') \neq 0$ or $\partial_2 G(u_1, v_1)(x_1, x') \neq 0$. Since $v = v_1$ for $x_1 < 0$, we infer from (4.2) and (4.3) that $\alpha = \beta$. Moreover, using the regularity of u, v, v_1 we get $\partial_2 F(u, v) - \partial_2 F(u, v_1) = b(x)(v(x) - v_1(x))$ and $\partial_2 G(u, v) - \partial_2 G(u, v_1) = c(x)(v(x) - v_1(x))$ where $b, c \in L^\infty(\mathbf{R}^N)$. Let $w(x) = v(x) - v_1(x)$. Using the second components of (4.2) and (4.3) and the fact that $\alpha = \beta$, we see that w satisfies the linear equation $-\Delta w(x) + a(x)w(x) = 0$ in $\mathbf{R}^N$, where $a = b + \alpha c \in L^\infty(\mathbf{R}^N)$. Since w vanishes on a half-space, by the Unique Continuation Principle we conclude that w vanishes everywhere, and this implies $v = v_1$ in $\mathbf{R}^N$. Thus we have shown that (u, v) is symmetric with respect to x_1 .

Repeating this argument with the variables $x_2, \dots, x_N$, we find a new origin O' such that (u, v) is symmetric with respect to $x_1, \dots, x_N$. Then as in the proof of Theorem 4.1 we show that (u, v) is symmetric with respect to any hyperplane Π containing O' , consequently (u, v) is radially symmetric with respect to the new origin O' . $\square$

Remark 4.3. Symmetrization inequalities for functions in the space $H^{1/2}(\mathbf{R}^N)$ have been proved in [3]. Therefore if $s = \frac{1}{2}$, the function F in Theorem 4.2 satisfies the cooperative condition

$\partial_{1,2}^2 F(u, v) \leq 0$ (see [5]), G has a special form and it is known in advance that the components u, v of the minimizer are nonnegative, then using symmetrization one can conclude that *there exists* a radially symmetric minimizer.

Remark 4.4. In the case $F(u, v) = u^2 + v^2$, $G(u, v) = u^2 v$, by using symmetrization and Riesz' inequality it has been proved in [3] that *there exists* a radially symmetric minimizer. The fact that F and G are homogeneous plays a crucial role in their proof.

As an example of application for Theorem 4.2, we consider the Hamiltonian system

$$\begin{cases} \frac{\partial u}{\partial t} = \frac{\partial}{\partial x_1} ((-\Delta)^{1/2} u + \partial_1 F(u, v)), \\ \frac{\partial v}{\partial t} = \frac{\partial}{\partial x_1} (-\Delta v + \partial_2 F(u, v)). \end{cases} \quad (4.5)$$

The generalized multidimensional Benjamin–Ono equation

$$\frac{\partial u}{\partial t} = \frac{\partial}{\partial x_1} ((-\Delta)^{1/2} u + g(u)) \quad (4.6)$$

with $g(u) = u^2$ and the generalized multidimensional Korteweg–de Vries equation

$$\frac{\partial v}{\partial t} = \frac{\partial}{\partial x_1} (-\Delta v + f(v)) \quad (4.7)$$

have been considered in [21] and in [4], respectively; in these papers, references giving the physical motivation for the above equations can also be found. System (4.5) can be considered a Hamiltonian coupling between (4.6) and (4.7).

Formally, the system (4.5) has the following conserved quantities:

$$\begin{aligned} E(u, v) &= \frac{1}{2} \int_{\mathbf{R}^N} |(-\Delta)^{1/4} u|^2 + |\nabla v|^2 dx + \int_{\mathbf{R}^N} F(u, v) dx \quad \text{and} \\ Q(u, v) &= \frac{1}{2} \int_{\mathbf{R}^N} (u^2 + v^2) dx. \end{aligned}$$

If we minimize $E(u, v)$ subject to the constraint $Q(u, v) = \lambda$, where $\lambda > 0$, then according to [9] the set S_λ containing the elements of $H^{\frac{1}{2}}(\mathbf{R}^N) \times H^1(\mathbf{R}^N)$ where the minimum is achieved is invariant and orbitally stable with respect to (4.5). Since any element $(\phi, \psi) \in S_\lambda$ satisfies the Euler–Lagrange system

$$\begin{cases} (-\Delta)^{1/2} \phi + \partial_1 F(\phi, \psi) + c\phi = 0, \\ -\Delta \psi + \partial_2 F(\phi, \psi) + c\psi = 0, \end{cases}$$

we see that (ϕ, ψ) gives rise to a travelling wave solution of (4.5) of the form $(u(t, x), v(t, x)) = (\phi(x_1 - ct, x'), \psi(x_1 - ct, x'))$, $x' \in \mathbf{R}^{N-1}$. As a consequence of Theorem 4.2, the elements (ϕ, ψ) obtained in this way are radially symmetric (after a translation).

4.2. Minimizers of the generalized Choquard functional

In this paragraph we consider the problem of minimizing the generalized Choquard functional

$$E(u) = \frac{1}{2} \int_{\mathbf{R}^N} |\nabla u|^2 dx - \int_{\mathbf{R}^N} \int_{\mathbf{R}^N} F(u(x)) \frac{1}{|x-y|^{N-2}} F(u(y)) dx dy + \int_{\mathbf{R}^N} H(u(x)) dx \quad (4.8)$$

subject to the constraint $Q(u) = \int_{\mathbf{R}^N} G(u(x)) dx = \text{constant} \neq 0$.

It is worth to note that the complex version of E ,

$$\begin{aligned} \tilde{E}(u) = & \frac{1}{2} \int_{\mathbf{R}^N} |\nabla u|^2 dx - \int_{\mathbf{R}^N} \int_{\mathbf{R}^N} F_1(|u(x)|^2) \frac{1}{|x-y|^{N-2}} F_1(|u(y)|^2) dx dy \\ & + \int_{\mathbf{R}^N} H_1(|u(x)|^2) dx \end{aligned}$$

is the Hamiltonian for the generalized Hartree equation

$$iu_t + \Delta u + 4 \left(\int_{\mathbf{R}^N} \frac{F_1(|u(y)|^2)}{|x-y|^{N-2}} dy \right) F'_1(|u|^2)(x) u(x) - 2H'_1(|u(x)|^2) u(x) = 0, \quad (4.9)$$

and $\tilde{Q}(u) = \int_{\mathbf{R}^N} |u^2(x)| dx$ is a conserved quantity for this evolution equation. The critical points of $\tilde{E} + \omega \tilde{Q}$ give rise to standing waves for (4.9). As far as minimization is concerned, using an argument of T. Cazenave and P.-L. Lions (see the proof of Theorem II.1 in [9, p. 555]), we may restrict ourselves to the real functionals $E(u)$ and $Q(u)$.

In the case $N = 3$, $F(u) = G(u) = u^2$ and $H(u) = 0$, the problem of minimizing $E(u)$ subject to $Q(u) = \lambda$ has been studied in [15], where the existence, the radial symmetry and the uniqueness of the minimizer have been proved. The symmetry was proved by using a sharp inequality for spherical rearrangements. This can still be used in our case if we know that the minimizer is nonnegative and if we assume that F is increasing on $[0, \infty)$ (because the equality $F(u^*) = (F(u))^*$ is needed). Using the results in Sections 2 and 3, we will show the radial symmetry of minimizers in dimension $N \geq 3$ under more general assumptions on F , G and H .

We begin by studying some properties of the nonlocal term appearing in (4.8).

Lemma 4.5. *Let $N \geq 3$ and let $F : \mathbf{R} \rightarrow \mathbf{R}$ be a function of class C^2 satisfying $F(0) = F'(0) = 0$ and*

$$|F'(x)| \leq C|x|^\sigma \quad \text{for } |x| \geq 1,$$

where $C > 0$ is a constant and $\sigma < \frac{4}{N-2}$. Then the singular integral operator

$$I(\varphi)(x) = \int_{\mathbf{R}^N} \frac{1}{|x-y|^{N-2}} \varphi(y) dy$$

and the functional

$$M(\varphi) = \int_{\mathbf{R}^N} \int_{\mathbf{R}^N} F(\varphi(x)) \frac{1}{|x-y|^{N-2}} F(\varphi(y)) dx dy$$

have the following properties:

- (i) I is continuous from $L^p(\mathbf{R}^N)$ to $L^q(\mathbf{R}^N)$ if $1 < p < q < \infty$ and $\frac{1}{q} = \frac{1}{p} - \frac{2}{N}$.
- (ii) If $1 \leq p_1 < \frac{N}{2} < p_2 \leq \infty$, then I is continuous from $L^{p_1}(\mathbf{R}^N) \cap L^{p_2}(\mathbf{R}^N)$ to $L^\infty(\mathbf{R}^N) \cap C^0(\mathbf{R}^N)$.
- (iii) If $1 \leq r_1 < \frac{2N}{N+2} < r_2 \leq 2$ and $\varphi \in L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)$, then

$$\widehat{I(\varphi)}(\xi) = \frac{4\pi^{\frac{N}{2}}}{\Gamma(\frac{N}{2}-1)} \cdot \frac{1}{|\xi|^2} \widehat{\varphi}(\xi) \quad \text{in } \mathcal{S}'(\mathbf{R}^N).$$

- (iv) M is well defined and differentiable on $H^1(\mathbf{R}^N)$ and

$$M'(u) \cdot \varphi = 2 \int_{\mathbf{R}^N} \left(\int_{\mathbf{R}^N} \frac{F(u(y))}{|x-y|^{N-2}} dy \right) F'(u(x)) \varphi(x) dx.$$

- (v) For any $u \in H^1(\mathbf{R}^N)$ we have

$$M(u) = c_N \int_{\mathbf{R}^N} \frac{1}{|\xi|^2} |\widehat{F(u)}(\xi)|^2 d\xi, \quad \text{where } c_N = \frac{1}{2^{N-2} \pi^{\frac{N}{2}} \Gamma(\frac{N}{2}-1)}.$$

Proof. (i) follows directly from Theorem 1 in [23, pp. 119–120].

(ii) We write $\frac{1}{|x|^{N-2}}$ as $a_1(x) + a_2(x)$, where $a_1(x) = \frac{1}{|x|^{N-2}} \chi_{\{|x|>1\}}$ and $a_2(x) = \frac{1}{|x|^{N-2}} \chi_{\{|x|\leq 1\}}$. Then we have $I(\varphi) = a_1 * \varphi + a_2 * \varphi$. It is obvious that $a_1 \in L^q(\mathbf{R}^N)$ for $q \in (\frac{N}{N-2}, \infty]$ and $a_2 \in L^q(\mathbf{R}^N)$ for $q \in [1, \frac{N}{N-2})$. Let p'_1 and p'_2 be the conjugate exponents of p_1 and p_2 . Then $p'_1 > \frac{N}{N-2}$ and $p'_2 < \frac{N}{N-2}$, so that $a_1 \in L^{p'_1}(\mathbf{R}^N)$ and $a_2 \in L^{p'_2}(\mathbf{R}^N)$. We infer that $I(\varphi)$ is continuous and by Young's inequality we get

$$\|I(\varphi)\|_{L^\infty} \leq \|a_1\|_{L^{p'_1}} \cdot \|\varphi\|_{L^{p_1}} + \|a_2\|_{L^{p'_2}} \cdot \|\varphi\|_{L^{p_2}}.$$

- (iv) First we consider the bilinear form

$$P(\varphi, \psi) = \int_{\mathbf{R}^N} \int_{\mathbf{R}^N} \varphi(x) \frac{1}{|x-y|^{N-2}} \overline{\psi(y)} dx dy.$$

Notice that P is well defined and continuous on $L^{\frac{2N}{N+2}}(\mathbf{R}^N) \times L^{\frac{2N}{N+2}}(\mathbf{R}^N)$. Indeed, it follows from (i) that I is well defined and continuous from $L^{\frac{2N}{N+2}}(\mathbf{R}^N)$ to $L^{\frac{2N}{N-2}}(\mathbf{R}^N)$ and we have

$$|P(\varphi, \psi)| = \left| \int_{\mathbf{R}^N} I(\varphi)(x) \overline{\psi(x)} dx \right| \leq \|I(\varphi)\|_{L^{\frac{2N}{N-2}}} \cdot \|\psi\|_{L^{\frac{2N}{N+2}}} \leq A_N \|\varphi\|_{L^{\frac{2N}{N+2}}} \|\psi\|_{L^{\frac{2N}{N+2}}}.$$

Without loss of generality we may assume that $\sigma > \frac{2}{N}$. From the assumptions on F we have $|F(u)| \leq C|u|^2$ if $|u| \leq 1$ and $|F(u)| \leq C|u|^{1+\sigma}$ if $|u| > 1$. It is well known that $H^1(\mathbf{R}^N)$ is continuously embedded in $L^p(\mathbf{R}^N)$ for $p \in [2, \frac{2N}{N-2}]$ and then it is standard (see, e.g. [26, Appendix A]) that $u \mapsto F(u)$ is continuously differentiable from $H^1(\mathbf{R}^N)$ to $L^q(\mathbf{R}^N)$ for $q \in [\max(1, \frac{2}{1+\sigma}), \frac{2N}{(N-2)(1+\sigma)}]$. In particular, $u \mapsto F(u)$ is continuously differentiable from $H^1(\mathbf{R}^N)$ to $L^{\frac{2N}{N+2}}(\mathbf{R}^N)$ (because $\frac{2}{1+\sigma} < \frac{2N}{N+2} < \frac{2N}{(N-2)(1+\sigma)}$). Since $M(u) = P(F(u), F(u))$, (iv) follows.

(iii) and (v). Let $K(x) = \frac{1}{|x|^{N-2}}$. Then $K \in \mathcal{S}'(\mathbf{R}^N)$ and it follows from Theorem 4.1 in [24, p. 160] or from Lemma 1 in [23, p. 117] that $\widehat{K}(\xi) = \frac{4\pi^{\frac{N}{2}}}{\Gamma(\frac{N}{2}-1)} \cdot \frac{1}{|\xi|^2}$. From Lemma 1 in [23, p. 117] we have

$$P(\varphi, \psi) = \frac{1}{(2\pi)^N} \int_{\mathbf{R}^N} \widehat{I(\varphi)}(\xi) \overline{\widehat{\psi}(\xi)} d\xi = c_N \int_{\mathbf{R}^N} \frac{1}{|\xi|^2} \widehat{\varphi}(\xi) \overline{\widehat{\psi}(\xi)} d\xi \quad (4.10)$$

whenever $\varphi, \psi \in \mathcal{S}(\mathbf{R}^N)$. We claim that (4.10) holds for any $\varphi, \psi \in L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)$ with $1 \leq r_1 < \frac{2N}{N+2} < r_2 \leq 2$. This assertion implies both (iii) and (v).

Now let us prove the claim. Since (4.10) holds on $\mathcal{S} \times \mathcal{S}$, the bilinear form P is continuous on $L^{\frac{2N}{N+2}}(\mathbf{R}^N) \times L^{\frac{2N}{N+2}}(\mathbf{R}^N)$ and $L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)$ is continuously embedded into $L^{\frac{2N}{N+2}}(\mathbf{R}^N)$, all we have to do is to show that the bilinear form

$$P_1(\varphi, \psi) = \int_{\mathbf{R}^N} \frac{1}{|\xi|^2} \widehat{\varphi}(\xi) \overline{\widehat{\psi}(\xi)} d\xi$$

is continuous on $(L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)) \times (L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N))$; then the claim follows by density of $\mathcal{S}$ in $L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)$.

Let r'_1, r'_2 be the conjugate exponents of r_1, r_2 and let q_1, q_2 be such that $\frac{1}{r'_1} + \frac{1}{q_1} = \frac{1}{2}$, respectively $\frac{1}{r'_2} + \frac{1}{q_2} = \frac{1}{2}$. Let $b_1(\xi) = \frac{1}{|\xi|} \chi_{\{|\xi| \leq 1\}}$ and $b_2(\xi) = \frac{1}{|\xi|} \chi_{\{|\xi| > 1\}}$. We have $2 \leq q_1 < N$ and $q_2 > N$, so that $b_1 \in L^{q_1}(\mathbf{R}^N)$ and $b_2 \in L^{q_2}(\mathbf{R}^N)$. Since the Fourier transform maps continuously $L^{r_1}(\mathbf{R}^N)$ into $L^{r'_1}(\mathbf{R}^N)$ and $L^{r_2}(\mathbf{R}^N)$ into $L^{r'_2}(\mathbf{R}^N)$, we have:

$$\begin{aligned} |P_1(\varphi, \psi)| &\leq \left| \int_{\{|\xi| \leq 1\}} \frac{1}{|\xi|^2} \widehat{\varphi}(\xi) \overline{\widehat{\psi}(\xi)} d\xi \right| + \left| \int_{\{|\xi| > 1\}} \frac{1}{|\xi|^2} \widehat{\varphi}(\xi) \overline{\widehat{\psi}(\xi)} d\xi \right| \\ &\leq \|b_1 \widehat{\varphi}\|_{L^2} \cdot \|b_1 \widehat{\psi}\|_{L^2} + \|b_2 \widehat{\varphi}\|_{L^2} \cdot \|b_2 \widehat{\psi}\|_{L^2} \\ &\leq \|b_1\|_{L^{q_1}}^2 \|\widehat{\varphi}\|_{L^{r'_1}} \|\widehat{\psi}\|_{L^{r'_1}} + \|b_2\|_{L^{q_2}}^2 \|\widehat{\varphi}\|_{L^{r'_2}} \|\widehat{\psi}\|_{L^{r'_2}} \end{aligned}$$

$$\leq C(N, r_1, r_2)(\|\varphi\|_{L^{r_1}} \|\psi\|_{L^{r_1}} + \|\varphi\|_{L^{r_2}} \|\psi\|_{L^{r_2}}).$$

This proves the continuity of P_1 and our claim. Thus the proof of Lemma 4.5 is complete. $\square$

Theorem 4.6. Let $N \geq 3$ and let $F, G, H: \mathbf{R} \rightarrow \mathbf{R}$ be C^2 functions satisfying the following assumptions:

(a) $F(0) = F'(0) = 0$ and there exist $\sigma < \frac{4}{N-2}$ and $C > 0$ such that

$$|F'(u)| \leq C|u|^\sigma \quad \text{if } |u| \geq 1.$$

(b) There exist $\sigma_1 \in [1, \frac{N+2}{N-2})$ and $C_1 > 0$ such that

$$|G'(u)| \leq C_1|u|^{\sigma_1} \quad \text{and} \quad |H'(u)| \leq C_1|u|^{\sigma_1} \quad \text{for any } u \in \mathbf{R}.$$

Moreover, if $\sigma_1 < 2$ then we assume that $\sigma_1 \geq \max(\frac{(N-2)(1+2\sigma)-4}{N}, 1)$.

(c) For any $\varepsilon > 0$, $G' \neq 0$ on $(-\varepsilon, 0)$ and on $(0, \varepsilon)$.

Then any minimizer $u \in H^1(\mathbf{R}^N)$ of the functional E given by (4.8) subject to the constraint $Q(u) = \lambda \neq 0$ is radially symmetric (after a translation in $\mathbf{R}^N$).

Proof. First of all, notice that the functionals E and Q are well defined and of class C^1 on $H^1(\mathbf{R}^N)$. Let $u \in H^1(\mathbf{R}^N)$ be a minimizer. We will show that, except for translation, u is symmetric with respect to x_1 . The same proof is valid for any other direction in $\mathbf{R}^N$ and the radial symmetry of u follows as in the proof of Theorem 4.1.

After a translation in the x_1 direction we may suppose that

$$\int_{\{x_1 < 0\}} G(u(x)) dx = \int_{\{x_1 > 0\}} G(u(x)) dx = \frac{\lambda}{2}.$$

As before, we define $u_1 = T_1 u$ and $u_2 = T_2 u$. We know that $u_1, u_2 \in H^1(\mathbf{R}^N)$. In view of assumption (a), it is obvious that $F(u) \in L^1(\mathbf{R}^N)$ and we have $T_1(F(u)) = F(u_1)$, $T_2(F(u)) = F(u_2)$, $Q(u_1) = Q(u_2) = \lambda$. Defining $W(\varphi) = \int_{\mathbf{R}^N} \frac{1}{|\xi|^2} |\widehat{\varphi}(\xi)|^2 d\xi$, from Lemma 4.5(v) we get

$$\begin{aligned} E(u_1) + E(u_2) - 2E(u) &= -[M(u_1) + M(u_2) - 2M(u)] \\ &= -c_N[W(T_1(F(u))) + W(T_2(F(u))) - 2W(F(u))]. \end{aligned} \quad (4.11)$$

Recall that by (2.38) we have for any $\varphi \in C_c^\infty(\mathbf{R}^N)$,

$$W(T_1\varphi) + W(T_2\varphi) - 2W(\varphi) = \frac{8}{\pi} \int_{\mathbf{R}^{N-1}} \frac{1}{|\xi'|} \left| \int_0^\infty \widehat{A\varphi}(\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 \right|^2 d\xi'. \quad (4.12)$$

To show that this identity also holds for $F(u)$ we need the following lemma.

Lemma 4.7. Let $N \geq 3$ and let r_1, r_2 be such that $1 < r_1 < \frac{2N}{N+2} < r_2 < 2$. The bilinear form

$$R(\varphi, \psi) = \int_{\mathbf{R}^{N-1}} \frac{1}{|\xi'|} \int_0^\infty \widehat{\varphi}(\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 \cdot \int_0^\infty \widehat{\psi}(\eta_1, \xi') \frac{\eta_1}{|\xi'|^2 + \eta_1^2} d\eta_1 d\xi'$$

is continuous on $(L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)) \times (L^{r'_1}(\mathbf{R}^N) \cap L^{r'_2}(\mathbf{R}^N))$.

Proof. Consider $\varphi, \psi \in L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)$. Then $\widehat{\varphi}, \widehat{\psi} \in L^{r'_1}(\mathbf{R}^N) \cap L^{r'_2}(\mathbf{R}^N)$, where r'_1 and r'_2 are the conjugate exponents of r_1 and r_2 . Using Hölder's inequality and the change of variable $\xi_1 = t|\xi'|$, we get for $\xi' \neq 0$ and $i = 1, 2$,

$$\begin{aligned} \left| \int_0^\infty \widehat{\varphi}(\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 \right| &\leq \left(\int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^{r'_i} d\xi_1 \right)^{\frac{1}{r'_i}} \left(\int_0^\infty \frac{\xi_1^{r_i}}{(|\xi'|^2 + \xi_1^2)^{r_i}} d\xi_1 \right)^{\frac{1}{r_i}} \\ &= |\xi'|^{\frac{1-r_i}{r_i}} \left(\int_0^\infty \frac{t^{r_i}}{(1+t^2)^{r_i}} dt \right)^{\frac{1}{r_i}} \left(\int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^{r'_i} d\xi_1 \right)^{\frac{1}{r'_i}} \\ &= C_i |\xi'|^{\frac{1-r_i}{r_i}} \left(\int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^{r'_i} d\xi_1 \right)^{\frac{1}{r'_i}}. \end{aligned} \quad (4.13)$$

A similar estimate holds for ψ . Let q_i be the conjugate exponent of $\frac{r'_i}{2}$, i.e. $q_i = \frac{r_i}{2-r_i}$. Using (4.13), Hölder's inequality and the estimate $\|\widehat{\varphi}\|_{L^{r'_i}} \leq A_i \|\varphi\|_{L^{r_i}}$ we have

$$\begin{aligned} &\left| \int_{B_{\mathbf{R}^{N-1}}(0,1)} \frac{1}{|\xi'|} \int_0^\infty \widehat{\varphi}(\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 \cdot \int_0^\infty \widehat{\psi}(\eta_1, \xi') \frac{\eta_1}{|\xi'|^2 + \eta_1^2} d\eta_1 d\xi' \right| \\ &\leq C_1^2 \int_{B_{\mathbf{R}^{N-1}}(0,1)} |\xi'|^{\frac{2-2r_1}{r_1}-1} \left(\int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^{r'_1} d\xi_1 \right)^{\frac{1}{r'_1}} \left(\int_0^\infty |\widehat{\psi}(\eta_1, \xi')|^{r'_1} d\eta_1 \right)^{\frac{1}{r'_1}} d\xi' \\ &\leq C_1^2 \left(\int_{B_{\mathbf{R}^{N-1}}(0,1)} |\xi'|^{\frac{q_1(2-3r_1)}{r_1}} d\xi' \right)^{\frac{1}{q_1}} \left(\int_{B_{\mathbf{R}^{N-1}}(0,1)} \int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^{r'_1} d\xi_1 d\xi' \right)^{\frac{1}{r'_1}} \\ &\quad \times \left(\int_{B_{\mathbf{R}^{N-1}}(0,1)} \int_0^\infty |\widehat{\psi}(\eta_1, \xi')|^{r'_1} d\eta_1 d\xi' \right)^{\frac{1}{r'_1}} \\ &\leq C_1^2 A_1^2 \left(\int_{B_{\mathbf{R}^{N-1}}(0,1)} |\xi'|^{\frac{q_1(2-3r_1)}{r_1}} d\xi' \right)^{\frac{1}{q_1}} \|\varphi\|_{L^{r_1}} \|\psi\|_{L^{r_1}} \end{aligned} \quad (4.14)$$

and

$$\begin{aligned}
 & \left| \int_{\{|\xi'|>1\}} \frac{1}{|\xi'|} \int_0^\infty \widehat{\varphi}(\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 \cdot \int_0^\infty \widehat{\psi}(\eta_1, \xi') \frac{\eta_1}{|\xi'|^2 + \eta_1^2} d\eta_1 d\xi' \right| \\
 & \leq C_2^2 \int_{\{|\xi'|>1\}} |\xi'|^{\frac{2-2r_2}{r_2}-1} \left(\int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^{r'_2} d\xi_1 \right)^{\frac{1}{r'_2}} \left(\int_0^\infty |\widehat{\psi}(\eta_1, \xi')|^{r'_2} d\eta_1 \right)^{\frac{1}{r'_2}} d\xi' \\
 & \leq C_2^2 \left(\int_{\{|\xi'|>1\}} |\xi'|^{\frac{q_1(2-3r_2)}{r_2}} d\xi' \right)^{\frac{1}{q_2}} \left(\int_{\{|\xi'|>1\}} \int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^{r'_2} d\xi_1 d\xi' \right)^{\frac{1}{r'_2}} \\
 & \quad \times \left(\int_{\{|\xi'|>1\}} \int_0^\infty |\widehat{\psi}(\eta_1, \xi')|^{r'_2} d\eta_1 d\xi' \right)^{\frac{1}{r'_2}} \\
 & \leq C_2^2 A_2^2 \left(\int_{\{|\xi'|>1\}} |\xi'|^{\frac{q_2(2-3r_2)}{r_2}} d\xi' \right)^{\frac{1}{q_2}} \|\varphi\|_{L^{r_2}} \|\psi\|_{L^{r_2}}. \tag{4.15}
 \end{aligned}$$

Since $1 < r_1 < \frac{2N}{N+2} < r_2 < 2$, a direct computation shows that $\int_{B_{\mathbf{R}^{N-1}}(0,1)} |\xi'|^{\frac{q_1(2-3r_1)}{r_1}} d\xi'$ and $\int_{\{|\xi'|>1\}} |\xi'|^{\frac{q_2(2-3r_2)}{r_2}} d\xi'$ are finite. From (4.14) and (4.15) we have

$$|R(\varphi, \psi)| \leq K (\|\varphi\|_{L^{r_1}} \|\psi\|_{L^{r_1}} + \|\varphi\|_{L^{r_2}} \|\psi\|_{L^{r_2}})$$

and Lemma 4.7 is proved. $\square$

Let r_1 and r_2 be as in Lemma 4.7. Since the maps $\varphi \mapsto T_1\varphi$ and $\varphi \mapsto T_2\varphi$ are obviously continuous from $L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)$ into itself and we have shown in the proof of Lemma 4.5 that the bilinear form $P_1(\varphi, \psi) = \int_{\mathbf{R}^N} \frac{1}{|\xi|^2} \widehat{\varphi}(\xi) \overline{\widehat{\psi}(\xi)} d\xi$ is continuous on this space, it follows that the left-hand side of (4.12) is continuous on $L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)$. By Lemma 4.7, the right-hand side of (4.12) also defines a continuous functional on $L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)$. Since (4.12) is valid for any $\varphi \in C_c^\infty(\mathbf{R}^N)$, by density we infer that (4.12) holds for any $\varphi \in L^{r_1}(\mathbf{R}^N) \cap L^{r_2}(\mathbf{R}^N)$. Recall that $u \in H^1(\mathbf{R}^N)$ and by the Sobolev embedding and assumption (a) we have $F(u) \in L^q(\mathbf{R}^N)$ for any $q \in [\max(1, \frac{2}{1+\sigma}), \frac{2N}{(N-2)(1+\sigma)}]$; hence (4.12) is valid for $F(u)$.

Since u is a minimizer, we must have $E(u_1) + E(u_2) - 2E(u) \geq 0$. From (4.11) and (4.12) we infer that necessarily

$$\int_{\mathbf{R}^{N-1}} \frac{1}{|\xi'|} \left| \int_0^\infty \mathcal{F}(A(F(u)))(\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 \right|^2 d\xi' = 0. \tag{4.16}$$

Contrary to our previous examples, (4.16) *does not* imply directly $AF(u) \equiv 0$. To see this, consider a function $\psi \in C_c^\infty(0, \infty)$ such that $\text{supp}(\psi) \subset [1, \infty)$, $\psi \not\equiv 0$ and $\int_0^\infty \frac{t}{1+t^2} \psi(t) dt = 0$.

(Such a function exists: for example, take two nonnegative functions $\psi_0, \psi_1 \in C_c^\infty(1, \infty)$ with disjoint supports and put $\psi_\tau = (1 - \tau)\psi_0 - \tau\psi_1$. There is some $\tau \in (0, 1)$ such that $\int_0^\infty \frac{t}{1+t^2} \psi_\tau(t) dt = 0$.) Extend ψ to an odd function defined on $\mathbf{R}$. Take $\alpha \in C_c^\infty(\mathbf{R}^{N-1})$ such that $\alpha \not\equiv 0$ and $\text{supp}(\alpha) \subset \mathbf{R}^{N-1} \setminus B(0, 1)$ and put $\widehat{f}(\xi_1, \xi') = \alpha(\xi')\psi(\frac{\xi_1}{|\xi'|})$. Then $\widehat{f} \in C_c^\infty(\mathbf{R}^N)$ (hence $f \in \mathcal{S}$), $f \not\equiv 0$ and f is odd with respect to the first variable. However, we have $\int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 = 0$ for any $\xi' \neq 0$ and consequently

$$\int_{\mathbf{R}^{N-1}} \frac{1}{|\xi'|} \left| \int_0^\infty \widehat{f}(\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 \right|^2 d\xi' = 0.$$

To show that u is symmetric with respect to x_1 , we argue as follows: since u and u_1 minimize E under the constraint $Q = \lambda$, these functions satisfy the Euler–Lagrange equations $E'(u) + \alpha Q'(u) = 0$, respectively $E'(u_1) + \beta Q'(u_1) = 0$ for some constants α and β , that is

$$-\Delta u - 2I(F(u))F'(u) + H'(u) + \alpha G'(u) = 0 \quad \text{in } \mathbf{R}^N, \quad (4.17)$$

$$-\Delta u_1 - 2I(F(u_1))F'(u_1) + H'(u_1) + \beta G'(u_1) = 0 \quad \text{in } \mathbf{R}^N. \quad (4.18)$$

We will show in the next lemma that u and u_1 are smooth functions. Then we prove that $I(F(u))(x) = I(F(u_1))(x)$ in the half-space $\{x_1 < 0\}$. Together with assumption (c), this implies that $\alpha = \beta$ in (4.17)–(4.18). Then we will be able to apply the Unique Continuation Principle to prove that $u = u_1$.

Lemma 4.8. *Let $u \in H^1(\mathbf{R}^N)$ be a solution of (4.17), where $F, G, H \in C^2(\mathbf{R})$ satisfy the assumptions (a) and (b) in Theorem 4.6. Then $u \in W^{3,p}(\mathbf{R}^N)$ for any $p \in [2, \infty)$. In particular, $u \in C^2(\mathbf{R}^N)$ and $D^\alpha u$ are continuous and bounded on $\mathbf{R}^N$ if $\alpha \in \mathbf{N}^N$, $|\alpha| \leq 2$.*

Proof. The proof relies on a classical boot-strap argument. We show first that $u \in L^\infty(\mathbf{R}^N)$. By the Sobolev embedding we have $u \in L^q(\mathbf{R}^N)$ for $q \in [2, \frac{2N}{N-2}]$. We will improve this estimate by an inductive argument to get the desired conclusion.

We consider only the case $N \geq 4$, the proof in the case $N = 3$ being similar. Assume that $u \in L^q(\mathbf{R}^N)$ for any $q \in [2, \beta]$, where $\beta \geq \frac{2N}{N-2}$. It is clear that $G'(u), H'(u) \in L^q(\mathbf{R}^N)$ for $q \in [\max(1, \frac{2}{\sigma_1}), \frac{\beta}{\sigma_1}]$ and $F(u) \in L^q(\mathbf{R}^N)$ for $q \in [1, \frac{\beta}{1+\sigma}]$. We distinguish two cases:

Case A. If $\frac{\beta}{1+\sigma} > \frac{N}{2}$, then $I(F(u)) \in L^q(\mathbf{R}^N)$ for any $q \in (\frac{N}{N-2}, \infty]$. We have $F'(u)\chi_{\{|u| \leq 1\}} \in L^q(\mathbf{R}^N)$ for $q \in [2, \infty]$, hence $I(F(u))F'(u)\chi_{\{|u| \leq 1\}} \in L^q(\mathbf{R}^N)$ for $q \in (1, \infty]$ if $N = 4$, respectively for $q \in [1, \infty]$ if $N \geq 5$ and $F'(u)\chi_{\{|u| > 1\}} \in L^q(\mathbf{R}^N)$ for $q \in [1, \frac{\beta}{\sigma}]$, hence $I(F(u))F'(u)\chi_{\{|u| > 1\}} \in L^q(\mathbf{R}^N)$ if $q \in [1, \frac{\beta}{\sigma}]$. Consequently $I(F(u))F'(u) \in L^q(\mathbf{R}^N)$ for $q \in (1, \frac{\beta}{\sigma})$ if $N = 4$, respectively for $q \in [1, \frac{\beta}{\sigma}]$ if $N \geq 5$. Notice that $\beta \geq \frac{2N}{N-2}$ and the second part of assumption (b) imply $\frac{\beta}{\sigma} \geq \frac{2}{\sigma_1}$. Using Eq. (4.17) we infer that $\Delta u \in L^q(\mathbf{R}^N)$ for $q \in [\max(1, \frac{2}{\sigma_1}), \min(\frac{\beta}{\sigma_1}, \frac{\beta}{\sigma})]$, $q \neq 1$ if $N = 4$. Let $q_3 = \min(\frac{\beta}{\sigma_1}, \frac{\beta}{\sigma})$. Notice that $q_3 \leq \beta$ because $\sigma_1 \geq 1$ and $\Delta u \in L^{q_3}(\mathbf{R}^N)$. If $q_3 > \frac{N}{2} \geq 2$, then $u \in L^{q_3}(\mathbf{R}^N)$, hence $u \in W^{2,q_3}(\mathbf{R}^N)$

and by the Sobolev embedding we get $u \in L^\infty(\mathbf{R}^N)$. If $q_3 = \frac{N}{2}$, then $u \in W^{2, \frac{N}{2}}(\mathbf{R}^N)$, consequently $u \in L^q(\mathbf{R}^N)$ for any $q \in [2, \infty)$ and repeating the above proof with $\tilde{\beta} > \beta$ we find $u \in L^\infty(\mathbf{R}^N)$. If $q_3 < \frac{N}{2}$, then necessarily $q_3 = \frac{\beta}{\sigma_1}$ (recall that $\frac{\beta}{\sigma} > \frac{\beta}{1+\sigma} > \frac{N}{2}$ because we are in Case A). By the Sobolev embedding we get $u \in L^{\beta_1}(\mathbf{R}^N)$, where $\frac{1}{\beta_1} = \frac{1}{q_3} - \frac{2}{N} = \frac{\sigma_1}{\beta} - \frac{2}{N}$, thus $\frac{1}{\beta_1} - \frac{1}{\beta} = \frac{\sigma_1-1}{\beta} - \frac{2}{N} \leq \frac{(\sigma_1-1)(N-2)-4}{2N} < 0$ by (b). Repeating the previous arguments with β replaced by β_1 , we find that either $u \in L^\infty(\mathbf{R}^N)$ or $u \in L^{\beta_2}(\mathbf{R}^N)$, where $\beta_2 > \beta_1$ and $\frac{1}{\beta_2} - \frac{1}{\beta_1} \leq \frac{(\sigma_1-1)(N-2)-4}{2N}$, and so on. After a finite number of steps we get $u \in L^\infty(\mathbf{R}^N)$.

Case B. If $\frac{\beta}{1+\sigma} \leq \frac{N}{2}$, we may suppose that $\frac{\beta}{1+\sigma} < \frac{N}{2}$. By Lemma 4.5(i), $I(F(u)) \in L^q(\mathbf{R}^N)$ for $q \in (\frac{N}{N-2}, (\frac{1+\sigma}{\beta} - \frac{2}{N})^{-1}]$. As in Case A we get $I(F(u))F'(u) \in L^q(\mathbf{R}^N)$ for $q \in [1, (\frac{1+2\sigma}{\beta} - \frac{2}{N})^{-1}]$, $q \neq 1$ if $N = 4$. By (a), (b) and the fact that $\beta \geq \frac{2N}{N-2}$ we have $(\frac{1+2\sigma}{\beta} - \frac{2}{N})^{-1} \geq \frac{2}{\sigma_1}$. Since $G'(u), H'(u) \in L^q(\mathbf{R}^N)$ for $q \in [\max(1, \frac{2}{\sigma_1}), \frac{\beta}{\sigma_1}]$, using (4.17) we get $\Delta u \in L^q(\mathbf{R}^N)$ for $q \in [\max(1, \frac{2}{\sigma_1}), q_4]$, $q \neq 1$ if $N = 4$, where $q_4 = \min(\frac{\beta}{\sigma_1}, (\frac{1+2\sigma}{\beta} - \frac{2}{N})^{-1})$. If $q_4 \geq \frac{N}{2}$ then, as above, we obtain $u \in L^\infty(\mathbf{R}^N)$. Otherwise by the Sobolev embedding we find $u \in L^{\beta_1}(\mathbf{R}^N)$, where $\frac{1}{\beta_1} = \frac{1}{q_4} - \frac{2}{N}$, thus $\frac{1}{\beta_1} - \frac{1}{\beta} \leq \max(\frac{(\sigma_1-1)(N-2)-4}{2N}, \frac{\sigma(N-2)-4}{N}) < 0$. Then we restart the process with β_1 instead of β . Continuing in this way, after a finite number of steps we obtain $u \in L^\infty(\mathbf{R}^N)$.

We have proved that $u \in L^q(\mathbf{R}^N)$ for any $q \in [2, \infty]$. Thus $F(u) \in L^1(\mathbf{R}^N) \cap L^\infty(\mathbf{R}^N)$, $I(F(u)) \in L^q(\mathbf{R}^N)$ for $q \in (\frac{N}{N-2}, \infty]$, $F'(u) \in L^2(\mathbf{R}^N) \cap L^\infty(\mathbf{R}^N)$, hence $I(F(u))F'(u) \in L^2(\mathbf{R}^N) \cap L^\infty(\mathbf{R}^N)$. Clearly $G'(u), H'(u) \in L^q(\mathbf{R}^N)$ for $q \in [\max(1, \frac{2}{\sigma_1}), \infty]$. Using (4.17) we have $\Delta u \in L^2(\mathbf{R}^N) \cap L^\infty(\mathbf{R}^N)$, thus $u \in W^{2,p}(\mathbf{R}^N)$ for any $p \in [2, \infty)$. In particular, $\frac{\partial u}{\partial x_i}$ are continuous and bounded on $\mathbf{R}^N$. Differentiating (4.17) with respect to x_i we get

$$\begin{aligned} & -\Delta \left(\frac{\partial u}{\partial x_i} \right) - 2I \left(F'(u) \frac{\partial u}{\partial x_i} \right) F'(u) - 2I(F(u))F''(u) \frac{\partial u}{\partial x_i} \\ & + G''(u) \frac{\partial u}{\partial x_i} + \alpha H''(u) \frac{\partial u}{\partial x_i} = 0 \quad \text{in } \mathbf{R}^N. \end{aligned}$$

It follows that $-\Delta(\frac{\partial u}{\partial x_i}) \in L^2(\mathbf{R}^N) \cap L^\infty(\mathbf{R}^N)$. Since obviously $\frac{\partial u}{\partial x_i} \in L^2(\mathbf{R}^N) \cap L^\infty(\mathbf{R}^N)$, we get $\frac{\partial u}{\partial x_i} \in W^{2,p}(\mathbf{R}^N)$, which implies $u \in W^{3,p}(\mathbf{R}^N)$ for any $p \in [2, \infty)$. $\square$

It follows from Lemma 4.8 that $F(u) \in C^2(\mathbf{R}^N)$ and $F(u) \in W^{2,p}(\mathbf{R}^N)$ for $p \in [1, \infty]$. Using Lemma 4.5(i) and (ii), it is easy to check that $I(F(u)) \in C^2(\mathbf{R}^N)$ and $I(F(u)) \in W^{2,p}(\mathbf{R}^N)$ for $p \in (\frac{N}{N-2}, \infty]$. In particular, $I(F(u)) \in \mathcal{S}'(\mathbf{R}^N)$ and Lemma 4.5(iii) implies $\mathcal{F}(I(F(u))) (\xi) = d_N \frac{1}{|\xi|^2} \widehat{F(u)}(\xi)$, where $d_N = \frac{4\pi^{\frac{N}{2}}}{\Gamma(\frac{N}{2}-1)}$. Setting $U = I(F(u))$ we have $-\Delta U = d_N F(u)$.

Next we show that $\frac{\partial U}{\partial x_1}(0, x') = \frac{\partial}{\partial x_1} I(F(u))(0, x') = 0$ for any $x' \in \mathbf{R}^{N-1}$. From (4.16) we infer that $\int_0^\infty \mathcal{F}(A(F(u))) (\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 = 0$ for almost every $\xi' \in \mathbf{R}^{N-1}$, that is $\int_{-\infty}^\infty \widehat{F(u)}(\xi_1, \xi') \frac{\xi_1}{|\xi'|^2 + \xi_1^2} d\xi_1 = 0$ a.e. $\xi' \in \mathbf{R}^{N-1}$, or equivalently

$$\int_{-\infty}^{\infty} \xi_1 \mathcal{F}(I(F(u)))(\xi_1, \xi') d\xi_1 = 0 \quad \text{for almost every } \xi' \in \mathbf{R}^{N-1}. \quad (4.19)$$

If $\frac{\partial}{\partial x_1} I(F(u))$ and $\mathcal{F}(\frac{\partial}{\partial x_1} I(F(u)))$ are in $L^1(\mathbf{R}^N)$, by the Fourier inversion theorem (4.19) is equivalent to $\frac{\partial}{\partial x_1} I(F(u))(0, x') = 0$, as desired.

Since we do not know whether $\frac{\partial}{\partial x_1} I(F(u)) \in L^1(\mathbf{R}^N)$ and $\mathcal{F}(\frac{\partial}{\partial x_1} I(F(u))) \in L^1(\mathbf{R}^N)$, we argue as follows: we take an arbitrary test function $\psi \in \mathcal{S}(\mathbf{R}^{N-1})$ and we put $\varphi_n(x_1) = \frac{n}{\sqrt{2\pi}} e^{-\frac{n^2 x_1^2}{2}}$.

Clearly, $\varphi_n(x_1) = n\varphi_1(nx_1)$, $\|\varphi_n\|_{L^1(\mathbf{R})} = 1$ and $\widehat{\varphi_n}(\xi_1) = e^{-\frac{\xi_1^2}{2n^2}}$. On one hand, we have, by using Lebesgue's Dominated Convergence Theorem,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_{\mathbf{R}^N} \varphi_n(x_1) \psi(x') \left[\frac{\partial}{\partial x_1} I(F(u)) \right] (x_1, x') dx \\ &= \lim_{n \rightarrow \infty} \int_{\mathbf{R}^N} \varphi_1(y_1) \psi(x') \left[\frac{\partial}{\partial x_1} I(F(u)) \right] \left(\frac{y_1}{n}, x' \right) dy_1 dx' \\ &= \int_{\mathbf{R}^{N-1}} \psi(x') \frac{\partial}{\partial x_1} (I(F(u)))(0, x') dx'. \end{aligned} \quad (4.20)$$

On the other hand, we have

$$\begin{aligned} & \int_{\mathbf{R}^N} \varphi_n(x_1) \psi(x') \left[\frac{\partial}{\partial x_1} I(F(u)) \right] (x_1, x') dx \\ &= \left\langle \frac{\partial}{\partial x_1} (I(F(u))), \varphi_n(x_1) \psi(x') \right\rangle_{S', S} \\ &= \left\langle \mathcal{F} \left(\frac{\partial}{\partial x_1} I(F(u)) \right), \mathcal{F}^{-1}(\varphi_n(x_1) \psi(x')) \right\rangle_{S', S} \\ &= \frac{1}{(2\pi)^N} \int_{\mathbf{R}^N} \frac{i d_N \xi_1}{|\xi|^2} \widehat{F(u)}(\xi) e^{-\frac{\xi_1^2}{2n^2}} \widehat{\psi}(-\xi') d\xi_1 d\xi'. \end{aligned} \quad (4.21)$$

Since $F(u) \in L^2(\mathbf{R}^N)$, for almost every $\xi' \in \mathbf{R}^{N-1}$ we have $\widehat{F(u)}(\cdot, \xi') \in L^2(\mathbf{R})$. For any such ξ' , arguing as in (4.13) we get

$$\int_{\mathbf{R}} \left| e^{-\frac{\xi_1^2}{2n^2}} \cdot \frac{\xi_1}{|\xi|^2} \widehat{F(u)}(\xi_1, \xi') \right| d\xi_1 \leq \int_{\mathbf{R}} \left| \frac{\xi_1}{\xi_1^2 + |\xi'|^2} \widehat{F(u)}(\xi_1, \xi') \right| d\xi_1 \leq \frac{C}{|\xi'|^{\frac{1}{2}}} \|\widehat{F(u)}(\cdot, \xi')\|_{L^2(\mathbf{R})},$$

where C does not depend on ξ' . Moreover, the Cauchy–Schwarz inequality gives

$$\int_{\mathbf{R}^{N-1}} \frac{C|\widehat{\psi}(-\xi')|}{|\xi'|^{\frac{1}{2}}} \|\widehat{F(u)}(\cdot, \xi')\|_{L^2(\mathbf{R})} d\xi' \leq C \left(\int_{\mathbf{R}^{N-1}} \frac{|\widehat{\psi}(-\xi')|^2}{|\xi'|} d\xi' \right)^{\frac{1}{2}} \|\widehat{F(u)}\|_{L^2(\mathbf{R}^N)} < \infty.$$

By the Dominated Convergence Theorem, we have for almost any $\xi' \in \mathbf{R}^{N-1}$

$$\int_{\mathbf{R}} \frac{\xi_1}{\xi_1^2 + |\xi'|^2} \widehat{F(u)}(\xi_1, \xi') e^{-\frac{\xi_1^2}{2n^2}} d\xi_1 \rightarrow \int_{\mathbf{R}} \frac{\xi_1}{\xi_1^2 + |\xi'|^2} \widehat{F(u)}(\xi_1, \xi') d\xi_1 = 0 \quad \text{as } n \rightarrow \infty.$$

Thus we may use Fubini's theorem, then the Dominated Convergence Theorem on $\mathbf{R}^{N-1}$ to obtain

$$\begin{aligned} & \int_{\mathbf{R}^N} \frac{\xi_1}{|\xi|^2} \widehat{F(u)}(\xi_1, \xi') e^{-\frac{\xi_1^2}{2n^2}} \psi(-\xi') d\xi_1 d\xi' \\ &= \int_{\mathbf{R}^{N-1}} \psi(-\xi') \int_{\mathbf{R}} \frac{\xi_1}{\xi_1^2 + |\xi'|^2} \widehat{F(u)}(\xi_1, \xi') e^{-\frac{\xi_1^2}{2n^2}} d\xi_1 d\xi' \\ &\rightarrow \int_{\mathbf{R}^{N-1}} \psi(-\xi') \cdot 0 d\xi' = 0 \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (4.22)$$

From (4.20)–(4.22) we infer that $\int_{\mathbf{R}^{N-1}} \psi(x') \frac{\partial}{\partial x_1} (I(F(u)))(0, x') dx' = 0$. Since $\psi \in \mathcal{S}(\mathbf{R}^{N-1})$ was arbitrary, we have $\frac{\partial}{\partial x_1} (I(F(u)))(0, \cdot) = 0$ in $\mathcal{S}'(\mathbf{R}^{N-1})$, hence $\frac{\partial}{\partial x_1} (I(F(u)))(0, x') = 0$ for any $x' \in \mathbf{R}^{N-1}$ because $\frac{\partial}{\partial x_1} (I(F(u)))$ is a continuous function.

We know that $F(u_1)$ is symmetric with respect to x_1 and a simple change of variables shows that the function $U_1 := I(F(u_1))$ is also symmetric with respect to x_1 . Clearly U_1 also belongs to $C^2(\mathbf{R}^N)$ and satisfies $-\Delta U_1 = -\Delta(I(F(u_1))) = d_N F(u_1)$. By symmetry we have $\frac{\partial U_1}{\partial x_1}(0, x') = 0$ for any $x' \in \mathbf{R}^{N-1}$. Since $u_1(x_1, x') = u(x_1, x')$ if $x_1 < 0$, we have proved that the functions U and U_1 are both solutions of the problem

$$\begin{cases} -\Delta W = d_N F(u) & \text{in } \{(x_1, x') \in \mathbf{R}^N \mid x_1 < 0\}, \\ W \in C^2(\mathbf{R}^N) \cap W^{2,p}(\mathbf{R}^N) & \text{for } p > \frac{N}{N-2}, \\ \frac{\partial W}{\partial x_1}(0, x') = 0 & \text{for any } x' \in \mathbf{R}^{N-1}. \end{cases} \quad (4.23)$$

It is not hard to see that the solution of (4.23) is unique. Consequently, $U(x_1, x') = U_1(x_1, x')$ if $x_1 < 0$. From (4.17) and (4.18) it is obvious that (u, U) and (u_1, U_1) solve the systems

$$\begin{cases} -\Delta u - 2UF'(u) + H'(u) + \alpha G'(u) = 0, \\ -\Delta U - d_N F(u) = 0 \end{cases} \quad \text{in } \mathbf{R}^N, \quad (4.24)$$

respectively

$$\begin{cases} -\Delta u_1 - 2U_1 F'(u_1) + H'(u_1) + \beta G'(u_1) = 0, \\ -\Delta U_1 - d_N F(u_1) = 0 \end{cases} \quad \text{in } \mathbf{R}^N. \quad (4.25)$$

We cannot have $u \equiv 0$ in the half-space $\{x_1 < 0\}$ because this would imply $\lambda = Q(u) = Q(u_1) = 0$. Since u is continuous, necessarily $u((-\infty, 0) \times \mathbf{R}^{N-1}) = u_1((-\infty, 0) \times \mathbf{R}^{N-1})$ contains an interval of the form $(-\varepsilon, 0)$ or $(0, \varepsilon)$ for some $\varepsilon > 0$. Now assumption (c), (4.24), (4.25) and the fact that $(u, U) = (u_1, U_1)$ on $(-\infty, 0) \times \mathbf{R}^{N-1}$ imply that $\alpha = \beta$ in (4.24)–(4.25). As a consequence, we see that $(u - u_1, U - U_1)$ solves a linear system whose coefficients belong to $L^\infty(\mathbf{R}^N)$. Since $(u, U) = (u_1, U_1)$ for $x_1 < 0$ and $(u, U), (u_1, U_1) \in W^{2,p}(\mathbf{R}^N, \mathbf{R}^2)$ if $p \geq 2$ and $p > \frac{N}{N-2}$, by using the Unique Continuation Principle we infer that $u = u_1$ (and $U = U_1$) in $\mathbf{R}^N$, that is u is symmetric with respect to x_1 .

Similarly we show that u is symmetric with respect to any other hyperplane Π which has the property that $\int_{\Pi_-} G(u(x)) dx = \int_{\Pi_+} G(u(x)) dx$, where Π_- and Π_+ are the two half-spaces determined by Π . As in the proof of Theorem 4.1 it follows that after a translation, u is radially symmetric. The proof of Theorem 4.6 is complete. $\square$

4.3. Standing waves for the Davey–Stewartson equation

We consider the Davey–Stewartson system

$$\begin{cases} iu_t + \Delta u = f(|u|^2)u - uv_{x_1}, \\ \Delta v = (|u|^2)_{x_1} \end{cases} \quad \text{in } \mathbf{R}^3, \quad (4.26)$$

which can be written as

$$iu_t = -\Delta u + f(|u|^2)u + R_1^2(|u|^2)u, \quad (4.27)$$

where R_1 is the Riesz transform defined by $\widehat{R_1 \varphi} = \frac{i\xi_1}{|\xi|} \widehat{\varphi}(\xi)$. Let $F_1(t) = \int_0^t f(\tau) d\tau$. It is easy to check that

$$\widetilde{E}(u) = \frac{1}{2} \int_{\mathbf{R}^3} |\nabla u|^2 dx + \frac{1}{2} \int_{\mathbf{R}^3} F_1(|u|^2) dx - \frac{1}{4} \int_{\mathbf{R}^3} |R_1(|u|^2)|^2 dx$$

is a Hamiltonian for (4.27) and $\widetilde{Q}(u) = \int_{\mathbf{R}^3} |u(x)|^2 dx$ is a conserved quantity for the same equation. The standing waves for (4.27) are precisely the critical points of $\widetilde{E} + \omega \widetilde{Q}$. As in the previous example, when we minimize $\widetilde{E}(u)$ subject to $\widetilde{Q}(u) = \text{constant}$, we may restrict ourselves to real functions u and to the real version of $\widetilde{E}$,

$$E(u) = \frac{1}{2} \int_{\mathbf{R}^3} |\nabla u|^2 dx + \int_{\mathbf{R}^3} F(u) dx - \frac{1}{4} \int_{\mathbf{R}^3} |R_1(u^2)|^2 dx.$$

We will consider a more general functional than $\widetilde{Q}$, namely $Q(u) = \int_{\mathbf{R}^3} G(u) dx$. If $G(u) = u^2$, in order to guarantee the boundedness from below of the functional E on the set

of functions satisfying $Q(u) = \lambda$, the function $F(u)$ is required to behave as $a|u|^\gamma$ for u large, with $a > 0$ and $\gamma > 4$. In the case $F(u) = a|u|^\gamma$, the Cauchy problem for the evolution equation (4.27) has been analysed in [12]. The global existence of solutions was proved in the case $a > 0$ and $\gamma > 4$, while in the case $\gamma = 4$ the global existence was proved if a is sufficiently large.

Still in the case of pure power $F(u) = a|u|^\gamma$, with $a > 0$ and $\gamma > 4$, the existence of minimizers of E subject to the constraint $Q(u) = \int_{\mathbf{R}^3} |u|^2 dx = \lambda$ can be proved by using the Concentration–Compactness Principle (see [17]) if λ is large enough (this assumption is needed to prevent vanishing).

In [10] the existence of ground states related to the problem (4.26) has been studied. However, our method cannot be used to prove the symmetry of these ground states because the nonlocal term appears in the constraint.

It is well known that R_1 is a linear continuous map from $L^p(\mathbf{R}^3)$ to $L^p(\mathbf{R}^3)$ for $1 < p < \infty$ (see [23]). If $u^2 \in L^2(\mathbf{R}^3)$, then $R_1(u^2) \in L^2(\mathbf{R}^3)$ and by Plancherel's theorem we get

$$\int_{\mathbf{R}^3} |R_1(u^2)|^2 dx = \frac{1}{(2\pi)^3} \int_{\mathbf{R}^3} |\widehat{R_1(u^2)}(\xi)|^2 d\xi = \frac{1}{(2\pi)^3} \int_{\mathbf{R}^3} \frac{\xi_1^2}{|\xi|^2} |\widehat{u^2}(\xi)|^2 d\xi. \quad (4.28)$$

We have the following symmetry result.

Theorem 4.9. *Let $u \in H^1(\mathbf{R}^3)$ be a solution of the minimization problem*

$$\begin{aligned} \text{minimize} \quad E(u) &= \frac{1}{2} \int_{\mathbf{R}^3} |\nabla u|^2 dx + \int_{\mathbf{R}^3} F(u) dx - \frac{1}{4} \int_{\mathbf{R}^3} |R_1(u^2)|^2 dx \\ \text{subject to} \quad Q(u) &= \int_{\mathbf{R}^3} G(u(x)) dx = \lambda \neq 0 \end{aligned}$$

under the following assumptions:

- (a) $F, G : \mathbf{R} \rightarrow \mathbf{R}$ are C^2 functions, $F(0) = F'(0) = 0$, $G(0) = G'(0) = 0$ and there exist $C > 0$, $\sigma < 5$ such that

$$|F'(u)| \leq C|u|^\sigma \quad \text{and} \quad |G'(u)| \leq C|u|^\sigma \quad \text{for } |u| \geq 1.$$

- (b) For any $\varepsilon > 0$, $G' \neq 0$ on $(-\varepsilon, 0)$ and on $(0, \varepsilon)$.

Then, after a translation, u is radially symmetric in the variables (x_2, x_3) (i.e. u is axially symmetric).

Proof. Making a translation in the x_2 direction if necessary, we may assume that

$$\int_{\{x_2 < 0\}} G(u(x)) dx = \int_{\{x_2 > 0\}} G(u(x)) dx = \frac{\lambda}{2}.$$

As before, we define u_1 and u_2 by

$$u_1(x_1, x_2, x_3) = \begin{cases} u(x_1, x_2, x_3) & \text{if } x_2 < 0, \\ u(x_1, -x_2, x_3) & \text{if } x_2 \geq 0, \end{cases} \quad u_2(x_1, x_2, x_3) = \begin{cases} u(x_1, -x_2, x_3) & \text{if } x_2 < 0, \\ u(x_1, x_2, x_3) & \text{if } x_2 \geq 0. \end{cases}$$

It is obvious that $Q(u_1) = Q(u_2) = \lambda$. Moreover, using (4.28) we get

$$\begin{aligned} E(u_1) + E(u_2) - 2E(u) &= -\frac{1}{4} \frac{1}{(2\pi)^3} \left[\int_{\mathbf{R}^3} \frac{\xi_1^2}{|\xi|^2} |\widehat{u}_1^2(\xi)|^2 d\xi + \int_{\mathbf{R}^3} \frac{\xi_1^2}{|\xi|^2} |\widehat{u}_2^2(\xi)|^2 d\xi \right. \\ &\quad \left. - 2 \int_{\mathbf{R}^3} \frac{\xi_1^2}{|\xi|^2} |\widehat{u}^2(\xi)|^2 d\xi \right]. \end{aligned} \quad (4.29)$$

Recall that by (2.40) and (2.41) we have the equality

$$\begin{aligned} &\int_{\mathbf{R}^N} \frac{\xi_j^2}{|\xi|^2} |\widehat{T_1\varphi}(\xi)|^2 d\xi + \int_{\mathbf{R}^N} \frac{\xi_j^2}{|\xi|^2} |\widehat{T_2\varphi}(\xi)|^2 d\xi - 2 \int_{\mathbf{R}^N} \frac{\xi_j^2}{|\xi|^2} |\widehat{\varphi}(\xi)|^2 d\xi \\ &= \frac{8}{\pi} \int_{\mathbf{R}^{N-1}} \frac{\xi_j^2}{|\xi'|} \left| \int_0^\infty \widehat{A\varphi}(\xi_1, \xi') \frac{\xi_1}{\xi_1^2 + |\xi'|^2} d\xi_1 \right|^2 d\xi' \end{aligned} \quad (4.30)$$

for any $\varphi \in C_c^\infty(\mathbf{R}^N)$, where $j \in \{2, \dots, N\}$. It is obvious that the left-hand side of (4.30) defines a continuous functional on $L^2(\mathbf{R}^N)$. By the next lemma, it follows that the right-hand side of (4.30) also defines a continuous functional on $L^2(\mathbf{R}^N)$. Then the density of $C_c^\infty(\mathbf{R}^N)$ in $L^2(\mathbf{R}^N)$ implies that (4.30) holds for any $\varphi \in L^2(\mathbf{R}^N)$.

Lemma 4.10. *Let $j \in \{2, \dots, N\}$. The bilinear form*

$$S_1(\varphi, \psi) = \int_{\mathbf{R}^{N-1}} \frac{\xi_j^2}{|\xi'|} \int_0^\infty \widehat{\varphi}(\xi_1, \xi') \frac{\xi_1}{\xi_1^2 + |\xi'|^2} d\xi_1 \cdot \int_0^\infty \overline{\widehat{\psi}(\eta_1, \xi')} \frac{\eta_1}{\eta_1^2 + |\xi'|^2} d\eta_1 d\xi'$$

is continuous on $L^2(\mathbf{R}^N) \times L^2(\mathbf{R}^N)$.

Proof. As in (4.13) we have

$$\left| \int_0^\infty \widehat{\varphi}(\xi_1, \xi') \frac{\xi_1}{\xi_1^2 + |\xi'|^2} d\xi_1 \right| \leq K \frac{1}{|\xi'|^{\frac{1}{2}}} \left(\int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^2 d\xi_1 \right)^{\frac{1}{2}},$$

where $K = (\int_0^\infty \frac{t^2}{(1+t^2)^2} dt)^{\frac{1}{2}}$. Consequently

$$\begin{aligned}
|S_1(\varphi, \psi)| &\leq K^2 \int_{\mathbf{R}^{N-1}} \frac{\xi_j^2}{|\xi'|^2} \left(\int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^2 d\xi_1 \right)^{\frac{1}{2}} \left(\int_0^\infty |\widehat{\psi}(\eta_1, \xi')|^2 d\eta_1 \right)^{\frac{1}{2}} d\xi' \\
&\leq K^2 \int_{\mathbf{R}^{N-1}} \left(\int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^2 d\xi_1 \right)^{\frac{1}{2}} \left(\int_0^\infty |\widehat{\psi}(\eta_1, \xi')|^2 d\eta_1 \right)^{\frac{1}{2}} d\xi' \\
&\leq K^2 \left(\int_{\mathbf{R}^{N-1}} \int_0^\infty |\widehat{\varphi}(\xi_1, \xi')|^2 d\xi_1 d\xi' \right)^{\frac{1}{2}} \cdot \left(\int_{\mathbf{R}^{N-1}} \int_0^\infty |\widehat{\psi}(\eta_1, \xi')|^2 d\eta_1 d\xi' \right)^{\frac{1}{2}} \\
&\leq K_1 \|\varphi\|_{L^2(\mathbf{R}^N)} \|\psi\|_{L^2(\mathbf{R}^N)}. \quad \square
\end{aligned}$$

Since $u^2, u_1^2, u_2^2 \in L^2(\mathbf{R}^3)$ (recall that $H^1(\mathbf{R}^3) \subset L^2(\mathbf{R}^3) \cap L^6(\mathbf{R}^3)$), by exchanging the roles of x_1 and x_2 and using (4.29) and (4.30) we find

$$\begin{aligned}
&E(u_1) + E(u_2) - 2E(u) \\
&= -\frac{1}{4} \frac{1}{(2\pi)^3} \frac{8}{\pi} \int_{\mathbf{R}^2} \frac{\xi_1^2}{\sqrt{\xi_1^2 + \xi_3^2}} \left| \int_0^\infty \widehat{A_2(u^2)}(\xi_1, \xi_2, \xi_3) \frac{\xi_2}{\xi_1^2 + \xi_2^2 + \xi_3^2} d\xi_2 \right|^2 d\xi_1 d\xi_3, \quad (4.31)
\end{aligned}$$

where $A_2\varphi = \frac{1}{2}(\varphi(x_1, x_2, x_3) - \varphi(x_1, -x_2, x_3))$.

Since u is a minimizer, we must have $E(u_1) + E(u_2) - 2E(u) \geq 0$, consequently the integral in the right-hand side of (4.31) must be zero, which is equivalent to

$$\int_0^\infty \widehat{A_2(u^2)}(\xi_1, \xi_2, \xi_3) \frac{\xi_2}{\xi_1^2 + \xi_2^2 + \xi_3^2} d\xi_2 = 0 \quad \text{a.e. } (\xi_1, \xi_3) \in \mathbf{R}^2. \quad (4.32)$$

In particular, u_1 and u_2 are also minimizers. However, as in the previous example, (4.32) is not sufficient to prove that $A_2(u^2) = 0$. In order to accomplish this task, we will use the Euler-Lagrange equation of u : since u minimizes E under the constraint $Q(u) = \lambda$, there exists a constant α such that $E'(u) + \alpha Q'(u) = 0$, that is

$$-\Delta u + F'(u) + R_1^2(u^2)u + \alpha G'(u) = 0. \quad (4.33)$$

Lemma 4.11. *If F and G satisfy assumption (a) in Theorem 4.9 and $u \in H^1(\mathbf{R}^3)$ is a solution of (4.33), then $u \in W^{3,p}(\mathbf{R}^3)$ for any $p \in [2, \infty)$. In particular, $u \in C^2(\mathbf{R}^3)$.*

Since R_1 and R_1^2 are linear continuous mappings from $L^p(\mathbf{R}^3)$ to $L^p(\mathbf{R}^3)$ for $1 < p < \infty$, the proof of Lemma 4.11 is standard, so we omit it.

Let $I(\varphi)(x) = \int_{\mathbf{R}^3} \frac{\varphi(y)}{|x-y|} dy$. Using Lemma 4.5 it is easy to see that $I(u^2) \in W^{2,p}(\mathbf{R}^3)$ for any $p \in (3, \infty]$ and $I(u^2)$ is a C^2 function. Moreover, we have

$$\mathcal{F}(R_1^2(u^2))(\xi) = -\frac{\xi_1^2}{|\xi|^2} \widehat{u^2}(\xi) = -\frac{1}{d_3} \xi_1^2 \widehat{I(u^2)}(\xi),$$

where $d_3 = \frac{4\pi^{\frac{3}{2}}}{\Gamma(\frac{1}{2})}$, thus $R_1^2(u^2) = \frac{1}{d_3} \frac{\partial^2}{\partial x_1^2} I(u^2)$. Equation (4.33) can be written as

$$-\Delta u + F'(u) + \frac{1}{d_3} \frac{\partial^2}{\partial x_1^2} (I(u^2))u + \alpha G'(u) = 0. \quad (4.34)$$

Arguing exactly as in the proof of Theorem 4.6, (4.32) implies that $\frac{\partial}{\partial x_2} (I(u^2))(x_1, 0, x_3) = 0$ for any $(x_1, x_3) \in \mathbf{R}^2$.

Since u_1 is also a minimizer, it satisfies the Euler–Lagrange equation

$$-\Delta u_1 + F'(u_1) + \frac{1}{d_3} \frac{\partial^2}{\partial x_1^2} (I(u_1^2))u_1 + \beta G'(u_1) = 0. \quad (4.35)$$

The conclusion of Lemma 4.11 is obviously valid for u_1 . Since u_1 is symmetric with respect to x_2 , $I(u_1^2)$ is also symmetric with respect to x_2 and, consequently, $\frac{\partial}{\partial x_2} (I(u_1^2))(x_1, 0, x_3) = 0$ for any $(x_1, x_3) \in \mathbf{R}^2$. We set $U = I(u^2)$ and $U_1 = I(u_1^2)$. Recall that $u(x_1, x_2, x_3) = u_1(x_1, x_2, x_3)$ if $x_2 < 0$; thus U and U_1 are both solutions of

$$\begin{cases} -\Delta W = u^2 & \text{in } \mathbf{R} \times (-\infty, 0) \times \mathbf{R}, \\ W \in C^2(\mathbf{R}^3) \cap W^{2,p}(\mathbf{R}^3) & \text{for } 3 < p \leq \infty, \\ \frac{\partial W}{\partial x_2}(x_1, 0, x_3) = 0 & \text{for any } (x_1, x_3) \in \mathbf{R}^2. \end{cases} \quad (4.36)$$

It is not hard to see that the solution of (4.36) is unique. Hence we must have $I(u^2) = I(u_1^2)$ in $\mathbf{R} \times (-\infty, 0] \times \mathbf{R}$. In the same way we obtain $I(u^2) = I(u_1^2)$ in $\mathbf{R} \times [0, \infty) \times \mathbf{R}$.

Now we focus our attention on u_1 . Making a translation in the x_3 direction if necessary, we may assume that $\int_{\{x_3 < 0\}} G(u_1(x)) dx = \int_{\{x_3 > 0\}} G(u_1(x)) dx = \frac{\lambda}{2}$. We define

$$\begin{aligned} w_1(x_1, x_2, x_3) &= \begin{cases} u_1(x_1, x_2, x_3) & \text{if } x_3 < 0, \\ u_1(x_1, x_2, -x_3) & \text{if } x_3 \geq 0, \end{cases} \\ w_2(x_1, x_2, x_3) &= \begin{cases} u_1(x_1, x_2, -x_3) & \text{if } x_3 < 0, \\ u_1(x_1, x_2, x_3) & \text{if } x_3 \geq 0. \end{cases} \end{aligned}$$

It is obvious that $Q(w_1) = Q(w_2) = \lambda$. Proceeding as above, we find the identity

$$\begin{aligned} E(w_1) + E(w_2) - 2E(u_1) \\ = -\frac{1}{4} \frac{1}{(2\pi)^3} \frac{8}{\pi} \int_{\mathbf{R}^2} \frac{\xi_1^2}{\sqrt{\xi_1^2 + \xi_2^2}} \left| \int_0^\infty \widehat{A_3(u_1^2)}(\xi_1, \xi_2, \xi_3) \frac{\xi_3}{\xi_1^2 + \xi_2^2 + \xi_3^2} d\xi_3 \right|^2 d\xi_1 d\xi_2, \end{aligned} \quad (4.37)$$

where $A_3\varphi = \frac{1}{2}(\varphi(x_1, x_2, x_3) - \varphi(x_1, x_2, -x_3))$. Since u_1 is a minimizer, it follows from (4.37) that w_1 and w_2 are also minimizers of E under the constraint $Q = \lambda$; hence w_1 and w_2 satisfy the conclusion of Lemma 4.11 and $I(w_1), I(w_2) \in C^2(\mathbf{R}^3) \cap W^{2,p}(\mathbf{R}^3)$ for $p \in (3, \infty]$. Moreover, the integral in the right-hand side of (4.37) must be zero. As previously, this gives

$\frac{\partial}{\partial x_3} I(u_1^2)(x_1, x_2, 0) = 0$ for any $(x_1, x_2) \in \mathbf{R}^2$. Proceeding as above, we find $I(u_1^2) = I(w_1^2)$ in $\mathbf{R}^2 \times (-\infty, 0]$ and $I(u_1^2) = I(w_2^2)$ in $\mathbf{R}^2 \times [0, \infty)$.

Now let us consider the function w_1 . It is clear that $w_1(x_1, -x_2, -x_3) = w_1(x_1, -x_2, x_3) = w_1(x_1, x_2, x_3)$, i.e. w_1 is symmetric with respect to x_2 and with respect to x_3 . Consider a plane Π in $\mathbf{R}^3$ containing the line $\{(x_1, 0, 0) \mid x_1 \in \mathbf{R}\}$ and let Π_+ and Π_- be the two half-spaces determined by Π . Since $(x_1, x_2, x_3) \mapsto (x_1, -x_2, -x_3)$ maps Π_+ onto Π_- , using the symmetry of w_1 we get $\int_{\Pi_+} G(w_1(x)) dx = \int_{\Pi_-} G(w_1(x)) dx = \frac{\lambda}{2}$. Let s_Π denote the symmetry in $\mathbf{R}^3$ with respect to Π . We define

$$r_1(x) = \begin{cases} w_1(x) & \text{if } x \in \Pi_-, \\ w_1(s_\Pi(x)) & \text{if } x \in \Pi_+ \end{cases} \quad \text{and} \quad r_2(x) = \begin{cases} w_1(s_\Pi(x)) & \text{if } x \in \Pi_-, \\ w_1(x) & \text{if } x \in \Pi_+. \end{cases}$$

Repeating the above arguments we obtain an integral identity analogous to (4.31) and (4.37) which implies that r_1 and r_2 also minimize E subject to the constraint $Q = \lambda$. Furthermore, using the fact that the integral in the right-hand side of this identity must vanish we find

$$\frac{\partial}{\partial n} I(w_1^2)(x_1, x_2, x_3) = 0 \quad \text{whenever } (x_1, x_2, x_3) \in \Pi, \quad (4.38)$$

where n is the unit normal to Π . Passing to cylindrical coordinates we write

$$I(w_1^2)(x_1, x_2, x_3) = I(w_1^2)(x_1, r \cos \theta, r \sin \theta) = \Phi(x_1, r, \theta),$$

where $r = \sqrt{x_2^2 + x_3^2}$. Since $I(w_1^2)$ is a C^2 function and (4.38) is valid for any plane Π containing $\{(x_1, 0, 0) \mid x_1 \in \mathbf{R}\}$, (4.38) is equivalent to $\frac{\partial \Phi}{\partial \theta} = 0$, that is Φ does not depend on θ , i.e. $I(w_1^2)(x_1, x_2, x_3) = \Phi_1(x_1, \sqrt{x_2^2 + x_3^2})$ for some function Φ_1 . In other words, we have proved that $I(w_1^2)$ is radially symmetric in the variables (x_2, x_3) . In the same way we show that $I(w_2^2)(x_1, x_2, x_3) = \Phi_2(x_1, \sqrt{x_2^2 + x_3^2})$ for some function Φ_2 . Since $I(u_1^2)$ is continuous on $\mathbf{R}^3$, $I(u_1^2) = I(w_1^2)$ in the half-space $\{x_3 < 0\}$ and $I(u_1^2) = I(w_2^2)$ in the half-space $\{x_3 > 0\}$, we have necessarily $\Phi_1 = \Phi_2$, and then $I(u_1^2)$ is radially symmetric in the variables (x_2, x_3) .

Similarly, there exists $k \in \mathbf{R}$ such that $\int_{\{x_3 < k\}} G(u_2(x)) dx = \int_{\{x_3 > k\}} G(u_2(x)) dx = \frac{\lambda}{2}$. (We have fixed the origin in such a way that $\int_{\{x_3 < 0\}} G(u_1(x)) dx = \int_{\{x_3 > 0\}} G(u_1(x)) dx = \frac{\lambda}{2}$ and nothing guarantees a priori that $k = 0$.) Arguing as above, we infer that $I(u_2^2)$ is radially symmetric with respect to the variables $(x_2, x_3 - k)$. Thus we have proved that there exist continuous functions η, γ defined on $\mathbf{R} \times [0, \infty)$ such that $I(u_1^2)(x_1, x_2, x_3) = \eta(x_1, \sqrt{x_2^2 + x_3^2})$ and $I(u_2^2)(x_1, x_2, x_3) = \gamma(x_1, \sqrt{x_2^2 + (x_3 - k)^2})$. Since $I(u_1^2)(x_1, 0, x_3) = I(u_2^2)(x_1, 0, x_3) = I(u_2^2)(x_1, 0, x_3)$, we get $\eta(x_1, |x_3|) = \gamma(x_1, |x_3 - k|)$ for any $x_1, x_3 \in \mathbf{R}$. In particular, if $k \geq 0$, for $t \geq 0$ we have $\eta(x_1, t + 2k) = \gamma(x_1, t + k) = \eta(x_1, t)$; that is, for any fixed x_1 , the function $\eta(x_1, \cdot)$ is periodic of period $2k$. On the other hand, we have $I(u_1^2), I(u_2^2) \in W^{2,p}(\mathbf{R}^N)$ for $p \in (3, \infty]$, thus $I(u_1^2)$ and $I(u_2^2)$ tend to zero at infinity, hence $\eta(x_1, t) \rightarrow 0$ and $\gamma(x_1, t) \rightarrow 0$ as $x_1^2 + t^2 \rightarrow \infty$. We infer that either $k = 0$, or $\eta \equiv 0$ in $\mathbf{R} \times [0, \infty)$. In both cases we get $\eta = \gamma$ on $\mathbf{R} \times [0, \infty)$ and $I(u_1^2) = I(u_2^2)$ in $\mathbf{R}^3$. Thus we have $I(u^2) = I(u_1^2) = I(u_2^2)$ on $\mathbf{R}^3$, and $I(u^2)$ is radially symmetric with respect to (x_2, x_3) .

Since $Q(u) = Q(u_1) = \lambda \neq 0$, we cannot have $u \equiv 0$ in the half-space $\{x_2 < 0\}$. Assumption (b) implies that there exists $(x_1, x_2, x_3) \in \mathbf{R}^3$, $x_2 < 0$ such that $G'(u(x_1, x_2, x_3)) \neq 0$. Since $u = u_1$ on $\{x_2 < 0\}$ and $I(u^2) = I(u_1^2)$ on $\mathbf{R}^3$, from (4.34) and (4.35) we infer that $\alpha = \beta$. Let $a(x) = \frac{1}{d_3} \frac{\partial^2}{\partial x_1^2} (I(u^2))(x) = \frac{1}{d_3} \frac{\partial^2}{\partial x_1^2} (I(u_1^2))(x)$. We know that a is a continuous and bounded function on $\mathbf{R}^3$. The functions u and u_1 both satisfy the equation $-\Delta w + F'(w) + a(x)w + \alpha G'(w) = 0$ in $\mathbf{R}^3$ and using the Unique Continuation Principle again we conclude that $u \equiv u_1$ in $\mathbf{R}^3$, i.e. u is symmetric with respect to x_2 .

In the same way we prove that u is symmetric with respect to x_3 (after possibly a translation). Proceeding as in the proof of Theorem 4.1 we can show that u is symmetric with respect to any plane containing the line $\{(x_1, 0, 0) \mid x_1 \in \mathbf{R}\}$, consequently u is radially symmetric with respect to (x_2, x_3) variables. $\square$

Remark 4.12. (i) We have stated and proved Theorem 4.9 in dimension $N = 3$ only for simplicity. Replacing the term $\int_{\mathbf{R}^3} |R_1(u^2)|^2(x) dx$ in $E(u)$ by $\int_{\mathbf{R}^N} |R_1(H(u))|^2(x) dx$ and making suitable assumptions on the function H , this result admits a straightforward generalization to $\mathbf{R}^N$, $N \geq 3$.

(ii) We do not know whether the minimizers in Theorem 4.9 are symmetric or not with respect to x_1 . Recall that by (2.42) we have

$$\begin{aligned} & \int_{\mathbf{R}^N} \frac{\xi_1^2}{|\xi|^2} |\widehat{T_1 \varphi}(\xi)|^2 d\xi + \int_{\mathbf{R}^N} \frac{\xi_1^2}{|\xi|^2} |\widehat{T_2 \varphi}(\xi)|^2 d\xi - 2 \int_{\mathbf{R}^N} \frac{\xi_1^2}{|\xi|^2} |\widehat{\varphi}(\xi)|^2 d\xi \\ &= -\frac{8}{\pi} \int_{\mathbf{R}^{N-1}} |\xi'| \left| \int_0^\infty \widehat{A \varphi}(\xi) \frac{\xi_1}{\xi_1^2 + |\xi'|^2} d\xi_1 \right|^2 d\xi' \end{aligned} \quad (4.39)$$

for any $\varphi \in C_c^\infty(\mathbf{R}^N)$. Clearly, the left-hand side of (4.39) is continuous on $L^2(\mathbf{R}^N)$. Proceeding as in Lemma 4.10, it is easy to see that the right-hand side of (4.39) also defines a continuous functional on $L^2(\mathbf{R}^N)$. Consequently, (4.39) holds for any $\varphi \in L^2(\mathbf{R}^N)$. Using (4.28) and (4.39) we have

$$E(T_1 u) + E(T_2 u) - 2E(u) = \frac{2}{\pi} \frac{1}{(2\pi)^N} \int_{\mathbf{R}^{N-1}} |\xi'| \left| \int_0^\infty \mathcal{F}(A(H(u)))(\xi) \frac{\xi_1}{|\xi|^2} d\xi_1 \right|^2 d\xi'. \quad (4.40)$$

The right-hand side in this integral identity is always nonnegative and (4.40) does not imply the symmetry of minimizers with respect to x_1 .

(iii) Let us change the sign of the nonlocal term appearing in Theorem 4.9, i.e. let us consider the minimization problem

$$\begin{aligned} \text{minimize } E_*(u) &= \frac{1}{2} \int_{\mathbf{R}^3} |\nabla u|^2 dx + \int_{\mathbf{R}^3} F(u) dx + \frac{1}{4} \int_{\mathbf{R}^3} |R_1(u^2)|^2 dx \\ \text{under the constraint } Q(u) &:= \int_{\mathbf{R}^3} G(u(x)) dx = \lambda. \end{aligned} \quad (4.41)$$

The minimizers of this problem (when they exist) give rise to standing waves for Eq. (4.27) where the sign of the nonlocal term $R_1^2(|u|^2)u$ has been reversed. Clearly, the integral identities that we have do not imply the symmetry of solutions of (4.41) with respect to x_2 and x_3 .

The symmetry of minimizers of (4.41) with respect to x_1 is also an open problem. As above, in this case we have the identity

$$E_*(T_1u) + E_*(T_2u) - 2E_*(u) = -\frac{2}{\pi} \frac{1}{(2\pi)^3} \int_{\mathbf{R}^2} |\xi'| \left| \int_0^\infty \mathcal{F}(A(u^2))(\xi) \frac{\xi_1}{|\xi|^2} d\xi_1 \right|^2 d\xi_2 d\xi_3. \quad (4.42)$$

If u is a minimizer, the right-hand side of (4.42) must vanish. As in the proof of Theorem 4.9, this implies $\frac{\partial}{\partial x_1} I(u^2)(0, x_2, x_3) = 0$ for any $(x_2, x_3) \in \mathbf{R}^2$. Repeating the argument already used in Theorem 4.9 we get $I(u^2) = I((T_1u)^2)$ on $\{x_1 \leq 0\}$ and $I(u^2) = I((T_2u)^2)$ on $\{x_1 \geq 0\}$. Moreover, if $\lambda \neq 0$ then u and $u_1 := T_1u$ satisfy the same Euler–Lagrange equation, namely

$$-\Delta w + F'(w) - \frac{1}{d_3} \frac{\partial^2}{\partial x_1^2} (I(w^2))w + \alpha G'(w) = 0. \quad (4.43)$$

Equivalently, defining $U = I(u^2)$ and $U_1 = I(u_1^2)$, we see that (u, U) and (u_1, U_1) are both solutions to the system

$$\begin{cases} -\Delta w + F'(w) - \frac{1}{d_3} \frac{\partial^2 W}{\partial x_1^2} w + \alpha G'(w) = 0, \\ -\Delta W = w^2. \end{cases} \quad (4.44)$$

Moreover, $(u, U) = (u_1, U_1)$ on $\{x_1 \leq 0\}$ and u, u_1 satisfy the conclusion of Lemma 4.11. We do not know whether this information together with the boundary condition $\frac{\partial U}{\partial x_1}(0, x_2, x_3) = \frac{\partial U_1}{\partial x_1}(0, x_2, x_3) = 0$ imply that $u \equiv u_1$.

Remark 4.13. If $N = 3$, the nonlocal term in Theorem 4.9 can be written as

$$\begin{aligned} \int_{\mathbf{R}^3} |R_1(u^2)|^2 dx &= \frac{1}{(2\pi)^3} \int_{\mathbf{R}^3} \frac{\xi_1^2}{|\xi|^2} |\widehat{u^2}(\xi)|^2 d\xi = -\frac{1}{d_3(2\pi)^3} \int_{\mathbf{R}^3} \mathcal{F}\left(\frac{\partial^2}{\partial x_1^2} I(u^2)\right)(\xi) \overline{\widehat{u^2}(\xi)} d\xi \\ &= -\frac{1}{d_3} \int_{\mathbf{R}^3} \frac{\partial^2}{\partial x_1^2} I(u^2)(x) \overline{u^2(x)} dx = -\frac{1}{d_3} \int_{\mathbf{R}^3} \int_{\mathbf{R}^3} u^2(x) K(x-y) u^2(y) dx dy, \end{aligned}$$

where $K(x) = \frac{\partial^2}{\partial x_1^2} \left(\frac{1}{|x|}\right) = \frac{2x_1^2 - x_2^2 - x_3^2}{(x_1^2 + x_2^2 + x_3^2)^{\frac{5}{2}}}$. Since this kernel changes sign, spherical rearrangements in the variables (x_2, x_3) combined with Riesz' inequality cannot be used to prove the symmetry of minimizers.

5. Some open problems

We close this paper speaking about several problems for which the methods described above (including ours) seem to fail.

First, let us come back to the two minimization problems considered in Theorem 4.1. As before, if u is a minimizer of any of these problems, we may assume that $\int_{\{x_1 < 0\}} G(u) dx = \int_{\{x_1 > 0\}} G(u) dx$ and we set $u_1 = T_1 u$ and $u_2 = T_2 u$. Assume that $s \in (1, \frac{3}{2})$. Then the identities (3.22) and (3.23) are still valid (see Corollary 3.4) and we get

$$E(u_1) + E(u_2) - 2E(u) = -\frac{16 \sin(s\pi)}{\pi^2} N_s^2(Au) \geq 0 \quad \text{in Case A,}$$

respectively

$$E(u_1) + E(u_2) - 2E(u) = -\frac{16 \sin(s\pi)}{\pi^2} \tilde{N}_s^2(Au) \geq 0 \quad \text{in Case B.}$$

It is easy to see that these integral identities work in the wrong direction. Are the minimizers still radially symmetric for $s \in (1, \frac{3}{2})$?

Another problem is to study the symmetry of minimizers of

$$E(u) = \frac{1}{2} \int_{\mathbf{R}^3} |\nabla u|^2 + \int_{\mathbf{R}^3 \times \mathbf{R}^3} \frac{1}{|x - y|} u(x)^2 u(y)^2 dx dy + \int_{\mathbf{R}^3} F(u(x)) dx$$

subject to the constraint

$$\int_{\mathbf{R}^3} u^2(x) dx = \lambda > 0.$$

In the particular case $F(u) = -C|u|^{8/3}$, this problem arises in connection with the Schrödinger–Poisson–Slater system [22]. Due to the repulsive effect of the nonlocal term, Riesz’ inequality as well as the Reflection method work in the wrong direction.

The last problem concerns the symmetry of minimizers of

$$E(u) = \int_{-\infty}^{+\infty} (u_x^2(x) + u^3(x)) dx - \gamma \int_{-\infty}^{+\infty} |\xi| |\widehat{u}(\xi)|^2 d\xi,$$

where $\gamma > 0$, subject to the constraint $\int_{-\infty}^{+\infty} u^2(x) dx = \lambda > 0$. These two functionals are conserved quantities for the Benjamin equation (see [1,2]). Symmetrization and reflection cannot be used due to the sign of the nonlocal term. Oscillating travelling waves for this equation have been found numerically; perhaps this is an indication that the minimizers of the problem above may change the sign.

References

- [1] J. Albert, J.L. Bona, J.-C. Saut, Model equations for waves in stratified fluids, *Proc. R. Soc. London Ser. A* 453 (1997) 1233–1260.
- [2] J. Albert, J.L. Bona, M. Restrepo, Solitary wave solutions of the Benjamin equation, *SIAM J. Appl. Math.* 59 (6) (1999) 2139–2161.
- [3] J. Angulo, F. Montenegro, Existence and evenness of solitary wave-solutions for an equation of short and long dispersive waves, *Nonlinearity* 13 (5) (2000) 1595–1611.
- [4] A. de Bouard, Stability and instability of some nonlinear dispersive solitary waves in higher dimensions, *Proc. Roy. Soc. Edinburgh Sect. A* 126 (1996) 89–112.
- [5] F. Brock, A general rearrangement inequality à la Hardy–Littlewood, *J. Inequal. Appl.* 5 (4) (2000) 309–320.
- [6] J. Brothers, W. Ziemer, Minimal rearrangements of Sobolev functions, *J. Reine Angew. Math.* 384 (1988) 153–179.
- [7] J. Busca, B. Sirakov, Symmetry results for semilinear elliptic systems in the whole space, *J. Differential Equations* 163 (1) (2000) 41–56.
- [8] H. Cartan, *Théorie élémentaire des fonctions analytiques d’une ou plusieurs variables complexes*, Hermann, Paris, 1961.
- [9] T. Cazenave, P.-L. Lions, Orbital stability of standing waves for some nonlinear Schrödinger equations, *Comm. Math. Phys.* 85 (1982) 549–561.
- [10] R. Cipelatti, On the existence of standing waves for a Davey–Stewartson system, *Comm. Partial Differential Equations* 17 (5–6) (1992) 967–988.
- [11] B. Gidas, W. Ni, L. Nirenberg, Symmetry of positive solutions of nonlinear elliptic equations in $\mathbf{R}^N$, in: *Adv. Math. Suppl. Stud.*, vol. 7A, Academic Press, 1981, pp. 369–402.
- [12] B. Guo, W. Baoxiang, The Cauchy problem for Davey–Stewartson systems, *Comm. Pure Appl. Math.* 52 (1999) 1477–1490.
- [13] L. Hörmander, *Linear Partial Differential Operators*, Springer-Verlag, New York, 1969.
- [14] B. Kawohl, Symmetry or not?, *Math. Intelligencer* 20 (1998) 16–22.
- [15] E. Lieb, Existence and uniqueness of the minimizing solution of Choquard’s nonlinear equation, *Studies Appl. Math.* 57 (2) (1977) 93–106.
- [16] E. Lieb, M. Loss, *Analysis*, Graduate Stud. Math., vol. 14, Amer. Math. Soc., Providence, RI, 1996.
- [17] P.-L. Lions, The Concentration–compactness principle in the calculus of variations, Part I, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 1 (1984) 109–145; Part II, 223–283.
- [18] O. Lopes, Radial symmetry of minimizers for some translation and rotation invariant functionals, *J. Differential Equations* 124 (1996) 378–388.
- [19] O. Lopes, Radial and nonradial minimizers for some radially symmetric functionals, *Electron. J. Differential Equations* 3 (1996) 1–14.
- [20] O. Lopes, Nonlocal variational problems arising in long wave propagation, *ESAIM Control Optim. Calc. Var.* 5 (2000) 501–528.
- [21] M. Mariş, On the existence, regularity and decay of solitary waves to a generalized Benjamin–Ono equation, *Nonlinear Anal.* 51 (2002) 1073–1085.
- [22] O. Sanchez, J. Soler, Long-time dynamics of the Schrödinger–Poisson–Slater system, *J. Stat. Phys.* 114 (1–2) (2004) 179–204.
- [23] E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Univ. Press, Princeton, NJ, 1970.
- [24] E.M. Stein, G. Weiss, *Fourier Analysis on Euclidean Spaces*, Princeton Univ. Press, Princeton, NJ, 1971.
- [25] W. Troy, Symmetry properties in systems of semilinear elliptic equations, *J. Differential Equations* 42 (3) (1981) 400–413.
- [26] M. Willem, *Minimax Theorems*, Progr. Nonlinear Differential Equations Appl., vol. 24, Birkhäuser, Basel, 1996.